

Forming localized dust concentrations in a dust ring

DM Tau case study

Hauyu Baobab Liu¹

Department of Physics, National Sun Yat-Sen University, No. 70, Lien-Hai Road, Kaohsiung City 80424, Taiwan, R.O.C.
e-mail: hyliau.nsysu@mail.nsysu.edu.tw

Received November 25, 2025; accepted November 26, 2025

ABSTRACT

Context. Previous high-angular-resolution 225 GHz (~ 1.3 mm) continuum observations of the transitional disk DM Tau have resolved an outer ring at 20–120 au radii that is weakly azimuthally asymmetric.

Aims. We aim to examine dust growth and filtration in the outer ring of DM Tau.

Methods. We performed $\sim 0''.06$ (~ 8.7 au) resolution Karl G. Jansky Very Large Array (JVLA) 40–48 GHz (~ 7 mm; Q band) continuum observations, along with complementary observations at lower frequencies. In addition, we analyzed the archival JVLA observations undertaken since 2010.

Results. Intriguingly, the Q band image resolved the azimuthally highly asymmetric, knotty dust emission sources close to the inner edge of the outer ring. Fitting the 8–700 GHz spectral energy distribution (SED) with two dust components indicates that the maximum grain size (a_{max}) in these knotty dust emission sources is likely $\gtrsim 300 \mu\text{m}$, whereas it is $\lesssim 50 \mu\text{m}$ in the rest of the ring. These results may be explained by a trapping of inwardly migrating "grown" dust close to the ring inner edge. The exact mechanism for developing the azimuthal asymmetry has not yet been identified, which may be due to planet-disk interaction that might also be responsible for the creation of the dust cavity and pressure bump. Otherwise, it may be due to the fluid instabilities and vortex formation as a result of shear motions. Finally, we remark that the asymmetries in DM Tau are difficult to diagnose from the $\gtrsim 225$ GHz observations, owing to a high optical depth at the ring. In other words, the apparent symmetric or asymmetric morphology of the transitional disks may be related to the optical depths of those disks at the observing frequency.

Key words. Protoplanetary disks – Planets and satellites: formation – (ISM:) dust, extinction – Radio continuum: ISM

1. Introduction

2. Data and reduction

2.1. Data

2.2. JVLA data processing

3. Results

4. Discussion

4.1. Spectral index distribution and model

4.2. Physical implications

4.2.1. Asymmetry in protoplanetary disks

4.2.2. Dust growth in an initially smooth background disk

5. Conclusion

Acknowledgements. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. This paper makes use of the following ALMA data: ADS/JAO.ALMA #2013.1.00498.S, #2013.1.00647.S, #2015.1.00296.S, #2016.1.00565.S, #2016.1.01042.S, #2017.1.01460.S, #2018.1.01755.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. The Submillimeter Array is a joint project between the Smithsonian Astrophysical Observatory and the Academia Sinica Institute of Astronomy and Astrophysics,

and is funded by the Smithsonian Institution and the Academia Sinica (Ho et al. 2004). H.B.L. is supported by the National Science and Technology Council (NSTC) of Taiwan (Grant Nos. 113-2112-M-110-022-MY3).

References

Ho, P. T. P., Moran, J. M., & Lo, K. Y. 2004, ApJ, 616, L1