

# 科技寫作(Scientific Writing)



Week 11 (Dec. 08 – Dec. 12)

Introduction Section

# 期末報告Requirement

1. 書面格式: (PDF) [RevTeX 4.2 Template and Sample](#)

2. 內容 :

(i) PDF-1 : Title + Abstract

(ii) PDF-2 += PDF-1 + all (Sub)section Titles

(iii) PDF-3 += First-sentences of all paragraphs in Method Section

(iv) PDF-4 += PDF-3 + At least 2 Figures for experimental results (with captions and some First-sentences in the Result section.

(v) PDF-5 += PDF-4 + First-sentences of all paragraphs in at least one Discussion subsection.

# 期末報告Requirement (口頭15mins-年會口頭報告長度)

1. 書面格式: (PDF) [RevTeX 4.2 Template and Sample](#)

2. 內容 :

(i) PDF-1 : Title + Abstract

(ii) PDF-2 += PDF-1 + all (Sub)section Titles

口頭: 3 mins概述研究(1)背景、(2)具體問題、(3)研究方法邏輯、  
(4)預期得到結果、(5)結果之預期physical implication

(iii) PDF-3 += First-sentences of all paragraphs in Method Section

(iv) PDF-4 += PDF-3 + At least 2 Figures for experimental results (with captions  
and some First-sentences in the Result section.

(v) PDF-5 += PDF-4 + First-sentences of all paragraphs in at least one Discussion subsection.

# 期末報告Requirement (口頭15mins-年會口頭報告長度)

1. 書面格式: (PDF) [RevTeX 4.2 Template and Sample](#)

2. 內容 :

(i) PDF-1 : Title + Abstract

(ii) PDF-2 += PDF-1 + all (Sub)section Titles

口頭: 3 mins概述研究(1)背景、(2)具體問題、(3)研究方法邏輯、  
(4)預期得到結果、(5)結果之預期physical implication

(iii) PDF-3 += First-sentences of all paragraphs in Method Section

口頭: 3 mins詳述研究方法

(iv) PDF-4 += PDF-3 + At least 2 Figures for experimental results (with captions  
and some First-sentences in the Result section.

(v) PDF-5 += PDF-4 + First-sentences of all paragraphs in at least one Discussion subsection.

# 期末報告Requirement (口頭15mins-年會口頭報告長度)

1. 書面格式: (PDF) [RevTeX 4.2 Template and Sample](#)

2. 內容 :

(i) PDF-1 : Title + Abstract

(ii) PDF-2 += PDF-1 + all (Sub)section Titles

口頭: 3 mins概述研究(1)背景、(2)具體問題、(3)研究方法邏輯、  
(4)預期得到結果、(5)結果之預期physical implication

(iii) PDF-3 += First-sentences of all paragraphs in Method Section

口頭: 3 mins詳述研究方法

(iv) PDF-4 += PDF-3 + At least 2 Figures for experimental results (with captions  
and some First-sentences in the Result section.

口頭: 6 mins詳述初步研究結果(聚焦於圖片講解)

(v) PDF-5 += PDF-4 + First-sentences of all paragraphs in at least one Discussion subsection.

# 期末報告Requirement (口頭15mins-年會口頭報告長度)

1. 書面格式: (PDF) [RevTeX 4.2 Template and Sample](#)

2. 內容 :

(i) PDF-1 : Title + Abstract

(ii) PDF-2 += PDF-1 + all (Sub)section Titles

口頭: 3 mins 概述研究(1)背景、(2)具體問題、(3)研究方法邏輯、  
(4)預期得到結果、(5)結果之預期physical implication

(iii) PDF-3 += First-sentences of all paragraphs in Method Section

口頭: 3 mins 詳述研究方法

(iv) PDF-4 += PDF-3 + At least 2 Figures for experimental results (with captions  
and some First-sentences in the Result section.

口頭: 6 mins 詳述初步研究結果(聚焦於圖片講解)

(v) PDF-5 += PDF-4 + First-sentences of all paragraphs in at least one Discussion subsection.

口頭: 3 mins 略述physical implication

報告前e-mail繳交PDF-1~5 (檔名：姓名 ScientificWriting 1(or2345).pdf )、  
評分會考慮是否超時(15 mins)及遲(未)到  
若針對同學的報告提問可額外加分

1. 書面格式: (PDF) [RevTeX 4.2 Template and Sample](#)

2. 內容：

(i) PDF-1 : Title + Abstract

(ii) PDF-2 += PDF-1 + all (Sub)section Titles

□頭: 3 mins概述研究(1)背景、(2)具體問題、(3)研究方法邏輯、  
(4)預期得到結果、(5)結果之預期physical implication

(iii) PDF-3 += First-sentences of all paragraphs in Method Section

□頭: 3 mins詳述研究方法

(iv) PDF-4 += PDF-3 + At least 2 Figures for experimental results (with captions  
and some First-sentences in the Result section.

□頭: 6 mins詳述初步研究結果(聚焦於圖片講解)

(v) PDF-5 += PDF-4 + First-sentences of all paragraphs in at least one Discussion subsection.

□頭: 3 mins略述physical implication

報告前e-mail繳交PDF-1~5 (檔名：姓名 ScientificWriting 1(or2345).pdf )、  
評分會考慮是否超時(15 mins)及遲(未)到  
若針對同學的報告提問可額外加分



論文全文：

<https://ui.adsabs.harvard.edu/abs/2024A%26A...685A..18L/abstract>

1. 書面格式: (PDF) [RevTeX 4.2 Template and Sample](#)

2. 內容：

(i) PDF-1 : Title + Abstract

(ii) PDF-2 += PDF-1 + all (Sub)section Titles

□頭: 3 mins概述研究(1)背景、(2)具體問題、(3)研究方法邏輯、  
(4)預期得到結果、(5)結果之預期physical implication

(iii) PDF-3 += First-sentences of all paragraphs in Method Section

□頭: 3 mins詳述研究方法

(iv) PDF-4 += PDF-3 + At least 2 Figures for experimental results (with captions  
and some First-sentences in the Result section.

□頭: 6 mins詳述初步研究結果(聚焦於圖片講解)

(v) PDF-5 += PDF-4 + First-sentences of all paragraphs in at least one Discussion subsection.

□頭: 3 mins略述physical implication



# Introduction section

A nightmare for the supervisor. 😞

# Introduction section

## 1. Who are your readers: Experts

# Introduction section

## 1. Who are your readers: Experts

A professional journal paper is NOT a textbook. DO NOT try to educate your readers with the introduction section. (Exception: when the major purpose of your paper is to clarify a long-standing misunderstanding of something basic.)

# Introduction section

## 1. Who are your readers: Experts

A professional journal paper is NOT a textbook. DO NOT try to educate your readers with the introduction section. (Exception: when the major purpose of your paper is to clarify a long-standing misunderstanding of something basic.)

In and only in a thesis, there can be a literature review (文獻回顧).

# Introduction section

## 1. Who are your readers: Experts

A professional journal paper is NOT a textbook. DO NOT try to educate your readers with the introduction section. (Exception: when the major purpose of your paper is to clarify a long-standing misunderstanding of something basic.)

In and only in a thesis, there can be a literature review (文獻回顧).

Introduction =! 文獻回顧

# Introduction section

1. Who are your readers: Experts
2. What are the purposes of the introduction section?

# Introduction section

1. Who are your readers: Experts
2. What are the purposes of the introduction section?
  - (1) Appetizing
  - (2) Providing context and logic (instead of providing knowledge!)

# Introduction section

1. Who are your readers: Experts
2. What are the purposes of the introduction section?
  - (1) Appetizing
  - (2) Providing context and logic (instead of providing knowledge!)

Don't be verbose!  
Otherwise, people may give up reading before they  
get to the main text!  
(no longer than 1 two-column page. Never!)



論文全文：

<https://ui.adsabs.harvard.edu/abs/2024A%26A...685A..18L/abstract>



# Introduction section

1. Who are your readers: Experts
2. What are the purposes of the introduction section?
  - (1) Appetizing
  - (2) Providing context and logic (instead of providing knowledge!)
    - (i) background
    - (ii) question
    - (iii) logic + method
    - (iv) how this paper is organized (optional)

# 論文寫作實例

## Introduction

1. 電子具有波動性？  $\lambda = h/p$
2. 雙狹縫實驗

## Method

1. 用什麼device作為屏幕
2. 如何generate具特定動量的電子束


## Results

1. 是否有對應到特定波長的干涉條紋？
2. 電子束動量熱誤差及系統性偏差？

## Discussion

### 呼應問題

1. (假設)電子束中的電子無交互作用。若此假設不對，是否可能在不考慮平面波干涉的情形解釋干涉條紋？
2. 干涉條紋間距與光子雙狹縫干涉理論比較，驗證(1)電子的波動性，及(2)電子動量與波長間的關係
3. 進行單電子雙狹縫干涉實驗以排除電子交互作用造成類似干涉的條文的可能性。



# 論文寫作實例

## Introduction

**Context:** The blackbody spectrum [1], photoelectric effect [2], Compton scattering [3], and single/double-slit(s) [4] experiments have demonstrated the wave-particle duality of photons [see 5 for a pedagogical review], the quanta of electromagnetic waves. The question is, can all kinds of massive and massless particles (e.g., electrons) diffract and interfere like how photons do? More specifically, are individual particles characterized by an effective wavelength, which is related to its momentum  $p$  by  $\lambda = \frac{h}{p}$  [6], where  $h$  is the Planck constant?

**Subject (sample):** Electrons are...

# 論文寫作實例

## Introduction

**Context:** The blackbody spectrum [1], photoelectric effect [2], Compton scattering [3], and single/double-slit(s) [4] experiments have demonstrated the wave-particle duality of photons [see 5 for a pedagogical review], the quanta of electromagnetic waves. The question is, can all kinds of massive and massless particles (e.g., electrons) diffract and interfere like how photons do? More specifically, are individual particles characterized by an effective wavelength, which is related to its momentum  $p$  by  $\lambda = \frac{h}{p}$  [6], where  $h$  is the Planck constant?

**Subject (sample):** Electrons are...

學位論文文獻回顧可更具體地以多個sentences敘述各實驗內容及公式怎麼來的。但在有網路及chatGPT的現代，也不是十分必要(指導教授說的算)。

# 論文寫作實例

## Introduction

**Context:** The blackbody spectrum [1], photoelectric effect [2], Compton scattering [3], and single/double-slit(s) [4] experiments have demonstrated the wave-particle duality of photons [see 5 for a pedagogical review], the quanta of electromagnetic waves. The question is, can all kinds of massive and massless particles (e.g., electrons) diffract and interfere like how photons do? More specifically, are individual particles characterized by an effective wavelength, which is related to its momentum  $p$  by  $\lambda = \frac{h}{p}$  [6], where  $h$  is the Planck constant?

**Subject (sample):** Electrons are...

# 論文寫作實例

## Introduction

**Context:** The blackbody spectrum [1], photoelectric effect [2], Compton scattering [3], and single/double-slit(s) [4] experiments have demonstrated the wave-particle duality of photons [see 5 for a pedagogical review], the quanta of electromagnetic waves. The question is, can all kinds of massive and massless particles (e.g., electrons 啟後) diffract and interfere like how photons 啟後 do? More specifically, are individual particles characterized by an effective wavelength, which is related to its momentum  $p$  by  $\lambda = \frac{h}{p}$  [6], where  $h$  is the Planck constant?

**Subject (sample):** Electrons are... 承先

段落注意承先啟後

# 論文寫作實例

## Introduction

**Context:** The blackbody spectrum [1], photoelectric effect [2], Compton scattering [3], and single/double-slit(s) [4] experiments have demonstrated the wave-particle duality of photons [see 5 for a pedagogical review], the quanta of electromagnetic waves. The question is, can all kinds of massive and massless particles (e.g., electrons 啟後) diffract and interfere like how photons 啟後 do? More specifically, are individual particles characterized by an effective wavelength, which is related to its momentum  $p$  by  $\lambda = \frac{h}{p}$  [6], where  $h$  is the Planck constant?

**Subject (sample):** Electrons are electrically charged, spin-1/2 particles. Contrast to the massless photons, each electron has a rest mass of  $9.109 \times 10^{-31}$  kg. In the non-relativistic limit, Their collisions are ....[7]; in the relativistic limit, ... [8].

呼應Compton scattering之敘述

# 論文寫作實例(課文式贅述)

## Introduction

**Context:** The blackbody spectrum [1], photoelectric effect [2], Compton scattering [3], and single/double-slit(s) [4] experiments have demonstrated the wave-particle duality of photons [see 5 for a pedagogical review], the quanta of electromagnetic waves. The question is, can all kinds of massive and massless particles (e.g., electrons 啟後) diffract and interfere like how photons 啟後 do? More specifically, are individual particles characterized by an effective wavelength, which is related to its momentum  $p$  by  $\lambda = \frac{h}{p}$  [6], where  $h$  is the Planck constant?

**Subject (sample):** ~~There are various kinds of massive particles, for example, proton, neutron, electron, Higgs boson, etc. Protons are ... Neutrons are ... Higgs bosons are ...~~ Electrons are electrically charged, spin-1/2 particles. Contrast to the massless photons, each electron has a rest mass of  $9.109 \times 10^{-31}$  kg. In the non-relativistic limit, Their collisions are ....[7]; in the relativistic limit, ... [8].



# 論文寫作實例

## Introduction

**Context:** The blackbody spectrum [1], photoelectric effect [2], Compton scattering [3], and single/double-slit(s) [4] experiments have demonstrated the wave-particle duality of photons [see 5 for a pedagogical review], the quanta of electromagnetic waves. The question is, can all kinds of massive and massless particles (e.g., electrons) diffract and interfere like how photons do? More specifically, are individual particles characterized by an effective wavelength, which is related to its momentum  $p$  by  $\lambda = \frac{h}{p}$  [6], where  $h$  is the Planck constant?

**Subject (sample):** Electrons are electrically charged, spin-1/2 particles. Contrast to the massless photons, each electron has a rest mass of  $9.109 \times 10^{-31}$  kg. In the non-relativistic limit, Their collisions are ....[7]; in the relativistic limit, ... [8].

**Logic:** To test the wave-like behavior of electrons, we performed the double-slit experiments for electron beams (e-beams). We control the momentum of the incoming e-beams to narrow ranges, such that the wavelengths can be concretely inferred from  $\lambda = \frac{h}{p}$ . We varied the centroid momentum and the incident angle, and examined how the patterns on the screen are varied correspondingly.

# 論文寫作實例

## Introduction

**Context:** The blackbody spectrum [1], photoelectric effect [2], Compton scattering [3], and single/double-slit(s) [4] experiments have demonstrated the wave-particle duality of photons [see 5 for a pedagogical review], the quanta of electromagnetic waves. The question is, can all kinds of massive and massless particles (e.g., electrons) diffract and interfere like how photons do? More specifically, are individual particles characterized by an effective wavelength, which is related to its momentum  $p$  by  $\lambda = \frac{h}{p}$  [6], where  $h$  is the Planck constant?

**Subject (sample):** Electrons are electrically charged, spin-1/2 particles. Contrast to the massless photons, each electron has a rest mass of  $9.109 \times 10^{-31}$  kg. In the non-relativistic limit, Their collisions are ....[7]; in the relativistic limit, ... [8].

**Logic:** To test the wave-like behavior of electrons, we performed the double-slit experiments for electron beams (e-beams). We control the momentum of the incoming e-beams to narrow ranges, such that the wavelengths can be concretely inferred from  $\lambda = \frac{h}{p}$ . We varied the centroid momentum and the incident angle, and examined how the patterns on the screen are varied correspondingly.

Section 2 describes our instrumental setup and the performance of the devices. The results are in Section 3. We detected bright and dark bands on the screen, which appear similar to the interference pattern seen in the double-slit experiments for photons. We explicitly compared the patterns we detected with the patterns theoretically expected from the interference of electromagnetic waves at the same wavelength (Section 4) and discuss the physical implications. The conclusion is in Section 5.

# 論文寫作實例

約1/4 page，略為extend後即為標準的introduction section。

## Forming localized dust concentrations in a dust ring

### DM Tau case study

Haiyu Baobab Liu<sup>1</sup>

Department of Physics, National Sun Yat-Sen University, No. 70, Lien-Hai Road, Kaohsiung City 80424, Taiwan, R.O.C.  
e-mail: hy1iu.nsysu@mail.nsysu.edu.tw

Received November 25, 2025; accepted November 26, 2025

#### ABSTRACT

**Context.** Previous high-angular-resolution 225 GHz ( $\sim 1.3$  mm) continuum observations of the transitional disk DM Tau have resolved an outer ring at 20–120 au radii that is weakly azimuthally asymmetric.

**Aims.** We aim to examine dust growth and filtration in the outer ring of DM Tau.

**Methods.** We performed  $\sim 0''.06$  ( $\sim 8.7$  au) resolution Karl G. Jansky Very Large Array (JVLA) 40–48 GHz ( $\sim 7$  mm; Q band) continuum observations, along with complementary observations at lower frequencies. In addition, we analyzed the archival JVLA observations undertaken since 2010.

**Results.** Intriguingly, the Q band image resolved the azimuthally highly asymmetric, knotty dust emission sources close to the inner edge of the outer ring. Fitting the 8–700 GHz spectral energy distribution (SED) with two dust components indicates that the maximum grain size ( $a_{\text{max}}$ ) in these knotty dust emission sources is likely  $\geq 300 \mu\text{m}$ , whereas it is  $\leq 50 \mu\text{m}$  in the rest of the ring. These results may be explained by a trapping of inwardly migrating “grown” dust close to the ring inner edge. The exact mechanism for developing the azimuthal asymmetry has not yet been identified, which may be due to planet-disk interaction that might also be responsible for the creation of the dust cavity and pressure bump. Otherwise, it may be due to the fluid instabilities and vortex formation as a result of shear motions. Finally, we remark that the asymmetries in DM Tau are difficult to diagnose from the  $\geq 225$  GHz observations, owing to a high optical depth at the ring. In other words, the apparent symmetric or asymmetric morphology of the transitional disks may be related to the optical depths of those disks at the observing frequency.

**Key words.** Protoplanetary disks – Planets and satellites: formation – (ISM:) dust, extinction – Radio continuum: ISM

1	<b>1. Introduction</b>	discuss the physical implications. The conclusion is in Section 5.	28
2	The blackbody spectrum [1], photoelectric effect [2], Compton scattering [3], and single/double-slit(s) [4] experiments have demonstrated the wave-particle duality of photons [see 5 for a pedagogical review], the quanta of electromagnetic waves. The question is, can all kinds of massive and massless particles (e.g., electrons) diffract and interfere like how photons do? More specifically, are individual particles characterized by an effective wavelength, which is related to its momentum $\lambda = h/p$ [6], where $h$ is the Planck constant?		29
3	Electrons are electrically charged, spin-1/2 particles. Contrast to the massless photons, each electron has a rest mass of $9.109 \times 10^{-31}$ kg. In the non-relativistic limit, Their collisions are ...[7]; in the relativistic limit, ...[8].		
4	To test the wave-like behavior of electrons, we performed the double-slit experiments for electron beams (e-beams). We control the momentum of the incoming e-beams to narrow ranges, such that the wavelengths can be concretely inferred from $\lambda = h/p$ . We varied the centroid momentum and the incident angle, and examined how the patterns on the screen are varied correspondingly. Section 2 describes our instrumental setup and the performance of the devices. The results are in Section 3. We detected bright and dark bands on the screen, which appear similar to the interference pattern seen in the double-slit experiments for photons. We explicitly compared the patterns we detected with the patterns theoretically expected from the interference of electromagnetic waves at the same wavelength (Section 4) and		
5	<b>2. Data and reduction</b>		30
6	2.1. Data		31
7	2.2. JVLA data processing		32
8	<b>3. Results</b>		33
9	<b>4. Discussion</b>		34
10	4.1. Spectral index distribution and model		35
11	4.2. Physical implications		36
12	4.2.1. Asymmetry in protoplanetary disks		37
13	4.2.2. Dust growth in an initially smooth background disk		38
14	<b>5. Conclusion</b>		39
15	<i>Acknowledgements.</i> The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. This paper makes use of the following ALMA data: ADS/JAO.ALMA #2013.1.00498.S, #2013.1.00647.S, #2015.1.00296.S, #2016.1.00565.S, #2016.1.01042.S, #2017.1.01460.S, #2018.1.01755.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. The		40
16			41
17			42
18			43
19			44
20			45
21			46
22			47
23			48