


An Introduction to Radio Interferometry


2-1 Aperture Synthesis





You can find relevant material
on my personal webpage


NSYSU EMI Online Lecture Series Haiyu Baobab Liu (吕浩宇),
Department of Physics

























(Gaussian) deconvolution theorem



When $\delta\theta \leq \sigma$, we fundamentally cannot distinguish whether there are two stars or a (slightly elongated) celestial object.



(Gaussian) deconvolution theorem


When $\Delta\theta \leq \sigma$, we fundamentally cannot distinguish whether there are two stars or a (slightly elongated) celestial object.

$$\sigma \sim \frac{\lambda}{D}$$

Observing wavelength

Size of the reflector (in one dimension)

(Gaussian) deconvolution theorem



When $\delta\theta \leq \sigma$, we fundamentally cannot distinguish whether there are two stars or a (slightly elongated) celestial object.

Observing wavelength

Size of the reflector (in one dimension)

$$\sigma \sim \frac{\lambda}{D}$$

$$1^\circ = 60' = 3600''$$

arcminute

arcsecond

Human eyes at visible band ($\lambda \sim 0.5 \cdot 10^{-6} \text{ m}$): $\sigma \sim 1'$

(Gaussian) deconvolution theorem

When $\delta\theta \leq \sigma$, we fundamentally cannot distinguish whether there are two stars or a (slightly elongated) celestial object.

$$\sigma \sim \frac{\lambda}{D}$$

Observing wavelength

Size of the reflector
(in one dimension)

Large $D \rightarrow$ small σ , i.e.,
 collecting light from a smaller angular area
 that is proportional to $(\lambda/D)^2$
 (i.e., we receive smaller flux density)
 but we can collect photons from a bigger surface area,
 which is approximately D^2 .

$$1^\circ = 60' = 3600''$$

arcminute

arcsecond

Human eyes at visible band ($\lambda \sim 0.5 \cdot 10^{-6} \text{ m}$): $\sigma \sim 1'$

A small telescope with $D \sim 1 \text{ cm}$ at visible band ($\lambda \sim 0.5 \cdot 10^{-6} \text{ m}$): $\sigma \sim 10''$

(Gaussian) deconvolution theorem

When $\delta\theta \leq \sigma$, we fundamentally cannot distinguish whether there are two stars or a (slightly elongated) celestial object.

$$\sigma \sim \frac{\lambda}{D}$$

Observing wavelength

Size of the reflector
(in one dimension)

Large $D \rightarrow$ small σ , i.e.,

collecting light from a smaller angular area
that is proportional to $(\lambda/D)^2$

(i.e., we receive smaller flux density)

but we can collect photons from a bigger surface area,
which is approximately D^2 .

$$1^\circ = 60' = 3600''$$

arcminute
arcsecond

Human eyes at visible band ($\lambda \sim 0.5 \cdot 10^{-6} \text{ m}$): $\sigma \sim 1'$

A small telescope with $D \sim 1 \text{ cm}$ at visible band ($\lambda \sim 0.5 \cdot 10^{-6} \text{ m}$): $\sigma \sim 10''$

To achieve $\sigma \sim 10''$ at (sub)millimeter band ($\lambda \sim 10^{-3} \text{ m}$):
Requires $D \sim 2,000 \text{ cm} = 20 \text{ meter}$

Applications: cosmic microwave background anisotropy,
black hole imaging and quasar (jet) physics,
interstellar molecular lines and dust emission, etc
JCMT 15-m telescope, IRAM-30m telescope, etc

To achieve $\sigma \sim 10''$ at centimeter band ($\lambda \sim 10^{-2} \text{ m}$):
Requires $D \sim 20,000 \text{ cm} = 200 \text{ meter}$

Applications: cosmic microwave background anisotropy,
complex organic molecules,
hyperfine line transitions (e.g., HI, NH₃)
chondules and pebbles in protoplanetary disks
Greenbank 100-m telescope, Effelsberg 100-m telescope,
Arecibo 300-m telescope (cannot move), FAST 500-m telescope (cannot move), etc

(Gaussian) deconvolution theorem

When $\delta\theta \leq \sigma$, we fundamentally cannot distinguish whether there are two stars or a (slightly elongated) celestial object.

$$\sigma \sim \frac{\lambda}{D}$$

Observing wavelength
Size of the reflector (in one dimension)

Large $D \rightarrow$ small σ , i.e.,
collecting light from a smaller angular area
that is proportional to $(\lambda/D)^2$
(i.e., we receive smaller flux density)
but we can collect photons from a bigger surface area,
which is approximately D^2 .

$$1^\circ = 60' = 3600''$$

arcminute

arcsecond

Human eyes at visible band ($\lambda \sim 0.5 \cdot 10^{-6} \text{ m}$): $\sigma \sim 1'$

A small telescope with $D \sim 1 \text{ cm}$ at visible band ($\lambda \sim 0.5 \cdot 10^{-6} \text{ m}$): $\sigma \sim 10''$

To achieve $\sigma \sim 10''$ at (sub)millimeter band ($\lambda \sim 10^{-3} \text{ m}$):
Requires $D \sim 2,000 \text{ cm} = 20 \text{ meter}$

Applications: cosmic microwave background anisotropy,
black hole imaging and quasar (jet) physics,
interstellar molecular lines and dust emission, etc
JCMT 15-m telescope, IRAM-30m telescope, etc


To achieve $\sigma \sim 10''$ at centimeter band ($\lambda \sim 10^{-2} \text{ m}$):
Requires $D \sim 20,000 \text{ cm} = 200 \text{ meter}$

Applications: cosmic microwave background anisotropy,
complex organic molecules,
hyperfine line transitions (e.g., HI, NH₃)
chondules and pebbles in protoplanetary disks

Greenbank 100-m telescope, Effelsberg 100-m telescope,
Arecibo 300-m telescope (cannot move), FAST 500-m telescope (cannot move), etc

NSYSU EMI Online Lecture Series Hanyu Baobab Liu (吕浩宇),
Department of Physics

Aperture Synthesis



$$1^\circ = 60' = 3600''$$

↖ arcminute
↗ arcsecond

Human eyes at visible band ($\lambda \sim 0.5 \cdot 10^{-6} \text{ m}$): $\sigma \sim 1'$

A small telescope with $D \sim 1 \text{ cm}$ at visible band ($\lambda \sim 0.5 \cdot 10^{-6} \text{ m}$): $\sigma \sim 10''$


To achieve $\sigma \sim 10''$ at (sub)millimeter band ($\lambda \sim 10^{-3} \text{ m}$):
Requires $D \sim 2,000 \text{ cm} = 20 \text{ meter}$

Applications: cosmic microwave background anisotropy,
black hole imaging and quasar (jet) physics,
interstellar molecular lines and dust emission, etc
JCMT 15-m telescope, IRAM-30m telescope, etc

To achieve $\sigma \sim 10''$ at centimeter band ($\lambda \sim 10^{-2} \text{ m}$):
Requires $D \sim 20,000 \text{ cm} = 200 \text{ meter}$

Applications: cosmic microwave background anisotropy,
complex organic molecules,
hyperfine line transitions (e.g., HI, NH₃)
chondules and pebbles in protoplanetary disks
Greenbank 100-m telescope, Effelsberg 100-m telescope,
Arecibo 300-m telescope (cannot move), FAST 500-m telescope (cannot move), etc

Aperture Synthesis



1. Angular resolution of a telescope: $\sigma \sim \frac{\lambda}{D}$
2. An interferometric array is an observing device that the D corresponds to the **largest separation** of antennae.
3. Interferometer measure the **fourier transformed intensity distribution**