

**Problem 16.10.** Let  $i$ ,  $j$ , and  $k$ , be the unit vectors of coordinates  $(1, 0, 0)$ ,  $(0, 1, 0)$ , and  $(0, 0, 1)$  in  $\mathbb{R}^3$ .

(1) Describe geometrically the rotations defined by the following quaternions:

$$p = (0, i), \quad q = (0, j).$$

Prove that the interpolant  $Z(\lambda) = p(p^{-1}q)^\lambda$  is given by

$$Z(\lambda) = (0, \cos(\lambda\pi/2)i + \sin(\lambda\pi/2)j).$$

Describe geometrically what this rotation is.

(2) Repeat Question (1) with the rotations defined by the quaternions

$$p = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}i\right), \quad q = (0, j).$$

Prove that the interpolant  $Z(\lambda)$  is given by

$$Z(\lambda) = \left(\frac{1}{2} \cos(\lambda\pi/2), \frac{\sqrt{3}}{2} \cos(\lambda\pi/2)i + \sin(\lambda\pi/2)j\right).$$

Describe geometrically what this rotation is.

(3) Repeat Question (1) with the rotations defined by the quaternions

$$p = \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}i\right), \quad q = \left(0, \frac{1}{\sqrt{2}}(i + j)\right).$$

Prove that the interpolant  $Z(\lambda)$  is given by

$$Z(\lambda) = \left(\frac{1}{\sqrt{2}} \cos(\lambda\pi/3) - \frac{1}{\sqrt{6}} \sin(\lambda\pi/3), \right. \\ \left. (1/\sqrt{2} \cos(\lambda\pi/3) + 1/\sqrt{6} \sin(\lambda\pi/3))i + \frac{2}{\sqrt{6}} \sin(\lambda\pi/3)j\right).$$

**Problem 16.11.** Prove that

$$w \times (u \times v) = (w \cdot v)u - (u \cdot w)v.$$

Conclude that

$$u \times (u \times v) = (u \cdot v)u - (u \cdot u)v.$$