

Then prove the desired result by writing the power series for  $e^A$  and regrouping terms so that the power series for  $\cos \theta$  and  $\sin \theta$  show up. In particular

$$\begin{aligned}
 e^A &= I_3 + \sum_{p \geq 1} \frac{A^p}{p!} = I_3 + \sum_{p \geq 0} \frac{A^{2p+1}}{(2p+1)!} + \sum_{p \geq 1} \frac{A^{2p}}{(2p)!} \\
 &= I_3 + \sum_{p \geq 0} \frac{(-1)^p \theta^{2p}}{(2p+1)!} A + \sum_{p \geq 1} \frac{(-1)^{p-1} \theta^{2(p-1)}}{(2p)!} A^2 \\
 &= I_3 + \frac{A}{\theta} \sum_{p \geq 0} \frac{(-1)^p \theta^{2p+1}}{(2p+1)!} - \frac{A^2}{\theta^2} \sum_{p \geq 1} \frac{(-1)^p \theta^{2p}}{(2p)!} \\
 &= I_3 + \frac{\sin \theta}{\theta} A - \frac{A^2}{\theta^2} \sum_{p \geq 0} \frac{(-1)^p \theta^{2p}}{(2p)!} + \frac{A^2}{\theta^2} \\
 &= I_3 + \frac{\sin \theta}{\theta} A + \frac{(1 - \cos \theta)}{\theta^2} A^2,
 \end{aligned}$$

as claimed. □

The above formulae are the well-known formulae expressing a rotation of axis specified by the vector  $(a, b, c)$  and angle  $\theta$ .

The Rodrigues formula can be used to show that the exponential map  $\exp: \mathfrak{so}(3) \rightarrow \mathbf{SO}(3)$  is surjective.

Given any rotation matrix  $R \in \mathbf{SO}(3)$ , we have the following cases:

- (1) The case  $R = I$  is trivial.
- (2) If  $R \neq I$  and  $\text{tr}(R) \neq -1$ , then

$$\exp^{-1}(R) = \left\{ \frac{\theta}{2 \sin \theta} (R - R^T) \mid 1 + 2 \cos \theta = \text{tr}(R) \right\}.$$

(Recall that  $\text{tr}(R) = r_{11} + r_{22} + r_{33}$ , the *trace* of the matrix  $R$ ).

Then there is a unique skew-symmetric  $B$  with corresponding  $\theta$  satisfying  $0 < \theta < \pi$  such that  $e^B = R$ .

- (3) If  $R \neq I$  and  $\text{tr}(R) = -1$ , then  $R$  is a rotation by the angle  $\pi$  and things are more complicated, but a matrix  $B$  can be found. We leave this part as a good exercise: see Problem 17.8.

The computation of a logarithm of a rotation in  $\mathbf{SO}(3)$  as sketched above has applications in kinematics, robotics, and motion interpolation.

As an immediate corollary of the Gram–Schmidt orthonormalization procedure, we obtain the  $QR$ -decomposition for invertible matrices.