

with  $\exp(0_3) = I_3$ .

(2) Prove that  $e^A$  is an orthogonal matrix of determinant  $+1$ , i.e., a rotation matrix.

(3) Prove that the exponential map  $\exp: \mathfrak{so}(3) \rightarrow \mathbf{SO}(3)$  is surjective. For this proceed as follows: Pick any rotation matrix  $R \in \mathbf{SO}(3)$ ;

(1) The case  $R = I$  is trivial.

(2) If  $R \neq I$  and  $\text{tr}(R) \neq -1$ , then

$$\exp^{-1}(R) = \left\{ \frac{\theta}{2 \sin \theta} (R - R^T) \mid 1 + 2 \cos \theta = \text{tr}(R) \right\}.$$

(Recall that  $\text{tr}(R) = r_{11} + r_{22} + r_{33}$ , the *trace* of the matrix  $R$ ).

Show that there is a unique skew-symmetric  $B$  with corresponding  $\theta$  satisfying  $0 < \theta < \pi$  such that  $e^B = R$ .

(3) If  $R \neq I$  and  $\text{tr}(R) = -1$ , then prove that the eigenvalues of  $R$  are  $1, -1, -1$ , that  $R = R^T$ , and that  $R^2 = I$ . Prove that the matrix

$$S = \frac{1}{2}(R - I)$$

is a symmetric matrix whose eigenvalues are  $-1, -1, 0$ . Thus  $S$  can be diagonalized with respect to an orthogonal matrix  $Q$  as

$$S = Q \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} Q^T.$$

Prove that there exists a skew symmetric matrix

$$U = \begin{pmatrix} 0 & -d & c \\ d & 0 & -b \\ -c & b & 0 \end{pmatrix}$$

so that

$$U^2 = S = \frac{1}{2}(R - I).$$

Observe that

$$U^2 = \begin{pmatrix} -(c^2 + d^2) & bc & bd \\ bc & -(b^2 + d^2) & cd \\ bd & cd & -(b^2 + c^2) \end{pmatrix},$$