

Unfortunately, even though  $L^{-1} = E_{n-1} \cdots E_2 E_1$ , the matrices  $E_k$  occur in the wrong order and the  $k$ th column of  $L^{-1}$  is *not* the  $k$ th column of  $E_k$ .

Here is an example illustrating the method.

**Example 8.2.** Given

$$A = A_1 = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & -1 & 0 & 1 \\ 1 & 1 & -1 & 0 \\ 1 & -1 & 0 & -1 \end{pmatrix},$$

we have the following sequence of steps: The first pivot is  $\pi_1 = 1$  in row 1, and we subtract row 1 from rows 2, 3, and 4. We get

$$A_2 = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & -2 & -1 & 1 \\ 0 & 0 & -2 & 0 \\ 0 & -2 & -1 & -1 \end{pmatrix} \quad L_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}.$$

The next pivot is  $\pi_2 = -2$  in row 2, and we subtract row 2 from row 4 (and add 0 times row 2 to row 3). We get

$$A_3 = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & -2 & -1 & 1 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix} \quad L_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{pmatrix}.$$

The next pivot is  $\pi_3 = -2$  in row 3, and since the fourth entry in column 3 is already a zero, we add 0 times row 3 to row 4. We get

$$A_4 = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & -2 & -1 & 1 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix} \quad L_3 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{pmatrix}.$$

The procedure is finished, and we have

$$L = L_3 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{pmatrix} \quad U = A_4 = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & -2 & -1 & 1 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix}.$$

It is easy to check that indeed

$$LU = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & -2 & -1 & 1 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & -1 & 0 & 1 \\ 1 & 1 & -1 & 0 \\ 1 & -1 & 0 & -1 \end{pmatrix} = A.$$

We now show how to extend the above method to deal with pivoting efficiently. This is the  $PA = LU$  factorization.