

matrix to rref, we can apply elementary row operations to the matrix  $[u_k \ \Gamma]$ , which consists of rows  $1, \dots, m$  of the tableau.

Once the new matrix  $\Gamma^+$  is obtained, the new reduced costs are given by the following proposition.

**Proposition 46.2.** *Given any Linear Program (P2) in standard form*

$$\begin{aligned} & \text{maximize} \quad cx \\ & \text{subject to} \quad Ax = b \text{ and } x \geq 0, \end{aligned}$$

where  $A$  is an  $m \times n$  matrix of rank  $m$ , if  $(u, K)$  is a basic (not necessarily feasible) solution of (P2) and if  $K^+ = (K - \{k^-\}) \cup \{j^+\}$ , with  $K = (k_1, \dots, k_m)$  and  $k^- = k_\ell$ , then for  $i = 1, \dots, n$  we have

$$c_i - c_{K^+} \gamma_{K^+}^i = c_i - c_K \gamma_K^i - \frac{\gamma_{k^-}^i}{\gamma_{j^+}^i} (c_{j^+} - c_K \gamma_K^{j^+}).$$

Using the reduced cost notation, the above equation is

$$(\bar{c}_{K^+})_i = (\bar{c}_K)_i - \frac{\gamma_{k^-}^i}{\gamma_{j^+}^i} (\bar{c}_K)_{j^+}.$$

*Proof.* Without any loss of generality and to simplify notation assume that  $K = (1, \dots, m)$  and write  $j$  for  $j^+$  and  $\ell$  for  $k_m$ . Since  $\gamma_K^i = A_K^{-1} A^i$ ,  $\gamma_{K^+}^i = A_{K^+}^{-1} A^i$ , and  $A_{K^+} = A_K E(\gamma_K^j)$ , we have

$$c_i - c_{K^+} \gamma_{K^+}^i = c_i - c_{K^+} A_{K^+}^{-1} A^i = c_i - c_{K^+} E(\gamma_K^j)^{-1} A_K^{-1} A^i = c_i - c_{K^+} E(\gamma_K^j)^{-1} \gamma_K^i,$$

where

$$E(\gamma_K^j)^{-1} = \begin{pmatrix} 1 & & & & -(\gamma_\ell^j)^{-1} \gamma_1^j & & \\ & \ddots & & & \vdots & & \\ & & 1 & & -(\gamma_\ell^j)^{-1} \gamma_{\ell-1}^j & & \\ & & & (\gamma_\ell^j)^{-1} & & & \\ & & & -(\gamma_\ell^j)^{-1} \gamma_{\ell+1}^j & 1 & & \\ & & & \vdots & & \ddots & \\ & & & -(\gamma_\ell^j)^{-1} \gamma_m^j & & & 1 \end{pmatrix}$$

where the  $\ell$ th column contains the  $\gamma$ s. Since  $c_{K^+} = (c_1, \dots, c_{\ell-1}, c_j, c_{\ell+1}, \dots, c_m)$ , we have

$$c_{K^+} E(\gamma_K^j)^{-1} = \left( c_1, \dots, c_{\ell-1}, \frac{c_j}{\gamma_\ell^j} - \sum_{k=1, k \neq \ell}^m c_k \frac{\gamma_k^j}{\gamma_\ell^j}, c_{\ell+1}, \dots, c_m \right),$$