

(or contraction, if $\lambda < 1$). When $\lambda = 0$, $H_{a,0}(x) = a$ for all $x \in E$, and $H_{a,0}$ is the constant affine map sending every point to a . If we assume $\lambda \neq 1$, note that $H_{a,\lambda}$ is never the identity, and since a is a fixed point, $H_{a,\lambda}$ is never a translation.

We now consider the set \widehat{E} of geometric transformations from E to E , consisting of the union of the (disjoint) sets of translations and dilatations of ratio $\lambda \neq 1$. We would like to give this set the structure of a vector space, in such a way that both E and \vec{E} can be naturally embedded into \widehat{E} . In fact, it will turn out that barycenters show up quite naturally too!

In order to “add” two dilatations H_{a_1,λ_1} and H_{a_2,λ_2} , it turns out that it is more convenient to consider dilatations of the form $H_{a,1-\lambda}$, where $\lambda \neq 0$. To see this, let us see the effect of such a dilatation on a point $x \in E$: We have

$$H_{a,1-\lambda}(x) = a + (1 - \lambda)\vec{ax} = a + \vec{ax} - \lambda\vec{ax} = x + \lambda\vec{ax}.$$

For simplicity of notation, let us denote $H_{a,1-\lambda}$ by $\langle a, \lambda \rangle$. Then, we have

$$\langle a, \lambda \rangle(x) = x + \lambda\vec{ax}.$$

Remarks:

- (1) Note that $H_{a,1-\lambda}(x) = H_{x,\lambda}(a)$.
- (2) Berger defines a map $h: E \rightarrow \vec{E}$ as a *vector field*. Thus, each $\langle a, \lambda \rangle$ can be viewed as the vector field $x \mapsto \lambda\vec{ax}$. Similarly, a translation u can be viewed as the constant vector field $x \mapsto u$. Thus, we could define \widehat{E} as the (disjoint) union of these two vector fields. We prefer our view in terms of geometric transformations.

Then, since

$$\langle a_1, \lambda_1 \rangle(x) = x + \lambda_1\vec{xa_1} \quad \text{and} \quad \langle a_2, \lambda_2 \rangle(x) = x + \lambda_2\vec{xa_2},$$

if we want to define $\langle a_1, \lambda_1 \rangle \widehat{+} \langle a_2, \lambda_2 \rangle$, we see that we have to distinguish between two cases:

- (1) $\lambda_1 + \lambda_2 = 0$. In this case, since

$$\lambda_1\vec{xa_1} + \lambda_2\vec{xa_2} = \lambda_1\vec{xa_1} - \lambda_1\vec{xa_2} = \lambda_1\vec{a_2a_1},$$

we let

$$\langle a_1, \lambda_1 \rangle \widehat{+} \langle a_2, \lambda_2 \rangle = \lambda_1\vec{a_2a_1},$$

where $\lambda_1\vec{a_2a_1}$ denotes the translation associated with the vector $\lambda_1\vec{a_2a_1}$.

- (2) $\lambda_1 + \lambda_2 \neq 0$. In this case, the points a_1 and a_2 assigned the weights $\lambda_1/(\lambda_1 + \lambda_2)$ and $\lambda_2/(\lambda_1 + \lambda_2)$ have a barycenter

$$b = \frac{\lambda_1}{\lambda_1 + \lambda_2}a_1 + \frac{\lambda_2}{\lambda_1 + \lambda_2}a_2,$$