

for every  $\epsilon > 0$ , there is some  $\eta > 0$ , such that, for every  $x \in E$ ,

$$\text{if } \|x - a\|_E \leq \eta, \text{ then } \|f(x) - f(a)\|_F \leq \epsilon.$$

It is worth noting that continuity is a topological notion, in the sense that equivalent metrics (or equivalent norms) define exactly the same notion of continuity.

**Definition 37.17.** If  $(E, \mathcal{O}_E)$  and  $(F, \mathcal{O}_F)$  are topological spaces, and  $f: E \rightarrow F$  is a function, for every nonempty subset  $A \subseteq E$  of  $E$ , we say that  $f$  is *continuous on  $A$*  if the restriction of  $f$  to  $A$  is continuous with respect to  $(A, \mathcal{U})$  and  $(F, \mathcal{O}_F)$ , where  $\mathcal{U}$  is the subspace topology induced by  $\mathcal{O}_E$  on  $A$ .

Given a product  $E_1 \times \cdots \times E_n$  of topological spaces, as usual, we let  $\pi_i: E_1 \times \cdots \times E_n \rightarrow E_i$  be the projection function such that,  $\pi_i(x_1, \dots, x_n) = x_i$ . It is immediately verified that each  $\pi_i$  is continuous.

Given a topological space  $(E, \mathcal{O})$ , we say that a point  $a \in E$  is *isolated* if  $\{a\}$  is an open set in  $\mathcal{O}$ . Then if  $(E, \mathcal{O}_E)$  and  $(F, \mathcal{O}_F)$  are topological spaces, any function  $f: E \rightarrow F$  is continuous at every isolated point  $a \in E$ . In the discrete topology, every point is isolated.

In a nontrivial normed vector space  $(E, \|\cdot\|)$  (with  $E \neq \{0\}$ ), no point is isolated. To show this, we show that every open ball  $B_0(u, \rho)$  contains some vectors different from  $u$ . Indeed, since  $E$  is nontrivial, there is some  $v \in E$  such that  $v \neq 0$ , and thus  $\lambda = \|v\| > 0$  (by (N1)). Let

$$w = u + \frac{\rho}{\lambda + 1}v.$$

Since  $v \neq 0$  and  $\rho > 0$ , we have  $w \neq u$ . Then,

$$\|w - u\| = \left\| \frac{\rho}{\lambda + 1}v \right\| = \frac{\rho\lambda}{\lambda + 1} < \rho,$$

which shows that  $\|w - u\| < \rho$ , for  $w \neq u$ .

The following proposition is easily shown.

**Proposition 37.10.** *Given topological spaces  $(E, \mathcal{O}_E)$ ,  $(F, \mathcal{O}_F)$ , and  $(G, \mathcal{O}_G)$ , and two functions  $f: E \rightarrow F$  and  $g: F \rightarrow G$ , if  $f$  is continuous at  $a \in E$  and  $g$  is continuous at  $f(a) \in F$ , then  $g \circ f: E \rightarrow G$  is continuous at  $a \in E$ . Given  $n$  topological spaces  $(F_i, \mathcal{O}_i)$ , for every function  $f: E \rightarrow F_1 \times \cdots \times F_n$ , then  $f$  is continuous at  $a \in E$  iff every  $f_i: E \rightarrow F_i$  is continuous at  $a$ , where  $f_i = \pi_i \circ f$ .*

One can also show that in a metric space  $(E, d)$ , the distance  $d: E \times E \rightarrow \mathbb{R}$  is continuous, where  $E \times E$  has the product topology. By the triangle inequality, we have

$$d(x, y) \leq d(x, x_0) + d(x_0, y_0) + d(y_0, y) = d(x_0, y_0) + d(x_0, x) + d(y_0, y)$$