

and $\varphi(u_i) = 1 \otimes u_i$. Next, if ρ is injective, by definition of the scalar multiplication in the A -module $\rho_*(\rho^*(M))$, we have $\varphi(a_1u_1 + \cdots + a_nu_n) = 0$ iff

$$\rho(a_1)\varphi(u_1) + \cdots + \rho(a_n)\varphi(u_n) = 0,$$

and since $(\varphi(u_1), \dots, \varphi(u_n))$ is a basis of $\rho^*(M)$, we must have $\rho(a_i) = 0$ for $i = 1, \dots, n$, which (by injectivity of ρ) implies that $a_i = 0$ for $i = 1, \dots, n$. Therefore, φ is injective. \square

In particular, if A is a subring of B , then ρ is the inclusion map and Proposition 35.41 shows that a basis of M becomes a basis of $M_{(B)}$ and that M is embedded into $M_{(B)}$. It is also easy to see that if M and N are two free A -modules and $f: M \rightarrow N$ is a linear map represented by the matrix X with respect to some bases (u_1, \dots, u_n) of M and (v_1, \dots, v_m) of N , then the B -linear map \bar{f} is also represented by the matrix X over the bases $(\varphi(u_1), \dots, \varphi(u_n))$ and $(\varphi(v_1), \dots, \varphi(v_m))$.

Proposition 35.41 yields another proof of the fact that any two bases of a free A -module have the same cardinality. Indeed, if \mathfrak{m} is a maximal ideal in the ring A , then we have the quotient ring homomorphism $\pi: A \rightarrow A/\mathfrak{m}$, and we get the A/\mathfrak{m} -module $\pi^*(M)$. If M is free, any basis (u_1, \dots, u_n) of M becomes the basis $(\varphi(u_1), \dots, \varphi(u_n))$ of $\pi^*(M)$; but A/\mathfrak{m} is a field, so the dimension n is uniquely determined. This argument also applies to an infinite basis $(u_i)_{i \in I}$. Observe that by Proposition 35.14, we have an isomorphism

$$\pi^*(M) = (A/\mathfrak{m}) \otimes_A M \approx M/\mathfrak{m}M,$$

so $M/\mathfrak{m}M$ is a vector space over the field A/\mathfrak{m} , which is the argument used in Theorem 35.1.

Proposition 35.42. *Given a ring homomorphism $\rho: A \rightarrow B$, for any two A -modules M and N , there is a unique isomorphism*

$$\rho^*(M) \otimes_B \rho^*(N) \approx \rho^*(M \otimes_A N),$$

such that $(1 \otimes u) \otimes (1 \otimes v) \mapsto 1 \otimes (u \otimes v)$, for all $u \in M$ and all $v \in N$.

The proof uses identities from Proposition 33.13. It is not hard but it requires a little gymnastic; a good exercise for the reader.