

Before defining the natural generalization of an inner product, it is convenient to define semilinear maps.

Definition 14.1. Given two vector spaces E and F over the complex field \mathbb{C} , a function $f: E \rightarrow F$ is *semilinear* if

$$\begin{aligned} f(u + v) &= f(u) + f(v), \\ f(\lambda u) &= \bar{\lambda}f(u), \end{aligned}$$

for all $u, v \in E$ and all $\lambda \in \mathbb{C}$.

Remark: Instead of defining semilinear maps, we could have defined the vector space \bar{E} as the vector space with the same carrier set E whose addition is the same as that of E , but whose multiplication by a complex number is given by

$$(\lambda, u) \mapsto \bar{\lambda}u.$$

Then it is easy to check that a function $f: E \rightarrow \mathbb{C}$ is semilinear iff $f: \bar{E} \rightarrow \mathbb{C}$ is linear.

We can now define sesquilinear forms and Hermitian forms.

Definition 14.2. Given a complex vector space E , a function $\varphi: E \times E \rightarrow \mathbb{C}$ is a *sesquilinear form* if it is linear in its first argument and semilinear in its second argument, which means that

$$\begin{aligned} \varphi(u_1 + u_2, v) &= \varphi(u_1, v) + \varphi(u_2, v), \\ \varphi(u, v_1 + v_2) &= \varphi(u, v_1) + \varphi(u, v_2), \\ \varphi(\lambda u, v) &= \lambda\varphi(u, v), \\ \varphi(u, \mu v) &= \bar{\mu}\varphi(u, v), \end{aligned}$$

for all $u, v, u_1, u_2, v_1, v_2 \in E$, and all $\lambda, \mu \in \mathbb{C}$. A function $\varphi: E \times E \rightarrow \mathbb{C}$ is a *Hermitian form* if it is sesquilinear and if

$$\varphi(v, u) = \overline{\varphi(u, v)}$$

for all $u, v \in E$.

Obviously, $\varphi(0, v) = \varphi(u, 0) = 0$. Also note that if $\varphi: E \times E \rightarrow \mathbb{C}$ is sesquilinear, we have

$$\varphi(\lambda u + \mu v, \lambda u + \mu v) = |\lambda|^2\varphi(u, u) + \lambda\bar{\mu}\varphi(u, v) + \bar{\lambda}\mu\varphi(v, u) + |\mu|^2\varphi(v, v),$$

and if $\varphi: E \times E \rightarrow \mathbb{C}$ is Hermitian, we have

$$\varphi(\lambda u + \mu v, \lambda u + \mu v) = |\lambda|^2\varphi(u, u) + 2\Re(\lambda\bar{\mu}\varphi(u, v)) + |\mu|^2\varphi(v, v).$$

Note that restricted to real coefficients, a sesquilinear form is bilinear (we sometimes say \mathbb{R} -bilinear).