

of *algebraic geometry*, a beautiful but formidable subject. For a taste of algebraic geometry, see Lang [109] or Dummit and Foote [54].

The duality theorem (Theorem 11.4) shows that the situation is much simpler if we restrict our attention to linear subspaces; in this case

$$U = \mathcal{I}(\mathcal{Z}(U)) \quad \text{and} \quad V = \mathcal{Z}(\mathcal{I}(V)).$$

Proposition 11.3. *We have $V \subseteq V^{00}$ for every subspace V of E , and $U \subseteq U^{00}$ for every subspace U of E^* .*

Proof. Indeed, for any $v \in V$, to show that $v \in V^{00}$ we need to prove that $u^*(v) = 0$ for all $u^* \in V^0$. However, V^0 consists of all linear forms u^* such that $u^*(y) = 0$ for all $y \in V$; in particular, for a fixed $v \in V$, we have $u^*(v) = 0$ for all $u^* \in V^0$, as required.

Similarly, for any $u^* \in U$, to show that $u^* \in U^{00}$ we need to prove that $u^*(v) = 0$ for all $v \in U^0$. However, U^0 consists of all vectors v such that $f^*(v) = 0$ for all $f^* \in U$; in particular, for a fixed $u^* \in U$, we have $u^*(v) = 0$ for all $v \in U^0$, as required. \square

We will see shortly that in finite dimension, we have $V = V^{00}$ and $U = U^{00}$.



However, even though $V = V^{00}$ is always true, when E is of infinite dimension, it is not always true that $U = U^{00}$.

Given a vector space E and a subspace U of E , by Theorem 3.7, every basis $(u_i)_{i \in I}$ of U can be extended to a basis $(u_j)_{j \in I \cup J}$ of E , where $I \cap J = \emptyset$.

11.3 The Duality Theorem and Some Consequences

We have the following important theorem adapted from E. Artin [6] (Chapter 1).

Theorem 11.4. (*Duality theorem*) *Let E be a vector space. The following properties hold:*

- (a) *For every basis $(u_i)_{i \in I}$ of E , the family $(u_i^*)_{i \in I}$ of coordinate forms is linearly independent.*
- (b) *For every subspace V of E , we have $V^{00} = V$.*
- (c) *For every subspace V of finite codimension m of E , for every subspace W of E such that $E = V \oplus W$ (where W is of finite dimension m), for every basis $(u_i)_{i \in I}$ of E such that (u_1, \dots, u_m) is a basis of W , the family (u_1^*, \dots, u_m^*) is a basis of the orthogonal V^0 of V in E^* , so that*

$$\dim(V^0) = \text{codim}(V).$$

Furthermore, we have $V^{00} = V$.