

Definition 44.6. Given any (nonempty) subset S of \mathbb{R}^n , the smallest convex set containing S is denoted by $\text{conv}(S)$ and called the *convex hull of S* (it is the intersection of all convex sets containing S).

It is essential not only to have a good understanding of $\text{conv}(S)$, but to also have good methods for computing it. We have the following simple but crucial result.

Proposition 44.1. For any family $S = (a_i)_{i \in I}$ of points in \mathbb{R}^n , the set V of convex combinations $\sum_{i \in I} \lambda_i a_i$ (where $\sum_{i \in I} \lambda_i = 1$ and $\lambda_i \geq 0$) is the convex hull $\text{conv}(S)$ of $S = (a_i)_{i \in I}$.

It is natural to wonder whether Proposition 44.1 can be sharpened in two directions: (1) Is it possible to have a fixed bound on the number of points involved in the convex combinations? (2) Is it necessary to consider convex combinations of all points, or is it possible to consider only a subset with special properties?

The answer is yes in both cases. In Case 1, Carathéodory's theorem asserts that it is enough to consider convex combinations of $n + 1$ points. For example, in the plane \mathbb{R}^2 , the convex hull of a set S of points is the union of all triangles (interior points included) with vertices in S . In Case 2, the theorem of Krein and Milman asserts that a convex set that is also compact is the convex hull of its extremal points (given a convex set S , a point $a \in S$ is *extremal* if $S - \{a\}$ is also convex).

We will not prove these theorems here, but we invite the reader to consult Gallier [73] or Berger [12].

Convex sets also arise as half-spaces cut out by affine hyperplanes.

Definition 44.7. An *affine form* $\varphi: \mathbb{R}^n \rightarrow \mathbb{R}$ is defined by some linear form $c \in (\mathbb{R}^n)^*$ and some scalar $\beta \in \mathbb{R}$ so that

$$\varphi(x) = cx + \beta \quad \text{for all } x \in \mathbb{R}^n.$$

If $c \neq 0$, the affine form φ specified by (c, β) defines the *affine hyperplane* (for short *hyperplane*) $H(\varphi)$ given by

$$H(\varphi) = \{x \in \mathbb{R}^n \mid \varphi(x) = 0\} = \{x \in \mathbb{R}^n \mid cx + \beta = 0\},$$

and the two (*closed*) *half-spaces*

$$\begin{aligned} H_+(\varphi) &= \{x \in \mathbb{R}^n \mid \varphi(x) \geq 0\} = \{x \in \mathbb{R}^n \mid cx + \beta \geq 0\}, \\ H_-(\varphi) &= \{x \in \mathbb{R}^n \mid \varphi(x) \leq 0\} = \{x \in \mathbb{R}^n \mid cx + \beta \leq 0\}. \end{aligned}$$

When $\beta = 0$, we call H a *linear hyperplane*.

Both $H_+(\varphi)$ and $H_-(\varphi)$ are convex and $H = H_+(\varphi) \cap H_-(\varphi)$.

For example, $\varphi: \mathbb{R}^2 \rightarrow \mathbb{R}$ with $\varphi(x, y) = 2x + y + 3$ is an affine form defining the line given by the equation $y = -2x - 3$. Another example of an affine form is $\varphi: \mathbb{R}^3 \rightarrow \mathbb{R}$