

In general, there is no closed-form formula for the exponential  $e^A$  of a matrix  $A$ , but for skew symmetric matrices of dimension 2 and 3, there are explicit formulae. Everyone should enjoy computing the exponential  $e^A$  where

$$A = \begin{pmatrix} 0 & -\theta \\ \theta & 0 \end{pmatrix}.$$

If we write

$$J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

then

$$A = \theta J$$

The key property is that

$$J^2 = -I.$$

**Proposition 9.22.** *If  $A = \theta J$ , then*

$$e^A = \cos \theta I + \sin \theta J = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

*Proof.* We have

$$\begin{aligned} A^{4n} &= \theta^{4n} I_2, \\ A^{4n+1} &= \theta^{4n+1} J, \\ A^{4n+2} &= -\theta^{4n+2} I_2, \\ A^{4n+3} &= -\theta^{4n+3} J, \end{aligned}$$

and so

$$e^A = I_2 + \frac{\theta}{1!} J - \frac{\theta^2}{2!} I_2 - \frac{\theta^3}{3!} J + \frac{\theta^4}{4!} I_2 + \frac{\theta^5}{5!} J - \frac{\theta^6}{6!} I_2 - \frac{\theta^7}{7!} J + \cdots.$$

Rearranging the order of the terms, we have

$$e^A = \left( 1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \frac{\theta^6}{6!} + \cdots \right) I_2 + \left( \frac{\theta}{1!} - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \frac{\theta^7}{7!} + \cdots \right) J.$$

We recognize the power series for  $\cos \theta$  and  $\sin \theta$ , and thus

$$e^A = \cos \theta I_2 + \sin \theta J,$$

that is

$$e^A = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix},$$

as claimed. □