

where  $P$  is a product of elementary row operations. Because the bottom row of  $A'$  is zero, the system  $A'x = 0$  has at most  $n - 1$  nontrivial equations, and by Proposition 8.17, this system has a nontrivial solution  $x$ . But then,  $Ax = P^{-1}A'x = 0$  with  $x \neq 0$ , contradicting the fact that the system  $Ax = 0$  is assumed to have only the trivial solution. Therefore, (d) implies (a) and the proof is complete.  $\square$

Proposition 8.18 yields a method for computing the inverse of an invertible matrix  $A$ : reduce  $A$  to the identity using elementary row operations, obtaining

$$E_p \cdots E_1 A = I.$$

Multiplying both sides by  $A^{-1}$  we get

$$A^{-1} = E_p \cdots E_1.$$

From a practical point of view, we can build up the product  $E_p \cdots E_1$  by reducing to row echelon form the augmented  $n \times 2n$  matrix  $(A, I_n)$  obtained by adding the  $n$  columns of the identity matrix to  $A$ . This is just another way of performing the Gauss–Jordan procedure.

Here is an example: let us find the inverse of the matrix

$$A = \begin{pmatrix} 5 & 4 \\ 6 & 5 \end{pmatrix}.$$

We form the  $2 \times 4$  block matrix

$$(A, I) = \begin{pmatrix} 5 & 4 & 1 & 0 \\ 6 & 5 & 0 & 1 \end{pmatrix}$$

and apply elementary row operations to reduce  $A$  to the identity. For example:

$$(A, I) = \begin{pmatrix} 5 & 4 & 1 & 0 \\ 6 & 5 & 0 & 1 \end{pmatrix} \longrightarrow \begin{pmatrix} 5 & 4 & 1 & 0 \\ 1 & 1 & -1 & 1 \end{pmatrix}$$

by subtracting row 1 from row 2,

$$\begin{pmatrix} 5 & 4 & 1 & 0 \\ 1 & 1 & -1 & 1 \end{pmatrix} \longrightarrow \begin{pmatrix} 1 & 0 & 5 & -4 \\ 1 & 1 & -1 & 1 \end{pmatrix}$$

by subtracting  $4 \times$  row 2 from row 1,

$$\begin{pmatrix} 1 & 0 & 5 & -4 \\ 1 & 1 & -1 & 1 \end{pmatrix} \longrightarrow \begin{pmatrix} 1 & 0 & 5 & -4 \\ 0 & 1 & -6 & 5 \end{pmatrix} = (I, A^{-1}),$$

by subtracting row 1 from row 2. Thus

$$A^{-1} = \begin{pmatrix} 5 & -4 \\ -6 & 5 \end{pmatrix}.$$

Proposition 8.18 can also be used to give an elementary proof of the fact that if a square matrix  $A$  has a left inverse  $B$  (resp. a right inverse  $B$ ), so that  $BA = I$  (resp.  $AB = I$ ), then  $A$  is invertible and  $A^{-1} = B$ . This is an interesting exercise, try it!