



Figure 24.14: Affine independence and linear independence

Definition 24.4 is reasonable, because by Proposition 24.4, the independence of the family $(\overrightarrow{a_i a_j})_{j \in (I - \{i\})}$ does not depend on the choice of a_i . A crucial property of linearly independent vectors (u_1, \dots, u_m) is that if a vector v is a linear combination

$$v = \sum_{i=1}^m \lambda_i u_i$$

of the u_i , then the λ_i are unique. A similar result holds for affinely independent points.

Proposition 24.5. *Given an affine space $\langle E, \vec{E}, + \rangle$, let (a_0, \dots, a_m) be a family of $m + 1$ points in E . Let $x \in E$, and assume that $x = \sum_{i=0}^m \lambda_i a_i$, where $\sum_{i=0}^m \lambda_i = 1$. Then, the family $(\lambda_0, \dots, \lambda_m)$ such that $x = \sum_{i=0}^m \lambda_i a_i$ is unique iff the family $(\overrightarrow{a_0 a_1}, \dots, \overrightarrow{a_0 a_m})$ is linearly independent.*

Proof. The proof is straightforward and is omitted. It is also given in Gallier [70]. \square

Proposition 24.5 suggests the notion of affine frame. Affine frames are the affine analogues of bases in vector spaces. Let $\langle E, \vec{E}, + \rangle$ be a nonempty affine space, and let (a_0, \dots, a_m) be a family of $m + 1$ points in E . The family (a_0, \dots, a_m) determines the family of m vectors $(\overrightarrow{a_0 a_1}, \dots, \overrightarrow{a_0 a_m})$ in \vec{E} . Conversely, given a point a_0 in E and a family of m vectors (u_1, \dots, u_m) in \vec{E} , we obtain the family of $m + 1$ points (a_0, \dots, a_m) in E , where $a_i = a_0 + u_i$, $1 \leq i \leq m$.

Thus, for any $m \geq 1$, it is equivalent to consider a family of $m + 1$ points (a_0, \dots, a_m) in E , and a pair $(a_0, (u_1, \dots, u_m))$, where the u_i are vectors in \vec{E} . Figure 24.14 illustrates the notion of affine independence.