

Under the Standard Margin Hypothesis, there is some index  $i_0$  such that  $0 < \lambda_{i_0} < K$  and there is some index  $j_0$  such that  $0 < \mu_{j_0} < K$ , and we obtain the value of  $b$  and  $\delta$  as

$$b = \frac{1}{2}(\langle w, \varphi(u_{i_0}) \rangle + \langle w, \varphi(v_{j_0}) \rangle)$$

$$\delta = \frac{1}{2}(\langle w, \varphi(u_{i_0}) \rangle - \langle w, \varphi(v_{j_0}) \rangle).$$

Using the above value for  $w$ , we obtain

$$b = \frac{\sum_{i=1}^p \lambda_i (\kappa(u_i, u_{i_0}) + \kappa(u_i, v_{j_0})) - \sum_{j=1}^q \mu_j (\kappa(v_j, u_{i_0}) + \kappa(v_j, v_{j_0}))}{2 \left( \begin{pmatrix} \lambda^\top & \mu^\top \end{pmatrix} K \begin{pmatrix} \lambda \\ \mu \end{pmatrix} \right)^{1/2}}.$$

It follows that the classification function

$$f(x) = \text{sgn}(\langle w, \varphi(x) \rangle - b)$$

is given by

$$f(x) = \text{sgn} \left( \sum_{i=1}^p \lambda_i (2\kappa(u_i, x) - \kappa(u_i, u_{i_0}) - \kappa(u_i, v_{j_0})) - \sum_{j=1}^q \mu_j (2\kappa(v_j, x) - \kappa(v_j, u_{i_0}) - \kappa(v_j, v_{j_0})) \right),$$

which is solely expressed in terms of the kernel  $\kappa$ .

Kernel methods for SVM are discussed in Schölkopf and Smola [145] and Shawe-Taylor and Christianini [159].

## 54.2 Solving SVM (SVM<sub>s1</sub>) Using ADMM

In order to solve (SVM<sub>s1</sub>) using ADMM we need to write the matrix corresponding to the constraints in equational form,

$$\sum_{i=1}^p \lambda_i - \sum_{j=1}^q \mu_j = 0$$

$$\sum_{i=1}^p \lambda_i + \sum_{j=1}^q \mu_j = 1$$

$$\lambda_i + \alpha_i = K, \quad i = 1, \dots, p$$

$$\mu_j + \beta_j = K, \quad j = 1, \dots, q.$$