

- (1) Consider combinations  $\sum_{i \in I} \lambda_i u_i$  for which

$$\sum_{i \in I} \lambda_i = 1.$$

These are called *affine combinations*. One should realize that every linear combination  $\sum_{i \in I} \lambda_i u_i$  can be viewed as an affine combination. For example, if  $k$  is an index not in  $I$ , if we let  $J = I \cup \{k\}$ ,  $u_k = 0$ , and  $\lambda_k = 1 - \sum_{i \in I} \lambda_i$ , then  $\sum_{j \in J} \lambda_j u_j$  is an affine combination and

$$\sum_{i \in I} \lambda_i u_i = \sum_{j \in J} \lambda_j u_j.$$

However, we get new spaces. For example, in  $\mathbb{R}^3$ , the set of all affine combinations of the three vectors  $e_1 = (1, 0, 0)$ ,  $e_2 = (0, 1, 0)$ , and  $e_3 = (0, 0, 1)$ , is the plane passing through these three points. Since it does not contain  $0 = (0, 0, 0)$ , it is not a linear subspace.

- (2) Consider combinations  $\sum_{i \in I} \lambda_i u_i$  for which

$$\lambda_i \geq 0, \quad \text{for all } i \in I.$$

These are called *positive* (or *conic*) *combinations*. It turns out that positive combinations of families of vectors are *cones*. They show up naturally in convex optimization.

- (3) Consider combinations  $\sum_{i \in I} \lambda_i u_i$  for which we require (1) *and* (2), that is

$$\sum_{i \in I} \lambda_i = 1, \quad \text{and} \quad \lambda_i \geq 0 \quad \text{for all } i \in I.$$

These are called *convex combinations*. Given any finite family of vectors, the set of all convex combinations of these vectors is a *convex polyhedron*. Convex polyhedra play a very important role in convex optimization.

**Remark:** The notion of linear combination can also be defined for infinite index sets  $I$ . To ensure that a sum  $\sum_{i \in I} \lambda_i u_i$  makes sense, we restrict our attention to families of finite support.

**Definition 3.5.** Given any field  $K$ , a family of scalars  $(\lambda_i)_{i \in I}$  has *finite support* if  $\lambda_i = 0$  for all  $i \in I - J$ , for some finite subset  $J$  of  $I$ .

If  $(\lambda_i)_{i \in I}$  is a family of scalars of finite support, for any vector space  $E$  over  $K$ , for any (possibly infinite) family  $(u_i)_{i \in I}$  of vectors  $u_i \in E$ , we define the linear combination  $\sum_{i \in I} \lambda_i u_i$  as the finite linear combination  $\sum_{j \in J} \lambda_j u_j$ , where  $J$  is any finite subset of  $I$  such that  $\lambda_i = 0$  for all  $i \in I - J$ . In general, results stated for finite families also hold for families of finite support.