



Figure 5.3: Second averages and second half differences.

$(w_1, w_2, w_3, w_4, w_5, w_6, w_7, w_8)$ is given by the matrix

$$W = \begin{pmatrix} 1 & 1 & 1 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & -1 & 0 & 0 & 0 \\ 1 & 1 & -1 & 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & -1 & 0 & 0 & -1 & 0 & 0 \\ 1 & -1 & 0 & 1 & 0 & 0 & 1 & 0 \\ 1 & -1 & 0 & 1 & 0 & 0 & -1 & 0 \\ 1 & -1 & 0 & -1 & 0 & 0 & 0 & 1 \\ 1 & -1 & 0 & -1 & 0 & 0 & 0 & -1 \end{pmatrix}.$$

The columns of this matrix are orthogonal, and it is easy to see that

$$W^{-1} = \text{diag}(1/8, 1/8, 1/4, 1/4, 1/2, 1/2, 1/2, 1/2)W^T.$$

A pattern is beginning to emerge. It looks like the second Haar basis vector w_2 is the “mother” of all the other basis vectors, except the first, whose purpose is to perform averaging. Indeed, in general, given

$$w_2 = \underbrace{(1, \dots, 1, -1, \dots, -1)}_{2^n},$$

the other Haar basis vectors are obtained by a “scaling and shifting process.” Starting from w_2 , the scaling process generates the vectors

$$w_3, w_5, w_9, \dots, w_{2^j+1}, \dots, w_{2^{n-1}+1},$$

such that w_{2^j+1} is obtained from w_{2^j+1} by forming two consecutive blocks of 1 and -1 of half the size of the blocks in w_{2^j+1} , and setting all other entries to zero. Observe that w_{2^j+1} has 2^j blocks of 2^{n-j} elements. The shifting process consists in shifting the blocks of 1 and -1 in w_{2^j+1} to the right by inserting a block of $(k-1)2^{n-j}$ zeros from the left, with $0 \leq j \leq n-1$ and $1 \leq k \leq 2^j$. Note that our convention is to use j as the scaling index and k as the shifting index. Thus, we obtain the following formula for w_{2^j+k} :

$$w_{2^j+k}(i) = \begin{cases} 0 & 1 \leq i \leq (k-1)2^{n-j} \\ 1 & (k-1)2^{n-j} + 1 \leq i \leq (k-1)2^{n-j} + 2^{n-j-1} \\ -1 & (k-1)2^{n-j} + 2^{n-j-1} + 1 \leq i \leq k2^{n-j} \\ 0 & k2^{n-j} + 1 \leq i \leq 2^n, \end{cases}$$