

Example 12.12. Consider polynomials over $[-1, 1]$, with the symmetric bilinear form

$$\langle f, g \rangle = \int_{-1}^1 \frac{1}{\sqrt{1-t^2}} f(t)g(t)dt.$$

We leave it as an exercise to prove that the above defines an inner product. It can be shown that the polynomials $T_n(x)$ given by

$$T_n(x) = \cos(n \arccos x), \quad n \geq 0,$$

(equivalently, with $x = \cos \theta$, we have $T_n(\cos \theta) = \cos(n\theta)$) are orthogonal with respect to the above inner product. These polynomials are the *Chebyshev polynomials*. Their norm is not equal to 1. Instead, we have

$$\langle T_n, T_n \rangle = \begin{cases} \frac{\pi}{2} & \text{if } n > 0, \\ \pi & \text{if } n = 0. \end{cases}$$

Using the identity $(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$ and the binomial formula, we obtain the following expression for $T_n(x)$:

$$T_n(x) = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} (x^2 - 1)^k x^{n-2k}.$$

The Chebyshev polynomials are defined inductively as follows:

$$\begin{aligned} T_0(x) &= 1 \\ T_1(x) &= x \\ T_{n+1}(x) &= 2xT_n(x) - T_{n-1}(x), \quad n \geq 1. \end{aligned}$$

Using these recurrence equations, we can show that

$$T_n(x) = \frac{(x - \sqrt{x^2 - 1})^n + (x + \sqrt{x^2 - 1})^n}{2}.$$

The polynomial T_n has n distinct roots in the interval $[-1, 1]$. The Chebyshev polynomials play an important role in approximation theory. They are used as an approximation to a best polynomial approximation of a continuous function under the sup-norm (∞ -norm).

The inner products of the last two examples are special cases of an inner product of the form

$$\langle f, g \rangle = \int_{-1}^1 W(t)f(t)g(t)dt,$$

where $W(t)$ is a *weight function*. If W is a continuous function such that $W(x) > 0$ on $(-1, 1)$, then the above bilinear form is indeed positive definite. Families of orthogonal