

**Theorem 17.15.** *Given a Euclidean space  $E$  of dimension  $n$ , for every skew-self-adjoint linear map  $f: E \rightarrow E$  there is an orthonormal basis  $(e_1, \dots, e_n)$  such that the matrix of  $f$  w.r.t. this basis is a block diagonal matrix of the form*

$$\begin{pmatrix} A_1 & & \cdots & \\ & A_2 & & \\ \vdots & \vdots & \ddots & \vdots \\ & & \cdots & A_p \end{pmatrix}$$

such that each block  $A_j$  is either 0 or a two-dimensional matrix of the form

$$A_j = \begin{pmatrix} 0 & -\mu_j \\ \mu_j & 0 \end{pmatrix},$$

where  $\mu_j \in \mathbb{R}$ , with  $\mu_j > 0$ . In particular, the eigenvalues of  $f_{\mathbb{C}}$  are pure imaginary of the form  $\pm i\mu_j$  or 0.

*Proof.* The case where  $n = 1$  is trivial. As in the proof of Theorem 17.12,  $f_{\mathbb{C}}$  has some eigenvalue  $z = \lambda + i\mu$ , where  $\lambda, \mu \in \mathbb{R}$ . We claim that  $\lambda = 0$ . First we show that

$$\langle f(w), w \rangle = 0$$

for all  $w \in E$ . Indeed, since  $f = -f^*$ , we get

$$\langle f(w), w \rangle = \langle w, f^*(w) \rangle = \langle w, -f(w) \rangle = -\langle w, f(w) \rangle = -\langle f(w), w \rangle,$$

since  $\langle -, - \rangle$  is symmetric. This implies that

$$\langle f(w), w \rangle = 0.$$

Applying this to  $u$  and  $v$  and using the fact that

$$f(u) = \lambda u - \mu v \quad \text{and} \quad f(v) = \mu u + \lambda v,$$

we get

$$0 = \langle f(u), u \rangle = \langle \lambda u - \mu v, u \rangle = \lambda \langle u, u \rangle - \mu \langle u, v \rangle$$

and

$$0 = \langle f(v), v \rangle = \langle \mu u + \lambda v, v \rangle = \mu \langle u, v \rangle + \lambda \langle v, v \rangle,$$

from which, by addition, we get

$$\lambda(\langle v, v \rangle + \langle v, v \rangle) = 0.$$

Since  $u \neq 0$  or  $v \neq 0$ , we have  $\lambda = 0$ .

Then going back to the proof of Theorem 17.12, unless  $\mu = 0$ , the case where  $u$  and  $v$  are orthogonal and span a subspace of dimension 2 applies, and the induction shows that all the blocks are two-dimensional or reduced to 0.  $\square$