

In addition,  $d \neq 0$ , and  $d$  is unique up to multiplication by a nonzero scalar in  $K$ .

As a consequence of Proposition 30.15, some polynomials  $f_1, \dots, f_n \in K[X]$  are relatively prime iff there exist  $u_1, \dots, u_n \in K[X]$  such that

$$u_1 f_1 + \dots + u_n f_n = 1.$$

The identity

$$u_1 f_1 + \dots + u_n f_n = 1$$

of part (2) of Proposition 30.15 is also called the *Bezout identity*.

We now consider the factorization of polynomials of a single variable into irreducible factors.

## 30.5 Factorization and Irreducible Factors in $K[X]$

**Definition 30.9.** Given a field  $K$ , a polynomial  $p \in K[X]$  is *irreducible or indecomposable or prime* if  $\deg(p) \geq 1$  and if  $p$  is not divisible by any polynomial  $q \in K[X]$  such that  $1 \leq \deg(q) < \deg(p)$ . Equivalently,  $p$  is irreducible if  $\deg(p) \geq 1$  and if  $p = q_1 q_2$ , then either  $q_1 \in K$  or  $q_2 \in K$  (and of course,  $q_1 \neq 0$ ,  $q_2 \neq 0$ ).

**Example 30.2.** Every polynomial  $aX + b$  of degree 1 is irreducible. Over the field  $\mathbb{R}$ , the polynomial  $X^2 + 1$  is irreducible (why?), but  $X^3 + 1$  is not irreducible, since

$$X^3 + 1 = (X + 1)(X^2 - X + 1).$$

The polynomial  $X^2 - X + 1$  is irreducible over  $\mathbb{R}$  (why?). It would seem that  $X^4 + 1$  is irreducible over  $\mathbb{R}$ , but in fact,

$$X^4 + 1 = (X^2 - \sqrt{2}X + 1)(X^2 + \sqrt{2}X + 1).$$

However, in view of the above factorization,  $X^4 + 1$  is irreducible over  $\mathbb{Q}$ .

It can be shown that the irreducible polynomials over  $\mathbb{R}$  are the polynomials of degree 1, or the polynomials of degree 2 of the form  $aX^2 + bX + c$ , for which  $b^2 - 4ac < 0$  (i.e., those having no real roots). This is not easy to prove! Over the complex numbers  $\mathbb{C}$ , the only irreducible polynomials are those of degree 1. This is a version of a fact often referred to as the “Fundamental theorem of Algebra”, or, as the French sometimes say, as “d’Alembert’s theorem”!

We already observed that for any two nonzero polynomials  $f, g \in K[X]$ ,  $f$  divides  $g$  iff  $(g) \subseteq (f)$ . In view of the definition of a maximal ideal given in Definition 30.4, we now prove that a polynomial  $p \in K[X]$  is irreducible iff  $(p)$  is a maximal ideal in  $K[X]$ .

**Proposition 30.16.** *A polynomial  $p \in K[X]$  is irreducible iff  $(p)$  is a maximal ideal in  $K[X]$ .*