

In general, for an overdetermined  $m \times n$  system  $Ax = b$ , what Gauss and Legendre discovered is that there are solutions  $x$  minimizing

$$\|Ax - b\|_2^2$$

(where  $\|u\|_2^2 = u_1^2 + \cdots + u_n^2$ , the square of the Euclidean norm of the vector  $u = (u_1, \dots, u_n)$ ), and that these solutions are given by the square  $n \times n$  system

$$A^\top Ax = A^\top b,$$

called the *normal equations*. Furthermore, when the columns of  $A$  are linearly independent, it turns out that  $A^\top A$  is invertible, and so  $x$  is unique and given by

$$x = (A^\top A)^{-1} A^\top b.$$

Note that  $A^\top A$  is a symmetric matrix, one of the nice features of the normal equations of a least squares problem. For instance, since the above problem in matrix form is represented as

$$\begin{pmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \end{pmatrix} \begin{pmatrix} d \\ c \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix},$$

the normal equations are

$$\begin{pmatrix} 3 & x_1 + x_2 + x_3 \\ x_1 + x_2 + x_3 & x_1^2 + x_2^2 + x_3^2 \end{pmatrix} \begin{pmatrix} d \\ c \end{pmatrix} = \begin{pmatrix} y_1 + y_2 + y_3 \\ x_1 y_1 + x_2 y_2 + x_3 y_3 \end{pmatrix}.$$

In fact, given any real  $m \times n$  matrix  $A$ , there is always a unique  $x^+$  of minimum norm that minimizes  $\|Ax - b\|_2^2$ , even when the columns of  $A$  are linearly dependent. How do we prove this, and how do we find  $x^+$ ?

**Theorem 23.1.** *Every linear system  $Ax = b$ , where  $A$  is an  $m \times n$  matrix, has a unique least squares solution  $x^+$  of smallest norm.*

*Proof.* Geometry offers a nice proof of the existence and uniqueness of  $x^+$ . Indeed, we can interpret  $b$  as a point in the Euclidean (affine) space  $\mathbb{R}^m$ , and the image subspace of  $A$  (also called the column space of  $A$ ) as a subspace  $U$  of  $\mathbb{R}^m$  (passing through the origin). Then it is clear that

$$\inf_{x \in \mathbb{R}^n} \|Ax - b\|_2^2 = \inf_{y \in U} \|y - b\|_2^2,$$

with  $U = \text{Im } A$ , and we claim that  $x$  minimizes  $\|Ax - b\|_2^2$  iff  $Ax = p$ , where  $p$  the orthogonal projection of  $b$  onto the subspace  $U$ .

Recall from Section 13.1 that the orthogonal projection  $p_U: U \oplus U^\perp \rightarrow U$  is the linear map given by

$$p_U(u + v) = u,$$