

**Example 4.1.** Let  $E = F = \mathbb{R}^2$ , with  $u_1 = (1, 0)$ ,  $u_2 = (0, 1)$ ,  $v_1 = (1, 1)$  and  $v_2 = (-1, 1)$ . The change of basis matrix  $P$  from the basis  $\mathcal{U} = (u_1, u_2)$  to the basis  $\mathcal{V} = (v_1, v_2)$  is

$$P = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$$

and its inverse is

$$P^{-1} = \begin{pmatrix} 1/2 & 1/2 \\ -1/2 & 1/2 \end{pmatrix}.$$

The old coordinates  $(x_1, x_2)$  with respect to  $(u_1, u_2)$  are expressed in terms of the new coordinates  $(x'_1, x'_2)$  with respect to  $(v_1, v_2)$  by

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix},$$

and the new coordinates  $(x'_1, x'_2)$  with respect to  $(v_1, v_2)$  are expressed in terms of the old coordinates  $(x_1, x_2)$  with respect to  $(u_1, u_2)$  by

$$\begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} = \begin{pmatrix} 1/2 & 1/2 \\ -1/2 & 1/2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}.$$

**Example 4.2.** Let  $E = F = \mathbb{R}[X]_3$  be the set of polynomials of degree at most 3, and consider the bases  $\mathcal{U} = (1, x, x^2, x^3)$  and  $\mathcal{V} = (B_0^3(x), B_1^3(x), B_2^3(x), B_3^3(x))$ , where  $B_0^3(x), B_1^3(x), B_2^3(x), B_3^3(x)$  are the *Bernstein polynomials* of degree 3, given by

$$B_0^3(x) = (1-x)^3 \quad B_1^3(x) = 3(1-x)^2x \quad B_2^3(x) = 3(1-x)x^2 \quad B_3^3(x) = x^3.$$

By expanding the Bernstein polynomials, we find that the change of basis matrix  $P_{\mathcal{V}, \mathcal{U}}$  is given by

$$P_{\mathcal{V}, \mathcal{U}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -3 & 3 & 0 & 0 \\ 3 & -6 & 3 & 0 \\ -1 & 3 & -3 & 1 \end{pmatrix}.$$

We also find that the inverse of  $P_{\mathcal{V}, \mathcal{U}}$  is

$$P_{\mathcal{V}, \mathcal{U}}^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1/3 & 0 & 0 \\ 1 & 2/3 & 1/3 & 0 \\ 1 & 1 & 1 & 1 \end{pmatrix}.$$

Therefore, the coordinates of the polynomial  $2x^3 - x + 1$  over the basis  $\mathcal{V}$  are

$$\begin{pmatrix} 1 \\ 2/3 \\ 1/3 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1/3 & 0 & 0 \\ 1 & 2/3 & 1/3 & 0 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \\ 0 \\ 2 \end{pmatrix},$$

and so

$$2x^3 - x + 1 = B_0^3(x) + \frac{2}{3}B_1^3(x) + \frac{1}{3}B_2^3(x) + 2B_3^3(x).$$