

Case for an EeV Gravitino

Emilian Dudas,^{1,*} Yann Mambrini,^{2,†} and Keith A. Olive^{3,‡}

¹*CPHT, Ecole Polytechnique, 91128 Palaiseau Cedex, France*

²*Laboratoire de Physique, Théorique Université Paris-Sud, F-91405 Orsay, France*

³*William I. Fine Theoretical Physics Institute, School of Physics and Astronomy,
University of Minnesota, Minneapolis, Minnesota 55455, USA*

(Received 26 April 2017; published 31 July 2017)

We consider the possibility that supersymmetry is broken above the inflationary mass scale and that the only “low” energy remnant of supersymmetry is the gravitino with a mass of the order of the EeV scale. The gravitino in this class of models becomes a candidate for the dark matter of the Universe. To avoid the overproduction of gravitinos from the decays of the next-to-lightest supersymmetric particle we argue that the supersymmetric spectrum must lie above the inflationary mass scale ($M_{\text{SUSY}} > 10^{-5} M_{\text{P}} \sim 10^{13}$ GeV). Since $m_{3/2} \approx M_{\text{SUSY}}^2 / M_{\text{P}}$, we expect $m_{3/2} \gtrsim 0.2$ EeV. Cosmological constraints then predict a relatively large reheating temperature between 10^{10} and 10^{12} GeV.

DOI: 10.1103/PhysRevLett.119.051801

Introduction.—To date, there is no significant experimental signal for weak scale (TeV) supersymmetry at the LHC [1]. In parallel, direct detection experiments such as XENON100 [2], LUX [3], or PandaX [4] have set strong limits on the elastic scattering cross section of neutralinos on nucleons exceeding common pre-run I LHC predictions [5,6]. This may indicate one of the following: (i) low energy supersymmetry is still around the corner waiting to be discovered at a slightly higher energy scale [6,7]; (ii) part of the supersymmetric spectrum lies at very high energy as in split supersymmetry [8]; (iii) essentially the entire supersymmetric spectrum lies at very high energy as in supersplit supersymmetry [9] (aka the standard model). Here, we consider the possibility that the only remnant of supersymmetry surviving down to energies significantly below the Planck scale is the gravitino. (We will consider the reduced Planck mass $M_{\text{P}}^2 = 1/8\pi G_N \approx 2.4 \times 10^{18}$ GeV throughout the Letter.)

The gravitino may either be an excellent dark matter candidate [10–19] or a severe cosmological problem [20,21]. If the gravitino is the lightest supersymmetric particle (LSP) and therefore a dark matter candidate, there is the risk of overproduction from the decay of the next-to-lightest supersymmetric particle (NLSP) [21–23]. In fact, as we discuss below, a combination of limits from big bang nucleosynthesis (BBN) [24] and the relic density allows us to place an upper limit of ≈ 4 TeV on the gravitino mass. However, if the sparticle spectrum lies above the inflationary mass scale, and none of the superpartners are ever produced after inflationary reheating, we derive a lower limit on the gravitino mass of 0.2 EeV and the gravitino may once again become a dark matter candidate. Note that such a spectrum implies that supersymmetry is nonlinearly realized [25]. Finally, the thermal production of gravitinos allows us to bound reheat temperature after inflation between a few $\times 10^{10}$ and 10^{12} GeV.

The Letter is organized as follows. We first discuss limits on the gravitino mass in typical supersymmetric models. We discuss both the limits from BBN and from NLSP decay. We then consider a high scale supersymmetric model where only the gravitino lies below the inflationary mass scale. We derive a new lower limit to the gravitino mass in this case. Assuming that the gravitino is the dark matter, we consider general consequences for inflationary models, particularly aspects of reheating.

Upper limits to the gravitino mass in typical SUSY scenarios.—The physics behind the limits on the gravitino mass can be very different depending on the specific mass range under consideration. With the exception of the cases of light (MeV, keV, or sub-keV) masses, typical gravitino masses discussed in the literature are in the 10–1000 GeV range similar to the masses expected for MSSM superpartners if the SUSY scale is related to the hierarchy problem. However, it is well known that a gravitino with $O(100)$ GeV mass is potentially problematic [20,21]. On the one hand, if it is not the LSP, it will decay to lighter sparticles, and if it is the LSP, the NLSP would decay to the gravitino. In either case, the lifetime may easily fall within the range of 100– 10^8 s and be subject to constraints from BBN [24,26–31]. For example, the decay rate of a neutralino NLSP to a gravitino and photon is given by [14,15,31]

$$\Gamma_{\text{decay}} \approx \frac{C^2}{16\pi} \frac{m_{\tilde{\chi}}^5}{m_{3/2}^2 M_{\text{P}}^2}, \quad (1)$$

where C depends on the neutralino diagonalization matrix and we have ignored phase space factors [and other factors of $O(1)$]. In the case of a gravitino LSP, there are typically strong constraints on the SUSY parameter space forcing one into regions where the NLSP is the tau slepton [28,29].

The BBN constraints begin to be relaxed when the lifetime of the NLSP becomes less than $O(100)$ s [26,27], and for a neutralino NLSP, we can use Eq. (1) to obtain a relation between the neutralino and gravitino masses,

$$\tau_\chi \lesssim 100 \text{ s} \Rightarrow m_\chi > 300 \text{ GeV} \left(\frac{m_{3/2}}{\text{GeV}} \right)^{2/5} \quad (2)$$

for $C \sim 1$. Thus, avoiding the limits from BBN will require a rather heavy SUSY spectrum for TeV scale (and above) gravitino masses. We note that the relaxation of the BBN bound at 100 s requires satisfying the upper bound on the density of decaying particles of roughly [26] $m_\chi n_\chi / n_\gamma \lesssim 7 \times 10^{-9}$ GeV. If we exceed this density, we must use the more strict BBN bound of $\tau_\chi \lesssim 0.1$ s. In this case, the lower limit on m_χ in Eq. (2) is increased by a factor of ~ 4 .

In addition to the BBN constraints, there is an additional constraint coming from the relic density of the NLSP whose decay contributes to the relic density of gravitinos [21–23]. The gravitino relic density from NLSP decays can be written simply as

$$\Omega_{3/2} h^2 = \frac{m_{3/2}}{m_\chi} \Omega_\chi h^2, \quad (3)$$

and, thus, the NLSP relic density is limited by

$$\Omega_\chi h^2 \lesssim 0.12 \frac{m_\chi}{m_{3/2}}, \quad (4)$$

where 0.12 is the approximate upper limit on the cold dark matter density from the PLANCK experiment [32]. As long as m_χ is not much greater than $m_{3/2}$, the NLSP density is constrained to be near the cold dark matter density. Even in the event that $m_\chi \gg m_{3/2}$, the relic density of the NLSP is still constrained by the BBN unless its lifetime is very short (< 0.1 s) as noted above.

Thus as we attempt to increase the mass of a gravitino LSP, we are forced to higher NLSP masses to insure both a relatively short lifetime and low relic density. For example, for $m_{3/2} = 2$ TeV, we must require $m_\chi \gtrsim 6$ TeV (20 TeV) to obtain $\tau_\chi < 100$ s (< 0.1 s). Generally, it is very difficult to obtain an acceptable neutralino relic density when the neutralino masses surpass the TeV scale [6,7]. In particular, the neutralino relic density in the TeV regime must be regulated by either some strong resonant process or coannihilation. Indeed, the strongest such process involves the coannihilation with the gluino [33–36]. Pushing the mass scales to their limit (when the neutralino and gluino masses are degenerate), an upper limit to the neutralino mass of roughly 8 TeV was found [34–36]. This translates [using Eq. (2)] to an upper bound on the gravitino mass of roughly $m_{3/2} < 4$ TeV.

High scale SUSY breaking and inflation—EeV scale gravitinos.—High scale SUSY.—In order to go beyond the derived upper limit on the gravitino mass of 4 TeV, we must make a more substantial departure from the common paradigm of weak scale supersymmetry. To this end, we consider the possibility for higher gravitino masses along with a very high SUSY breaking scale, leaving only the gravitino surviving at low energies as a dark matter candidate.

As we demonstrated in the previous section, a gravitino mass in excess of 4 TeV, would require a SUSY spectrum in excess of 8 TeV in order to obtain NLSP lifetimes short enough to be compatible with constraints from BBN. However, even in the limit of degenerate neutralinos and gluinos, strong coannihilations are insufficient to lower the NLSP relic density to acceptable levels. Further increasing the SUSY mass scale, weakens the interaction strengths, lowering the annihilation (and coannihilation) cross sections, leading to an overabundance. Without resorting to some unknown form of dilution, one possibility for larger gravitino masses is to move the SUSY matter spectrum to such high scales, so that SUSY particles were never part of the thermal bath after inflation.

To completely remove the supersymmetric particle spectrum from the thermal history, we must assume that the SUSY mass spectrum is larger than both the inflationary reheating temperature, T_R , and the inflaton mass, m_ϕ , so as to prevent SUSY particles from being produced by either thermal processes during reheating or by the decay of the inflaton. Here, we will not tie ourselves to a particular inflationary model, but note that in many models considered, the inflaton mass is set by amplitude of density perturbations seen in the microwave background, and yields a value of roughly 3×10^{13} GeV. When we need to refer to a specific example, we consider a no-scale supergravity model of inflation [37] which leads to Starobinsky-like inflation [38].

If we denote as F the order parameter for supersymmetry breaking, then typical soft SUSY masses will be proportional to F ,

$$M_{\text{SUSY}} = \frac{F}{\Lambda_{\text{mess}}}, \quad (5)$$

where Λ_{mess} is the mass scale associated with the mediators of supersymmetry breaking. (These messengers could in principle also play a role in restoring unification at high scale.) We expect $\Lambda_{\text{mess}} \geq M_{\text{SUSY}}$. Thus $M_{\text{SUSY}} > m_\phi$ translates to $F > m_\phi^2$. The gravitino mass is also determined by F [39],

$$m_{3/2} = \frac{F}{\sqrt{3} M_P}. \quad (6)$$

And, hence, we have a *lower* bound on the gravitino mass given by

$$m_{3/2} > \frac{m_\phi^2}{\sqrt{3}M_P} \simeq 0.2 \text{ EeV}. \quad (7)$$

Thus, we have a gravitino mass gap between 4 TeV and 0.2 EeV which remains cosmologically problematic. As a toy model, one could consider supersymmetry breaking using a Polonyi-like field, z [40] with superpotential $W = F(z + \Delta)$, where Δ is a constant. In this case, the Goldstino is the fermionic component of z eaten by the gravitino. However, the transmission of supersymmetry breaking to the observable sector cannot be of standard gravity type. One option is (general) gauge mediation, with a high scale for messenger masses and supersymmetry breaking.

Gravitino production.—Clearly the LHC bounds can be satisfied if the sparticle mass spectrum lies above a few TeV. The direct detection limits can also be satisfied as the spectrum approaches its upper limit [7]. It is also possible that the dark matter lies beyond the MSSM and has weaker couplings to matter, e.g., through a t -channel exchange of a massive Z' or Higgs boson as shown in Ref. [41] or invoking a pseudoscalar or pure axial mediator to velocity suppress σ_N^{scat} [42,43]. Furthermore, if the dark matter couples too weakly with the standard model, it will never reach thermal equilibrium as its production rate is $(dn/dt) = n_\gamma^2 \langle \sigma v \rangle$. The particle is frozen in during the process of thermalization. The weak coupling of the dark sector with the standard model can be due to either an effectively small coupling (of the order of 10^{-10}) [44] or because the mass of the mediator between the two sectors is very large, as in the case of nonequilibrium thermal dark matter (NETDM) models [45].

By increasing the SUSY mass scale, we have also removed most of the standard gravitino production mechanisms. Namely, both NSLP decay, and the thermal production from standard model annihilations such as gluon, gluon \rightarrow gluino, gravitino are no longer kinematically allowed. The rate for the latter is well known [46,47] and scales as $\Gamma \sim T^3 M_{\text{SUSY}}^2 / M_P^2 m_{3/2}^2$, where we have assumed predominantly goldstino production in the limit $m_{3/2} \ll M_{\text{SUSY}}$. In this case, the gravitino abundance is approximately $n_{3/2}/n_\gamma \sim \Gamma/H \sim T M_{\text{SUSY}}^2 / M_P m_{3/2}^2$, where we have simply taken the Hubble parameter as T^2/M_P .

In the limit that the SUSY mass scale is above the inflationary scale, there remains, however, (at least) two sources of gravitino production. Inflaton decay to gravitinos [47,48], and thermal production of two gravitinos from the thermal bath (gluon, gluon \rightarrow gravitino, gravitino) [49], as this is only kinematically allowed channel. A careful computation of the gravitino production rate was derived in [49]

$$R = n^2 \langle \sigma v \rangle \simeq 21.65 \times \frac{T^{12}}{F^4}, \quad (8)$$

where n is the number density of incoming states and we see that the rate has a strong dependence on temperature and is even stronger than the NETDM case [45] where the dependence is $R(T) \propto T^8$. This dependence can be easily ascertained on dimensional grounds. Recall that $n \propto T^3$, and for gravitino production, we expect $\langle \sigma v \rangle \propto T^6/F^4$. The consequences of such a high temperature dependence are important: we expect that all gravitino production will occur early and rapidly in the reheating process. This differs from the feebly coupled case [44], where the smallness of the dark matter coupling to the standard model bath renders the production rate slower.

From the rate $R(T)$, we can determine that $\Gamma \sim R/n \sim T^9/M_P^4 m_{3/2}^4$ (again assuming $m_{3/2} \ll M_{\text{SUSY}}$) leading to a gravitino abundance $n_{3/2}/n_\gamma \sim \Gamma/H \sim T^7/M_P^3 m_{3/2}^4$. More precisely, we find,

$$\Omega_{3/2} h^2 \simeq 0.11 \left(\frac{0.1 \text{ EeV}}{m_{3/2}} \right)^3 \left(\frac{T_{RH}}{2.0 \times 10^{10} \text{ GeV}} \right)^7 \quad (9)$$

In the absence of direct inflaton decays, a gravitino at the lower mass limit (7) would require a reheating temperature of roughly 3×10^{10} GeV, above the upper limit allowed by the relic abundance constraint ($T_R \lesssim 10^7$ GeV) in the more common thermal scenario [46], thus favoring thermal leptogenesis [50,51].

Consequences for inflationary models.—The reheating temperature appearing in Eq. (9) is generated by the decay of an inflaton field ϕ of mass m_ϕ and width Γ_ϕ . We assume that the decay and thermalization occur instantaneously at the time t_ϕ , $\Gamma_\phi t_\phi = 2\Gamma_\phi/3H = c$, where $c \approx 1.2$ is a constant. In this case, the reheating temperature is given by [47,52]

$$T_{RH} = \left(\frac{10}{g_s} \right)^{1/4} \left(\frac{2\Gamma_\phi M_P}{\pi c} \right)^{1/2} = 0.55 \frac{y_\phi}{2\pi} \left(\frac{m_\phi M_P}{c} \right)^{1/2}, \quad (10)$$

where we have defined a standard “Yukawa-like” coupling y_ϕ of the inflaton field to the thermal bath, $\Gamma_\phi = (y_\phi^2/8\pi)m_\phi$ and g_s is the effective number of light degrees of freedom in this case set by the standard model, $g_s = 427/4$. We can then reexpress the relic abundance (9) as a function of y_ϕ :

$$\Omega_{3/2} h^2 \simeq 0.11 \left(\frac{0.1 \text{ EeV}}{m_{3/2}} \right)^3 \left(\frac{m_\phi}{3 \times 10^{13} \text{ GeV}} \right)^{7/2} \times \left(\frac{y_\phi}{2.9 \times 10^{-5}} \right)^7, \quad (11)$$

where we have set $c = 1.2$. The cosmological constraint is plotted in Fig. (1) in the $(m_{3/2}, y_\phi)$ plane, where we show the region allowed by PLANCK [32]. The black (solid) line represents the PLANCK constraint $\Omega h^2 = 0.11$.




FIG. 1. Region of the parameter space allowed by PLANCK constraints [32] in the plane $(m_{3/2}, y_\phi)$, for different values of the branching ratio $B_{3/2}$ and $m_\phi = 3 \times 10^{13}$ GeV (see the text for details).

One immediately sees the linear increase in the Yukawa coupling y_ϕ with increasing gravitino mass in order to counterbalance the weakening of the effective coupling $1/F$ responsible for its production in the thermal bath.

A large inflaton-matter coupling produces a high reheating temperature, which in turn increases the gravitino abundance. Then, as one can see from Eq. (11), the solid curve in Fig. 1 is an upper bound on y_ϕ to avoid an overabundant gravitino. In fact, one can extract an upper bound on y_ϕ independent of $m_{3/2}$ simply requiring $m_{3/2} < T_{RH}$, a necessary condition for the gravitino to be thermally produced. The condition $m_{3/2} < T_{RH}$ implemented in Eq. (11) with the expression (10) gives

$$y_\phi \lesssim 1.6 \times 10^{-3} \left(\frac{3 \times 10^{13} \text{ GeV}}{m_\phi} \right)^{1/2}, \quad (12)$$

shown as the horizontal dashed line in the Fig. 1. We can then extract the maximum reheating temperature $T_{RH} \lesssim 1.1 \times 10^{12}$ GeV. Combined with the condition (7) $m_{3/2} > 0.2$ EeV, the relic abundance constraint (9) gives

$$2.7 \times 10^{10} \text{ GeV} \lesssim T_{RH} \lesssim 1.1 \times 10^{12} \text{ GeV}, \quad (13)$$

which is a strong prediction of our model.

Gravitino production by inflaton decay.—It is also possible to produce gravitinos through the direct decay of the inflaton. For example, in no-scale supergravity models of inflation, the decay of the inflaton to gravitinos is highly suppressed. In simple models, there is no coupling at the tree level [53]. However, it is possible to couple the inflaton to moduli without spoiling the inflationary potential [47,48]. We can parametrize the decay to a pair of gravitinos as $\Gamma_{3/2} = m_\phi (y_{3/2}^2 / 72\pi)$.

The branching ratio of decays to gravitinos is then

$$B_{3/2} = \Gamma_{3/2} / \Gamma_\phi = \frac{|y_{3/2}|^2}{9y_\phi^2}. \quad (14)$$

Using the result from Ref. [47] for the gravitino abundance produced by inflaton decay at the epoch of reheating, we get

$$\frac{n_{3/2}}{n_\gamma} \approx 3.6 B_{3/2} \frac{(\Gamma_\phi M_P)^{1/2}}{m_\phi} \approx 0.7 B_{3/2} y_\phi \left(\frac{M_P}{m_\phi} \right)^{1/2}, \quad (15)$$

corresponding to

$$\Omega_{3/2}^{\text{decay}} h^2 = 0.11 \left(\frac{B_{3/2}}{1.3 \times 10^{-13}} \right) \left(\frac{y_\phi}{2.9 \times 10^{-5}} \right) \times \left(\frac{m_{3/2}}{0.1 \text{ EeV}} \right) \left(\frac{3 \times 10^{13} \text{ GeV}}{m_\phi} \right)^{1/2}. \quad (16)$$

today.

The condition (7) is then translated into

$$B_{3/2} y_\phi = \frac{|y_{3/2}|^2}{9|y_\phi|} \lesssim 1.9 \times 10^{-18} \left(\frac{0.1 \text{ EeV}}{m_{3/2}} \right) \quad (17)$$

for $m_\phi = 3 \times 10^{13}$ GeV. Contrary to the case of thermal gravitino production, our limit to the coupling y_ϕ is strengthened as $m_{3/2}$ is increased when gravitino production occurs through inflaton decay. Since the density through the decay of the inflaton is proportional to $n_\phi B_{3/2} m_{3/2}$, where $m_\phi n_\phi$ is the inflaton energy density, the limit on the coupling is improved when either the branching ratio or the gravitino mass is increased.

This result is also shown in Fig. 1 where we clearly see the changing in the slope for a larger value of $B_{3/2} > 10^{-19}$ where the direct production from inflaton decay may dominate over the thermal production. We note that the constraints obtained on the inflaton coupling to gravitinos are strong. We recall, however, that in no-scale models of inflation [47,48,53] and in classes of inflationary models with so-called stabilized field [54,55], this coupling is naturally very small. Finally, we point out that in the case of the direct production of the gravitino through inflaton decay, both the $\pm 3/2$ and the $\pm 1/2$ components of the gravitino populate the Universe, whereas in the case of thermal production [Eq. (9)] only the longitudinal Goldstino component contributes to the relic abundance.

Perspectives and conclusions.—In many ways, it seems quite natural that a particle with only gravitational interactions should make up the dark matter of the Universe. We have seen that in the generic context with gravitino dark matter where the supersymmetric particle spectrum thermalizes with standard model bath, an upper limit to the mass of the gravitino of ≈ 4 TeV is obtained. However, if

one makes the minimal hypothesis that the supersymmetric spectrum lies above the inflaton mass, a new cosmologically allowed window opens for gravitino mass above 0.2 EeV. Indeed, despite the weakness of its coupling, the gravitino can be produced directly from the thermal bath by the exchange of virtual heavy superpartners (or, equivalently, by higher dimensional operators). It can also be produced directly from the inflaton decay. In order to obtain gravitino dark matter from the thermal bath, we predict a relatively large reheating temperature $\gtrsim 10^{10}$ GeV, compatible with the thermal leptogenesis scenario. If stable, this gravitino is virtually undetectable as it is the only R -parity odd state ever present in the Universe after inflation. If unstable through an R -parity violating coupling, the decay of the gravitino would produce EeV-like monochromatic photons or neutrinos, which are not yet observable by present experiments.

The authors acknowledge Karim Benakli and Yifan Chen for useful discussions. This work was supported by the France-US PICS No. 06482. Y. Mambrini acknowledges partial support from the European Union's Horizon 2020 research and innovation programme (under the Marie Skłodowska-Curie Grant Agreements No. 690575 and No. 674896), and the ERC advanced Grants Higgs@LHC. E.D. acknowledges partial support from the ANR Black-dS-String. The work of K. A. O. was supported in part by DOE Grant No. DE-SC0011842 at the University of Minnesota.

*Emilian.Dudas@cpht.polytechnique.fr

†yann.mambrini@th.u-psud.fr

‡olive@physics.umn.edu

- [1] G. Aad *et al.* (ATLAS Collaboration), *J. High Energy Phys.* **09** (2014) 176; **10** (2015) 054; *Phys. Rev. D* **94**, 032003 (2016); S. Chatrchyan *et al.* (CMS Collaboration), *J. High Energy Phys.* **06** (2014) 055; V. Khachatryan *et al.* (CMS Collaboration), *J. High Energy Phys.* **10** (2016) 006.
- [2] E. Aprile *et al.* (XENON100 Collaboration), *Phys. Rev. Lett.* **109**, 181301 (2012).
- [3] D. S. Akerib *et al.* (LUX Collaboration), *Phys. Rev. Lett.* **118**, 021303 (2017).
- [4] C. Fu *et al.* (PandaX-II Collaboration), *Phys. Rev. Lett.* **118**, 071301 (2017).
- [5] O. Buchmueller, R. Cavanaugh, A. De Roeck, J. R. Ellis, H. Flaecher, S. Heinemeyer, G. Isidori, K. A. Olive, F. J. Ronga, and G. Weiglein, *Eur. Phys. J. C* **64**, 391 (2009).
- [6] E. A. Bagnaschi *et al.*, *Eur. Phys. J. C* **75**, 500 (2015).
- [7] J. Ellis, J. L. Evans, F. Luo, N. Nagata, K. A. Olive, and P. Sandick, *Eur. Phys. J. C* **76**, 8 (2016); K. A. Olive, *Proc. Sci.*, PLANCK2015 (2015) 093 [arXiv:1510.06412].
- [8] J. D. Wells, arXiv:hep-ph/0306127; N. Arkani-Hamed and S. Dimopoulos, *J. High Energy Phys.* **06** (2005) 073; G. F. Giudice and A. Romanino, *Nucl. Phys.* **B699**, 65 (2004); **B706**, 487 (2005); N. Arkani-Hamed, S. Dimopoulos, G. F. Giudice, and A. Romanino, *Nucl. Phys.* **B709**, 3 (2005); J. D. Wells, *Phys. Rev. D* **71**, 015013 (2005).
- [9] P. J. Fox, D. E. Kaplan, E. Katz, E. Poppitz, V. Sanz, M. Schmaltz, M. D. Schwartz, and N. Weiner, arXiv:hep-th/0503249; L. J. Hall and Y. Nomura, *J. High Energy Phys.* **03** (2010) 076.
- [10] H. Pagels and J. R. Primack, *Phys. Rev. Lett.* **48**, 223 (1982).
- [11] D. V. Nanopoulos, K. A. Olive, and M. Srednicki, *Phys. Lett. B* **127**, 30 (1983).
- [12] J. R. Ellis, J. S. Hagelin, D. V. Nanopoulos, K. A. Olive, and M. Srednicki, *Nucl. Phys.* **B238**, 453 (1984).
- [13] K. A. Olive, D. N. Schramm, and M. Srednicki, *Nucl. Phys.* **B255**, 495 (1985).
- [14] J. R. Ellis, K. A. Olive, Y. Santoso, and V. C. Spanos, *Phys. Lett. B* **588**, 7 (2004).
- [15] J. L. Feng, S. Su, and F. Takayama, *Phys. Rev. D* **70**, 063514 (2004); **70**, 075019 (2004).
- [16] F. D. Steffen, *J. Cosmol. Astropart. Phys.* **09** (2006) 001.
- [17] W. Buchmuller, L. Covi, K. Hamaguchi, A. Ibarra, and T. Yanagida, *J. High Energy Phys.* **03** (2007) 037; W. Buchmuller, *AIP Conf. Proc.* **1200**, 155 (2010).
- [18] S. Bailly, K. Y. Choi, K. Jedamzik, and L. Roszkowski, *J. High Energy Phys.* **05** (2009) 103.
- [19] L. Covi, J. Hasenkamp, S. Pokorski, and J. Roberts, *J. High Energy Phys.* **11** (2009) 003.
- [20] P. Fayet, Phenomenology of supersymmetry, in *Proceedings of 17th Rencontres de Moriond* (1982), C82-03-14, p. 483; S. Weinberg, *Phys. Rev. Lett.* **48**, 1303 (1982); J. R. Ellis, A. D. Linde, and D. V. Nanopoulos, *Phys. Lett. B* **118**, 59 (1982); L. M. Krauss, *Nucl. Phys.* **B227**, 556 (1983); M. Y. Khlopov and A. D. Linde, *Phys. Lett. B* **138**, 265 (1984); M. Kawasaki, F. Takahashi, and T. T. Yanagida, *Phys. Rev. D* **74**, 043519 (2006).
- [21] J. L. Evans, M. A. G. Garcia, and K. A. Olive, *J. Cosmol. Astropart. Phys.* **03** (2014) 022.
- [22] T. Moroi, M. Yamaguchi, and T. Yanagida, *Phys. Lett. B* **342**, 105 (1995).
- [23] M. Kawasaki, T. Moroi, and T. Yanagida, *Phys. Lett. B* **370**, 52 (1996).
- [24] D. Lindley, *Astrophys. J.* **294**, 1 (1985); J. R. Ellis, D. V. Nanopoulos, and S. Sarkar, *Nucl. Phys.* **B259**, 175 (1985); M. Kawasaki and T. Moroi, *Prog. Theor. Phys.* **93**, 879 (1995); M. Kawasaki, K. Kohri, and T. Moroi, *Phys. Rev. D* **63**, 103502 (2001); R. H. Cyburt, J. Ellis, B. D. Fields, and K. A. Olive, *Phys. Rev. D* **67**, 103521 (2003); J. R. Ellis, K. A. Olive, and E. Vangioni, *Phys. Lett. B* **619**, 30 (2005); D. G. Cerdeno, K. Y. Choi, K. Jedamzik, L. Roszkowski, and R. Ruiz de Austri, *J. Cosmol. Astropart. Phys.* **06** (2006) 005; K. Jedamzik, K. Y. Choi, L. Roszkowski, and R. Ruiz de Austri, *J. Cosmol. Astropart. Phys.* **07** (2006) 007; S. Bailly, K. Jedamzik, and G. Moulta, *Phys. Rev. D* **80**, 063509 (2009); K. Jedamzik and M. Pospelov, *New J. Phys.* **11**, 105028 (2009); M. Pospelov and J. Pradler, *Annu. Rev. Nucl. Part. Sci.* **60**, 539 (2010).
- [25] D. V. Volkov and V. P. Akulov, *Phys. Lett.* **46B**, 109 (1973); for applications of nonlinear supersymmetry to inflation, see, e.g., I. Antoniadis, E. Dudas, S. Ferrara, and A. Sagnotti, *Phys. Lett. B* **733**, 32 (2014); S. Ferrara, R. Kallosh, and A. Linde, *J. High Energy Phys.* **10** (2014) 143; G. Dall'Agata and F. Zwirner, *J. High Energy Phys.* **10** (2014) 172.

- [26] M. Kawasaki, K. Kohri, and T. Moroi, *Phys. Rev. D* **71**, 083502 (2005).
- [27] R. H. Cyburt, J. Ellis, B. D. Fields, F. Luo, K. A. Olive, and V. C. Spanos, *J. Cosmol. Astropart. Phys.* **10** (2009) 021; **10** (2010) 032; **05** (2013) 014.
- [28] R. H. Cyburt, J. R. Ellis, B. D. Fields, K. A. Olive, and V. C. Spanos, *J. Cosmol. Astropart. Phys.* **11** (2006) 014; R. H. Cyburt, J. Ellis, B. D. Fields, F. Luo, K. A. Olive, and V. C. Spanos, *J. Cosmol. Astropart. Phys.* **12** (2012) 037.
- [29] J. Pradler and F. D. Steffen, *Phys. Lett. B* **666**, 181 (2008).
- [30] J. L. Feng, A. Rajaraman, and F. Takayama, *Phys. Rev. D* **68**, 063504 (2003); J. L. Feng, S. F. Su, and F. Takayama, *Phys. Rev. D* **70**, 063514 (2004).
- [31] M. Kawasaki, K. Kohri, T. Moroi, and A. Yotsuyanagi, *Phys. Rev. D* **78**, 065011 (2008).
- [32] P. A. R. Ade *et al.* (Planck Collaboration), *Astron. Astrophys.* **594**, A13 (2016).
- [33] S. Profumo and C. E. Yaguna, *Phys. Rev. D* **69**, 115009 (2004); I. Gogoladze, R. Khalid, and Q. Shafi, *Phys. Rev. D* **79**, 115004 (2009); D. Feldman, Z. Liu, and P. Nath, *Phys. Rev. D* **80**, 015007 (2009); M. Adeel Ajaib, T. Li, Q. Shafi, and K. Wang, *J. High Energy Phys.* **01** (2011) 028; K. Harigaya, M. Ibe, and T. T. Yanagida, *J. High Energy Phys.* **12** (2013) 016; K. Harigaya, K. Kaneta, and S. Matsumoto, *Phys. Rev. D* **89**, 115021 (2014); M. Low and L. T. Wang, *J. High Energy Phys.* **08** (2014) 161; J. L. Evans and K. A. Olive, *Phys. Rev. D* **90**, 115020 (2014); S. Raza, Q. Shafi, and C. S. Ün, *Phys. Rev. D* **92**, 055010 (2015).
- [34] A. De Simone, G. F. Giudice, and A. Strumia, *J. High Energy Phys.* **06** (2014) 081.
- [35] J. Ellis, F. Luo, and K. A. Olive, *J. High Energy Phys.* **09** (2015) 127.
- [36] J. Ellis, J. L. Evans, F. Luo, and K. A. Olive, *J. High Energy Phys.* **02** (2016) 071.
- [37] J. Ellis, D. V. Nanopoulos, and K. A. Olive, *Phys. Rev. Lett.* **111**, 111301 (2013); **111**, 129902 (2013); *J. Cosmol. Astropart. Phys.* **10** (2013) 009; J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, and K. A. Olive, *Classical Quantum Gravity* **33**, 094001 (2016).
- [38] A. A. Starobinsky, *Phys. Lett.* **91B**, 99 (1980).
- [39] S. Deser and B. Zumino, *Phys. Rev. Lett.* **38**, 1433 (1977).
- [40] J. Polonyi, Hungary Central Institute Research Report No. KFKI-1977-93 (1977).
- [41] G. Arcadi, M. Dutra, P. Ghosh, M. Lindner, Y. Mambrini, M. Pierre, S. Profumo, and F. S. Queiroz, [arXiv:1703.07364](https://arxiv.org/abs/1703.07364).
- [42] G. Arcadi, Y. Mambrini, and F. Richard, *J. Cosmol. Astropart. Phys.* **03** (2015) 018; J. Kearney, N. Orlofsky, and A. Pierce, *Phys. Rev. D* **95**, 035020 (2017).
- [43] O. Lebedev and Y. Mambrini, *Phys. Lett. B* **734**, 350 (2014); G. Arcadi, Y. Mambrini, M. H. G. Tytgat, and B. Zaldivar, *J. High Energy Phys.* **03** (2014) 134.
- [44] L. J. Hall, K. Jedamzik, J. March-Russell, and S. M. West, *J. High Energy Phys.* **03** (2010) 080; X. Chu, T. Hambye, and M. H. G. Tytgat, *J. Cosmol. Astropart. Phys.* **05** (2012) 034; X. Chu, Y. Mambrini, J. Quevillon, and B. Zaldivar, *J. Cosmol. Astropart. Phys.* **01** (2014) 034.
- [45] Y. Mambrini, K. A. Olive, J. Quevillon, and B. Zaldivar, *Phys. Rev. Lett.* **110**, 241306 (2013).
- [46] M. Bolz, A. Brandenburg, and W. Buchmuller, *Nucl. Phys.* **B606**, 518 (2001); **B790**, 336(E) (2008); J. Pradler and F. D. Steffen, *Phys. Rev. D* **75**, 023509 (2007); V. S. Rychkov and A. Strumia, *Phys. Rev. D* **75**, 075011 (2007).
- [47] J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, K. A. Olive, and M. Peloso, *J. Cosmol. Astropart. Phys.* **03** (2016) 008.
- [48] J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, and K. A. Olive, *J. Cosmol. Astropart. Phys.* **10** (2015) 003.
- [49] K. Benakli, Y. Chen, E. Dudas, and Y. Mambrini, *Phys. Rev. D* **95**, 095002 (2017).
- [50] G. F. Giudice, A. Notari, M. Raidal, A. Riotto, and A. Strumia, *Nucl. Phys.* **B685**, 89 (2004).
- [51] J. Heisig, *J. Cosmol. Astropart. Phys.* **04** (2014) 023.
- [52] J. Pradler and F. D. Steffen, *Phys. Lett. B* **648**, 224 (2007).
- [53] M. Endo, K. Kadota, K. A. Olive, F. Takahashi, and T. T. Yanagida, *J. Cosmol. Astropart. Phys.* **02** (2007) 018.
- [54] M. Kawasaki, M. Yamaguchi, and T. Yanagida, *Phys. Rev. Lett.* **85**, 3572 (2000).
- [55] Y. Ema, K. Mukaida, K. Nakayama, and T. Terada, *J. High Energy Phys.* **11** (2016) 184.