

Gravitino dark matter with trilinear couplings to explain the anomalous positron fraction

Johannes Buchner,^a Edson Carquin,^a Marco A. Díaz,^a
Germán A. Gómez-Vargas,^{a,b} Boris Panes,^c Nicolás Viaux^a

^aInstituto de Física, Pontificia Universidad Católica de Chile, Avenida Vicuña Mackenna 4860, Santiago, Chile

^bIstituto Nazionale di Fisica Nucleare, Sezione di Roma “Tor Vergata”, I-00133 Roma, Italy

^cInstituto de Física, Universidade de São Paulo, R. do Matão 187, São Paulo, SP, 05508-900, Brazil

E-mail: jbuchner@astro.puc.cl, edson.carquin@fis.puc.cl, mad@susy.fis.puc.cl,
ggomezv@uc.cl, bapanes@if.usp.br, nviaux@fis.puc.cl

Abstract. The flux of charged leptons measured by the space-based detectors PAMELA, AMS-02, CALET, DAMPE and Fermi-LAT presents anomalous behavior as energy increase. In particular AMS-02 observations provide compelling evidence for a new source of positrons and electrons. Its origin is still unknown, but standard and successful scenarios include the contribution of dark matter and unresolved astrophysical sources, such as nearby pulsars. On the one hand, it has been shown that explanations based mostly on dark matter emission tend to overproduce gamma-rays, entering in conflict with measurements of the extra-galactic gamma-ray background (EGB). On the other hand, explanations using pulsars have shown good performance to solve the data, but these ideas are still under scrutiny. In general, a mixture of sources would signal a good approach for future research. However, before going deeply into that research direction, in this work we would like to present a dark matter scenario, which contains a gravitino as dark matter candidate with trilinear couplings to standard model particles, that is able to solve both the lepton data as well as the bounds from the EGB. Besides, it also offers some connections to other observables such as neutrino physics. These results may serve as a probe of concept for the dark matter only explanation of the positron anomaly, but more conservatively speaking it could motivates the use of this model for future mixed searches.

Keywords: dark matter experiments, cosmic ray experiments, gamma ray experiments, dark matter theory

1 Introduction

The continuous and systematic analysis of the data collected by cosmic-ray detectors is one of the most direct ways to check and test the predictions of particle physics models that contain a candidate for dark matter (DM) which can annihilate or decay to Standard Model (SM) particles. In particular, we suggest [1] for a review about this subject focused on the gravitino dark matter scenario and [2] for an updated and more general discussion.

Furthermore, this discussion becomes specially relevant since contemporary experiments show the presence of persistent and striking anomalies concerning the detection of cosmic-rays. For instance, the energy spectrum of electrons and positrons measured by experiments such as AMS [3–5], CALET [6], DAMPE [7] and Fermi-LAT [8] show noticeable discrepancies when compared to predictions based on standard astrophysical sources, such as cosmic-ray interactions or emissions from pulsars. Therefore, these experimental signals strongly suggest that extra sources of (primary) positrons are required in order to make sense of the data.

It has been shown that the positron anomaly measured by several experiments, from PAMELA to AMS, can be well explained by considering a population of DM in agreement with standard density profiles, such as NFW, that can annihilate or decay to charged leptons both as primary or secondary particles. This picture, however, seem to be in conflict with the measurements of the extra galactic gamma-ray background (EGB) derived from Fermi-LAT and other gamma-ray detectors [9–14]. From a revision of the latest literature concerning this subject, we can conclude that the explanation of the positron excess would necessarily require some modifications of the DM density profile properties or the contribution of astrophysical sources with particular properties, such as near pulsars with proper diffusion coefficients [Refs to HAWK and Hooper](#).

Indeed, in a previous work, we have checked that a DM scenario considering a gravitino as the lightest super-symmetric particle, that is able to decay to SM particles only through bilinear R-parity violation (BRpV) couplings is in serious conflicts with EGB limits [Ref to us](#). Furthermore, the parameter space that prefers the explanation of AMS positron fraction is not useful to explain any of the neutrino masses or mixing angles, which is one of the big motivations to introduce BRpV to start with. This is one more piece of information that may indicate that other solutions should be considered in order to explain the positron anomaly.

In order to reduce the troubles to explain the cosmic-ray anomalous data mostly with DM contributions, here we would like to argue that the gravitino scenario could be strongly alleviated if we simply move from bilinear to trilinear RpV couplings. In this scenario we can consistently explain the data from AMS, CALET and the EGB limits derived from Fermi-LAT data. However, when we include the data from DAMPE the things get tougher again. Nonetheless, in this model we also can connect the life-time and branching fractions of gravitino decays that are able to explain the positron anomaly to the scale of neutrino physics. Ultimately, we acknowledge that this model probably should be complemented in order to properly account for cosmic-ray observations and neutrino physics.

The paper is organized as follows, in the first section we discuss the data from different cosmic ray detectors that we consider for our analysis. In the second section we introduce our model of dark matter. In the third section we compare trilinear and bilinear observables that show why we can expect some improvement of the gravitino DM scenario. In the fourth section we present the results of our statistical analysis and consequences. Finally, in the fifth section we conclude and elaborate about future prospects regarding these issues.

2 Data resources

Motivated by the anomalous measurement of the positron fraction done by experiments such as AMS [3], in this work we study a model of dark matter that try to accomodate old and recent data concerning the detection of electrons and positrons, which are originated in astrophysical environments. Also we use the measurements of gamma-rays in order to check the results of our models. In particular, we consider the following data releases:

- D_1 : The positron fraction measured by AMS between 0.5 and 500 GeV [3],
- D_2 : The independent measurement of the electron and positron fluxes by AMS between 0.5 and 700 GeV [4],
- D_3 : The measurement of the sum of electron and positron spectrum measured by AMS between 0.5 GeV and 1 TeV [5],
- D_4 : The extended measurement of the sum of electron and positron spectrum by CALET between 11 GeV and 4.8 TeV [6],
- D_5 : The direct detection of the spectrum of electrons plus positrons measured by DAMPE between 25 GeV and 4.6 TeV [7] and
- D_6 : The spectrum of isotropic diffuse gamma-ray emission between 100 MeV and 820 GeV measured by Fermi-LAT [8], from which we derive the limits on the EGB.

where we have used the symbol D_i to identify each dataset, which is going to be used in the section that discuss the statistical analysis of these datasets once we define our model.

3 Gravitino model and effective decay channels

In order to define our dark matter scenario we consider a super-symmetric (SUSY) extension of the SM with a low energy spectrum characterized by a gravitino as the lightest SUSY particle (LSP), which is able to decay to standard model particles only through trilinear leptonic R-parity violating couplings. We suggest to follow [1, 15] for details about the superpotential and lagrangian formulation of the complete model, the corresponding Feynman diagrams and computations of related observables.

In this scenario, the decay of the gravitino LSP can be achieved in two steps. Initially, we have the R-Parity conserved interactions between the gravitino, one SM fermion and the corresponding scalar super-partner, which in principle do not allow the direct decay of the gravitino. In order to allow these decays we may consider the RpV interactions between the scalar super-partners and pairs of SM fermions.

The latter part is what we want to highlight here, since this contains the trilinear couplings that ultimately determine the channels allowed for gravitino decays. The superpotential that generate the RpV interactions can be written in terms of the left-handed superfields for the leptons (L), quarks (Q) and Higgs of hypercharge 1/2 (H) and the right-handed superfields for the charged leptons (E^c), up and down type quarks (U^c , D^c). In practice, this superpotential is given by the following expression,

$$W_{\mathcal{RP}} = \sum_{ijk} \left(\frac{1}{2} \lambda_{ijk} L_i L_j E_k^c + \lambda'_{ijk} L_i Q_j D_k^c + \lambda''_{ijk} U_i D_j D_k^c + \mu_i H L_i \right) \quad (3.1)$$

Channel	Final State	Details	Acronym
c1	$e^+\mu^-\nu$	antielectron-muon-neutrino	AEMuNue
c2	$e^+\tau^-\nu$	antielectron-tau-neutrino	AETauNue
c3	$e^-\mu^+\nu$	electron-antimuon-neutrino	EAMuNue
c4	$\mu^-\mu^+\nu$	muon-antimuon-neutrino	MuAMuNue
c5	$\tau^-\tau^+\nu$	tau-antitau-neutrino	TauATauNue
c6	$\tau^-\mu^+\nu$	tau-antimuon-neutrino	TauAMuNue
c7	$\tau^+\mu^-\nu$	antitau-muon-neutrino	ATauMuNue
c8	$e^-e^+\nu$	electron-antielectron-neutrino	EAENue
c9	$e^-\tau^+\nu$	electron-antitau-neutrino	EATauNue

Table 1. Independent channels considering prompt final states. Notice that we use ν to indicate any flavor of neutrinos.

i,j,k being flavor indices, λ_{ijk} , λ'_{ijk} , λ''_{ijk} dimensionless coupling constants and μ_i dimension one parameters. Basically, for our analysis we just consider the first term of the potential since we focus on the purely leptonic decays of the gravitinos.

This scenario can be compatible with the measurements of the large hadron collider by considering a gravitino mass above 1-2 TeV [Refs.](#) In [\[15\]](#) we can also find the analytical expressions for the gravitino decay associated to each individual trilinear coupling, which we are going to use in [section 7](#) to find a relationship between the gravitino life-time required to fit the positron anomaly and the scale of neutrino masses.

For the computation of the electron-positron spectrum at Earth we consider an approach similar to our previous work [\[16\]](#), therefore we suggest to follow this work and references therein for the computations of the total flux including propagation effects (we restrict the current analysis to the MED case only). On the other hand, we need to explain some details about the particular final state channels that appear in our scenario. In practice we can start by considering only the gravitino final state channels that in principle could produce different spectra of electrons and positrons, which are given in [table 1](#). We can notice that each channel can contain any neutrino flavor [This is not critical but we need to check the validity of the statement about the neutrino flavors](#), since the final state is equivalent, therefore we can see that the relation between branching fractions and trilinear couplings is not necessarily direct.

Considering the effective channels given in [table 1](#), we can model the amount of electrons, positrons, or γ rays, labeled as η , produced by gravitino decay as:

$$\Phi_G^\eta(E) = \frac{1}{m_G \tau_G} \sum_{j=1}^9 Br_j \frac{dN_j^\eta}{dE} D_{\text{factor}}^\eta, \quad (3.2)$$

with m_G and τ_G the mass and lifetime of the gravitino respectively. The term $\eta = e, p, \gamma$ for electron, positron or gamma-ray flux correspondingly. The D_{factor}^η is proportional to the density of dark matter in the case of $\eta = \gamma$, in the other cases is a more complex term that depends on the dark matter density and the propagation of charged particles in the Galaxy. The term $\frac{dN_j^\eta}{dE}$ is the amount of electrons, positrons, or gamma-rays per energy produced by decay of a gravitino per energy and propagated at the Earth position.

3.1 Electron-positron spectrum

Now, let us focus on the spectrum of charged leptons. In principle, we can get each Br_j as a function of the free parameters of our model, such as the trilinear couplings λ_{ijk} or the mass of scalars [at some point we have to do this computation, for which we suggest to follow hep-ph/0107286](#), but in general we can consider the branching fractions as the effective free parameters for the fit of charged lepton measurements with the condition that $\sum_i Br_i = 1$.

Interestingly, we can still reduce a bit more the number of degrees of freedom of our model by considering that some of these final states generate the same spectrum of electrons (positrons). [Here we could extend a bit more if necessary since we have all the material to back this assumptions](#). Thus, we can group the decay channels as follows,

$$\begin{aligned}\Phi_G^e(E) &\propto \frac{1}{m_G \tau_G} \left[(Br_1 + Br_4 + Br_7) \frac{dN_1^e}{dE} + (Br_2 + Br_5 + Br_6) \frac{dN_2^e}{dE} + \right. \\ &\quad \left. (Br_3 + Br_8 + Br_9) \frac{dN_3^e}{dE} \right] \\ \Phi_G^e(E) &\propto \frac{1}{m_G \tau_G} \left[\alpha_1 \frac{dN_1^e}{dE} + \alpha_2 \frac{dN_2^e}{dE} + \alpha_3 \frac{dN_3^e}{dE} \right]\end{aligned}$$

where $\alpha_1 = Br_1 + Br_4 + Br_7$, $\alpha_2 = Br_2 + Br_5 + Br_6$ and $\alpha_3 = Br_3 + Br_8 + Br_9$ with $\alpha_1 + \alpha_2 + \alpha_3 = 1$. Thus, we just need to define two independent effective branching fractions for the fit of electrons. Similarly, for the positron spectrum we have that

$$\begin{aligned}\Phi_G^p(E) &\propto \frac{1}{m_G \tau_G} \left[(Br_1 + Br_2 + Br_8) \frac{dN_1^p}{dE} + (Br_3 + Br_4 + Br_6) \frac{dN_3^p}{dE} + \right. \\ &\quad \left. (Br_5 + Br_7 + Br_9) \frac{dN_3^p}{dE} \right]\end{aligned}$$

Besides, we can use some equivalences between branching fractions of conjugated decay channels, $Br_1 = Br_3$, $Br_2 = Br_9$ and $Br_6 = Br_7$ to rewrite the positron spectrum as

$$\begin{aligned}\Phi_G^p(E) &\propto \frac{1}{m_G \tau_G} \left[(Br_9 + Br_3 + Br_8) \frac{dN_1^p}{dE} + (Br_1 + Br_4 + Br_7) \frac{dN_3^p}{dE} + \right. \\ &\quad \left. (Br_2 + Br_5 + Br_6) \frac{dN_5^p}{dE} \right] \\ \Phi_G^p(E) &\propto \frac{1}{m_G \tau_G} \left[\alpha_1 \frac{dN_3^p}{dE} + \alpha_2 \frac{dN_5^p}{dE} + \alpha_3 \frac{dN_1^p}{dE} \right]\end{aligned}$$

Finally, we can use that the electron spectrum from a given channel must be equal to the positron spectrum of the conjugated one to find that

$$\Phi_G^p(E) \propto \frac{1}{m_G \tau_G} \left[\alpha_1 \frac{dN_1^e}{dE} + \alpha_2 \frac{dN_2^e}{dE} + \alpha_3 \frac{dN_3^e}{dE} \right] \quad (3.3)$$

$$\Phi_G^p(E) = \Phi_G^e(E) \quad (3.4)$$

Therefore, for the fit of AMS-02, CALET or DAMPE we just need two α 's and three independent spectra. Furthermore, we get automatically the electron-positron symmetry for (gravitino) dark matter decays which is expected from general arguments considering charge conjugation symmetry.

3.2 Gamma-ray spectrum

Analogously to the electron-positron flux, we may discuss the total contribution of gravitino decays to the EGB measured at earth by considering the following expression

$$\Phi_G^\gamma(E) \propto \frac{1}{m_G \tau_G} \sum_{i=1}^9 Br_i \frac{dN_i^\gamma}{dE} \quad (3.5)$$

In this case we are not going to exploit the potential similarities between the gamma-ray energy spectra arising from different decay channels, if they exist at all. Instead, we are going to use the results of the previous section to find scenarios where the gamma-ray spectrum is indeed minimized, in order to maximize the chance to be compatible with the EGB measurements.

As we only can fix α_1 , α_2 and α_3 from the fit of charged leptons we have some freedom to choose the individual branching fractions to generate the photon flux. Also we must notice that for the photon spectrum we do not have coincidences between the spectrum of different channels such as $e^+ \mu^- \nu$ and $e^+ \tau^- \nu$, as we had for positrons. Therefore we are free to choose BR_1 to BR_9 with the conditions that

$$\begin{aligned} \alpha_1 &= Br_1 + Br_4 + Br_7 \\ \alpha_2 &= Br_2 + Br_5 + Br_6 \\ \alpha_3 &= Br_3 + Br_8 + Br_9 \end{aligned}$$

In order to decrease the possible number of photons to be produced we may choose Br_i in the following way,

$$\begin{aligned} BR_4 &= \alpha_1, \quad Br_1 = BR_3 = 0 \\ BR_5 &= \alpha_2, \quad Br_2 = BR_9 = 0 \\ BR_8 &= \alpha_3, \quad Br_6 = BR_7 = 0 \end{aligned}$$

which can be justified from the analysis of the corresponding gamma-ray spectra obtained for the corresponding channels. Basically, we prioritize the channels that produce the least amount of photons per gravitino decay. [Maybe, this choice needs further justification but in principle it has allowed us to find compatible points, which can be seen from the plots shown in the analysis part.](#)

4 Statistical data analysis

We fit measurements related to electron and positron fluxes at Earth, so we define backgrounds for each of these fluxes with power laws:

$$\Phi_B^p(E) = C_p E^{-\gamma_p}, \quad (4.1)$$

for positrons, and for electrons:

$$\Phi_B^e(E) = C_e E^{-\gamma_e}. \quad (4.2)$$

It has been shown that is not possible to reproduce the rise in electron and positron flux above ≈ 200 GeV modeling the fluxes with decreasing power laws. Therefore, we must include a source term injecting electron and positrons at high energies. Our source term candidate is the decay of gravitino dark matter into standard model particles, whose general expression is given in Eq. (3.2). It is worth noticing that gravitino decay yields equal amounts of electron and positron, as shown in Eq. (3.4).

Considering the definition of our background model and the analysis of the gravitino sector of the previous section, we end with eight free parameters that we need to fix to fully determine signal and background for electron and positron fluxes. Therefore, in order to compare with data, for instance positron measurements $\Phi_D^p(E_i)$, we define the following likelihood form:

$$\log \mathcal{L}_{\text{Positrons}} = -\frac{1}{2} \sum_i \left(\frac{(\Phi_D^p(E_i) - \Phi_M^p(\theta_p, E_i))^2}{(\sigma_D^2 + j \times \Phi_D^{p2}(\theta_p, E_i))} - \frac{1}{(\sigma_D^2 + j \times \Phi_D^{p2}(\theta_p, E_i))} \right), \quad (4.3)$$

with σ_D the statistical uncertainty of the measurement. The model is defined as:

$$\Phi_M^p(\theta, E) = \Phi_B^p(C_p, \gamma_p, E) + \Phi_G^p(m_G, \tau_G, \alpha_1, \alpha_2, E). \quad (4.4)$$

We introduce the parameter j in the likelihood to increase the total uncertainty as a fraction of the model, which allow us to account for possible systematic effects and correlations among the different data sets used in our following analysis. The likelihood functions considering other measurements can be defined analogously [Appendix or reference to the jupyter-notebook](#). In order to minimize the full likelihood, find the best fit values and confidence regions of our unknown parameters, we use the Bayesian inference Python package `PyMultinest` [17]. We analyze our datasets in four different cases by considering sets of 3 or 4 measurements. These cases are given by:

- Case 1: $D_1 + D_2$
- Case 2: $D_1 + D_2 + D_3$
- Case 3: $D_1 + D_2 + D_4$
- Case 4: $D_1 + D_2 + D_5$

In this way we are going to consider most of the data that we currently know about the detection of charged leptons. Also we are going to be able to study how each of them can affect the determination of gravitino and background parameters and their consequences to the comparison with the EGB measured by Fermi-LAT, which is our final target.

Best Fit	Case 1	Case 2	Case 3	Case 4
C_p [1/GeV cm ² s str]	14.90	14.74	14.93	14.37
γ_p	3.11	3.10	3.11	3.09
C_e [1/GeV cm ² s str]	426.10	421.77	422.08	422.67
γ_e	3.27	3.27	3.27	3.27
m_G [GeV]	1281	2274	3604	3751
τ_G [10 ²⁶ s]	4.61	3.59	2.27	2.29
α_1	0.03	0.36	0.15	0
α_2	0.54	0.58	0.82	0.68

Table 2. Best fit parameters for the different cases defined in previous section.

5 Results

The results of our statistical analysis are summarized in the following table 2 and the figures shown at the end of the section. In general we can see that the required life-time to fit the charged-lepton data is around 4×10^{26} s. This value is sufficiently high to predict a gamma-ray flux which is statistically compatible with the limits of the EGB. In this computation we have not considered the extragalactic component of gravitino decays or the inverse-compton mechanism, which will enhance the contribution to the total amount of gamma-rays produced by our candidate dark matter. By considering these issues we may suggest that this scenario can accomodate most but not all the anomalous signal in the charged lepton measurements and the rest should be supply by astrophysical components. Below, we details the results of our effective analysis for different choices of measured data.

5.1 $D_1 + D_2$

We find many points that can explain the lepton data and while not overshoot the γ -ray extragalactic background. With the bilinear model in previous work, we could not have that situation.

5.2 $D_1 + D_2 + D_3$

We get similar results than in the previous case, but adding the electron-positron sum from the same detector (in the 0.5 GeV to 1 TeV range PRL 113, 221102) may be a problem. Not entirely sure what it means, but the parameter related to this new measurement $\log(k)$ is significantly larger than the other three (see slide 8), maybe some correlation with the other data. An interpretation is that to produce the observed points we need to degrade the likelihood in the electron+positron data in a fraction of the model.

5.3 $D_1 + D_2 + D_4$

In this case we do not use the electron-positron sum from AMS02, but the values reported by CALET. The CALorimetric Electron Telescope is an instrument placed in the International Space Station, as well as AMS02. In arXiv:1712.01711 CALET team reported a measurement of the cosmic-ray electron + positron spectrum from 10 GeV to 3 TeV. The CALET spectrum is compatible with results from AMS02, but with Fermi and DAMPE reports. Including CALET data (but e+p AMS02) makes the fit to prefers higher gravitino masses and reduces the parameter space allowed for reproducing the lepton data while keeping the gamma-ray flux below the Fermi estimation.

5.4 $D_1 + D_2 + D_5$

We include here e+p spectrum from DAMPE. This measurement is in tension with AMS02. We find no points compatible with lepton and gamma-ray data.

6 Comparison between trilinear and bilinear observables

Here we want to support our previous results from a comparative analysis between TRpV and BRpV decay channels. The big difference between these scenarios is directly connected to the type of decay channels that are allowed for both cases.

In BRpV we only have access to two-body decays containing gauge bosons and leptons ($G \rightarrow Z\nu$ and $G \rightarrow W^\pm l^\mp$) or a higgs particle plus neutrinos ($G \rightarrow H\nu$). In particular, in our previous work we have shown that the preferred channel to explain the positron anomaly is given by $G \rightarrow W^\pm \tau^\mp$ which is one of the channels that produce most photons associated to final state radiation. Instead, in TRpV we just have access to direct three-body decays including two charged leptons plus a neutrino ($G \rightarrow l^\pm l^\mp \nu$) as shown in table 1. In the previous section we have shown that the preferred channels are given by $G \rightarrow \tau^\pm l^\mp \nu$, with l^\mp including the three flavours, but also the contributions of other channels are necessary.

Interestingly, from the analysis of the propagated spectrum of charged leptons associated to these channels, which are shown in Fig. (1), it is possible to see that the contribution to charged leptons in TRpV is enhanced in comparison to BRpV. In order to support this argument we have computed the total spectrum of charged leptons for both cases considering branching fractions that fit well the positron anomaly, but considering a normalized and common life-time. In simply words, in TRpV we need a lower rate of gravitino decays in comparison to BRpV to fit the positron anomaly, which automatically suggests that the contribution to photons could be reduced in the former case.




Figure 1. Charged lepton spectrum of trilinear (left) and bilinear (center) decay channels. In the last figure (right) we show the contribution to charged leptons from both cases considering branching fractions that fit well the positron anomaly.

Furthermore, it also can be shown that for the analogous exercise but now using the photon spectrum, which is shown in Fig. (2), the contribution to photons from TRpV is decreased in comparison to BRpV. Therefore, there are two factors that contribute to the

better performance of TRpV in comparison to BRpV. For a given gravitino decay rate, there are both an enhancement of the charged lepton flux and a reduction of the contribution to photons that confabulate to reduce the tensions with the EGB measurements.




Figure 2. Photon spectrum of trilinear (full top row) and bilinear (left bottom panel) decay channels. In the last figure (right bottom panel) we show the contribution to photons from both cases considering branching fractions that fit well the positron anomaly.

7 Discussion about the gravitino lifetime and neutrino masses

The full expressions for the gravitino decay width, considering trilinear R-Parity violation, are given in hep-ph/017286. For instance, from these expressions we can get an approximated formula for the leptonic decay $\Gamma(\tilde{G} \rightarrow \nu_i e_j \bar{e}_k)$ by assuming that the mass of the sleptons that mediate the three body decay are equal, such that $m_{\tilde{\nu}_{iL}} = m_{\tilde{e}_{jL}} = m_{\tilde{e}_{kR}} = \tilde{m}$, and expand in taylor series around the variable m_G/\tilde{m} to obtain

$$\Gamma(\tilde{G} \rightarrow \nu_i e_j \bar{e}_k) \approx \frac{1}{96(2\pi)^3} \frac{\lambda_{ijk}^2}{8M_\star^2} \frac{m_G^7}{\tilde{m}^4}, \quad (7.1)$$

where $M_\star = (8\pi G_N)^{1/2} = 2.4 \times 10^{18} \text{ GeV}$ is the reduced Planck mass. This result shows that the decay width (lifetime) decreases (increases) rapidly as we increase \tilde{m} , as expected. We




Figure 3. Gravitino life time in Trilinear RpV for $\lambda_{ijk} = 0.01$, $m_G = 1 \text{ TeV}$. For simplicity we use m_1, m_2, m_3 and m instead of $m_{\tilde{\nu}_{iL}}, m_{\tilde{e}_{jL}}, m_{\tilde{e}_{kR}}$ and \tilde{m} .

expect that a similar behavior should be obtained even when the mass of sleptons are not equal.

Indeed, we have verified this expectation numerically, by evaluating the full expression given in hep-ph/017286 using the maximum numerical precision in Mathematica. For instance, in Fig. 3 we plot the gravitino lifetime as a function of $m_{\tilde{\nu}_{iL}}$ for $m_{\tilde{e}_{jL}} = m_{\tilde{\nu}_{iL}}/2$ and $m_{\tilde{e}_{kR}} = m_{\tilde{\nu}_{iL}}/5$. Also, in the same figure we plot the lifetime derived from Eq. 7.1 evaluated at $\tilde{m} = m_{\tilde{\nu}_{iL}}/2$ in order to check that both approaches, exact computation and approximated formula, behave quite similarly.

Therefore, we can confidently derive the following expression for the gravitino lifetime,

$$\tau_G \approx 10^{26} \text{ s} \left(\frac{1}{\lambda_{ijk}\lambda_{ijk}} \right) \left(\frac{\tilde{m}}{2 \times 10^7 \text{ GeV}} \right)^4 \left(\frac{1 \text{ TeV}}{m_G} \right)^7 \quad (7.2)$$

where we have normalized with respect to 10^{26} s since this is the order of magnitude required by experiments such as AMS-02 and Fermi-LAT in order to fit the electron positron data in the first case or to avoid gamma ray constraints in the second.

In trilinear RpV the neutrino mass matrix receives contributions from 1-loop diagrams that contain both a charged lepton and the corresponding slepton. Indeed, we have derived the following (preliminary) expression

$$M_{ij}^{\nu(1)} \approx \frac{1}{16\pi^2} \sum_{gr} s_{\tilde{l}} c_{\tilde{l}} (\lambda_{igr} \lambda_{jrg} + \lambda_{jgr} \lambda_{irg}) m_g \ln \frac{m_{\tilde{l}_{r2}}^2}{m_{\tilde{l}_{r1}}^2}$$

where i and j are neutrino generation indices that run from 1 to 3. g is a charged lepton index that also run from 1 to 3, as well as r which is a slepton index. Thus, it can be seen that for order one $s_{\tilde{l}}$, $c_{\tilde{l}}$ and $\ln(m_{\tilde{l}_{r2}}^2/m_{\tilde{l}_{r1}}^2)$ we can get neutrino masses around the eV scale for $\lambda_{ijk} \approx 0.01$ even for $m_g \approx m_e$.

Indeed, by following the expressions given in hep-ph/0410242 for the contribution of λ' trilinear terms, we can get by analogy that the dominant term in the leptonic sector is

$$\begin{aligned} M_{ij}^{\nu(1)} &\approx \frac{1}{8\pi^2} \lambda_{i23} \lambda_{j32} \frac{m_\mu m_\tau A_\tau}{\tilde{m}^2} \\ &\approx 2 \times 10^{-2} \text{eV} \lambda_{i23} \lambda_{j32} \left(\frac{10^8 \text{ GeV}}{\tilde{m}} \right) \\ &\approx 2 \times 10^{-2} \text{eV} (\lambda_{i23} \lambda_{j32})^{1/4} \left(\frac{\tau_G}{10^{26} \text{ s}} \right)^{1/4} \left(\frac{m_G}{2 \text{ TeV}} \right)^{7/4} \end{aligned}$$

where A_τ is a free parameter that can be considered of order \tilde{m} , as it is done in hep-ph/0410242. Thus, if we consider this formula together with Eq. (7.2) we see that we can have contributions to the neutrino mass matrix of order 10^{-2} eV for trilinear couplings and a scalar mass which are compatible with $\tau_G \approx 10^{26}$ s.

Acknowledgments

The authors are thankful to Andrea Albert, Borut Bajc, Marco Cirelli, Michael Grefe, Luis Labarga, Carlos Munoz, Paolo Panci, Frank Steffen, and Gabriela Zaharijas for useful comments, and Marco Ajello for providing the EGB model contributions of Fig. (8). This work was supported by Conicyt Anillo grant ACT1102. GAGV thanks for the support of the Spanish MINECO's Consolider-Ingenio 2010 Programme under grant MultiDark CSD2009-00064 also the partial support by MINECO under grant FPA2012-34694. BP also thanks for the support of the State of São Paulo Research Foundation (FAPESP). The work of NV was supported by CONICYT FONDECYT/POSTDOCTORADO/3140559.

A Appendix

A.1 Results of the fit

In this section we show the plots concerning the statistical analysis of Cases 1 to 4, which are detailed in the main text. The black lines indicate the best fit curve, while the shade regions show 1 and 2 σ confidence level regions.




Figure 4. Best fit results for positron flux. The cases D_1 to D_4 are ordered from left to right and from top to bottom.




Figure 5. Best fit results for electron flux. The cases D_1 to D_4 are ordered from left to right and from top to bottom.




Figure 6. Best fit results for positron fraction. The cases D_1 to D_4 are ordered from left to right and from top to bottom.




Figure 7. Best fit results for electron plu positron flux. The cases D_1 to D_4 are ordered from left to right and from top to bottom.




Figure 8. Best fit results for photon flux compared to EGB limits. The cases D_1 to D_4 are ordered from left to right and from top to bottom.

References

- [1] M. Grefe, *Unstable Gravitino Dark Matter - Prospects for Indirect and Direct Detection*, [arXiv:1111.6779](#).
- [2] D. Hooper, *TASI Lectures on Indirect Searches For Dark Matter*, [arXiv:1812.0202](#).
- [3] **AMS** Collaboration, L. Accardo et al., *High Statistics Measurement of the Positron Fraction in Primary Cosmic Rays of 0.5–500 GeV with the Alpha Magnetic Spectrometer on the International Space Station*, *Phys. Rev. Lett.* **113** (2014) 121101.
- [4] **AMS** Collaboration, M. Aguilar et al., *Electron and Positron Fluxes in Primary Cosmic Rays Measured with the Alpha Magnetic Spectrometer on the International Space Station*, *Phys. Rev. Lett.* **113** (2014) 121102.
- [5] **AMS** Collaboration, M. Aguilar et al., *Precision Measurement of the $(e^+ + e^-)$ Flux in Primary Cosmic Rays from 0.5 GeV to 1 TeV with the Alpha Magnetic Spectrometer on the International Space Station*, *Phys. Rev. Lett.* **113** (2014) 221102.
- [6] O. Adriani et al., *Extended Measurement of the Cosmic-Ray Electron and Positron Spectrum from 11 GeV to 4.8 TeV with the Calorimetric Electron Telescope on the International Space Station*, *Phys. Rev. Lett.* **120** (2018), no. 26 261102, [[arXiv:1806.0972](#)].
- [7] **DAMPE** Collaboration, G. Ambrosi et al., *Direct detection of a break in the teraelectronvolt cosmic-ray spectrum of electrons and positrons*, *Nature* **552** (2017) 63–66, [[arXiv:1711.1098](#)].
- [8] **The Fermi LAT** Collaboration, M. Ackermann et al., *The spectrum of isotropic diffuse gamma-ray emission between 100 MeV and 820 GeV*, [arXiv:1410.3696](#).
- [9] M. Grefe, *Neutrino signals from gravitino dark matter with broken R-parity*, [arXiv:1111.6041](#).
- [10] M. Cirelli, E. Moulin, P. Panci, P. D. Serpico, and A. Viana, *Gamma ray constraints on decaying dark matter*, *Phys. Rev. D* **86** (Oct., 2012) 083506, [[arXiv:1205.5283](#)].
- [11] S. Ando and K. Ishiwata, *Constraints on decaying dark matter from the extragalactic gamma-ray background*, [arXiv:1502.0200](#).
- [12] M. Laletin, *A no-go theorem for the dark matter interpretation of the positron anomaly*, *Frascati Phys. Ser.* **63** (2016) 7–12, [[arXiv:1607.0204](#)].
- [13] W. Liu, X.-J. Bi, S.-J. Lin, and P.-F. Yin, *Constraints on dark matter annihilation and decay from the isotropic gamma-ray background*, *Chin. Phys.* **C41** (2017), no. 4 045104, [[arXiv:1602.0101](#)].
- [14] K. Belotsky, R. Budaev, A. Kirillov, and M. Laletin, *Fermi-LAT kills dark matter interpretations of AMS-02 data. Or not?*, *JCAP* **1701** (2017), no. 01 021, [[arXiv:1606.0127](#)].
- [15] G. Moreau and M. Chemtob, *R-parity violation and the cosmological gravitino problem*, *Phys. Rev. D* **65** (2002) 024033, [[hep-ph/0107286](#)].
- [16] E. Carquin, M. A. Diaz, G. A. Gomez-Vargas, B. Panes, and N. Viaux, *Confronting recent AMS-02 positron fraction and Fermi-LAT extragalactic γ -ray background measurements with gravitino dark matter*, *Phys. Dark Univ.* **11** (2016) 1–10, [[arXiv:1501.0593](#)].
- [17] J. Buchner, A. Georgakakis, K. Nandra, L. Hsu, C. Rangel, M. Brightman, A. Merloni, M. Salvato, J. Donley, and D. Kocevski, *X-ray spectral modelling of the AGN obscuring region in the CDFS: Bayesian model selection and catalogue*, *Astron. Astrophys.* **564** (2014) A125, [[arXiv:1402.0004](#)].