

LSAPy: Land Suitability Analysis in Python

Baptiste Hamon ¹

¹ Department of Civil and Environmental Engineering, University of Canterbury, Christchurch, New Zealand

DOI: [10.xxxxxx/draft](https://doi.org/10.1002/xxxxx.draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

LSAPy is a highly customizable Python library designed to streamline and enhance Land Suitability Analysis (LSA) workflows. The package implements a fuzzy-logic approach and provides three core objects—SuitabilityFunction, SuitabilityCriteria, and LandSuitabilityAnalysis—that work together to deliver a flexible and user-defined LSA framework. By relying on xarray objects for computation ([Hoyer & Hamman, 2017](#)), LSAPy seamlessly integrates with the broader Python ecosystem, such as dask for efficient parallel processing and matplotlib for data visualisation. Its modular design addresses some limitations of existing LSA tools by offering greater flexibility, reproducibility, and scalability for research and practical applications.

Statement of need

In the past decades, several software programs have been developed to perform land evaluation or suitability analysis, including ALES ([Johnson & Cramb, 1991](#)), Micro-LEIS ([D. De La Rosa et al., 1992](#); [D. De La Rosa et al., 2004](#)), LEIGIS ([Kalogirou, 2002](#)), ALSE ([Elsheikh et al., 2013](#)) and general-purpose GIS platforms such as ArcGIS and QGIS. While each of these tools offers distinct advantages, a limitation that often arises is the software-imposed constraints and the lack of freedom left to users ([Asaad et al., 2022](#); [Chen et al., 2022](#); [Elsheikh et al., 2013](#)). These limitations manifest in several ways. Desktop GIS solutions, for example, often present challenges related to operating system dependency, the cost of proprietary software (e.g., ArcGIS), and difficulties in integration with broader analytical frameworks ([Chen et al., 2022](#)). Furthermore, many specialised programs (e.g., Micro-LEIS, LEIGIS, ALSE) do not permit modification of the land characteristics used in the analysis ([Asaad et al., 2022](#); [Elsheikh et al., 2013](#)). This rigidity is problematic, as the predefined land characteristics may not apply to all crops or may require augmentation with additional parameters. Additionally, some tools (e.g., LEIGIS) support only a limited selection of crops for evaluation ([Elsheikh et al., 2013](#)).

Recently, two libraries – ALUES ([Asaad et al., 2022](#)) and PyLUSAT ([Chen et al., 2022](#)) – have been developed to address some of these limitations. ALUES, an R package, supports the evaluation of 56 crops and allows the addition of new ones. While it addresses some of the limitations discussed above, it falls short in others. For example, it relies on fixed criteria grouped into three unmodifiable categories ([Asaad et al., 2022](#)). PyLUSAT, a Python package, enables land suitability analysis using vector data, which offers specific advantages ([Chen et al., 2022](#)). Nevertheless, raster-based approaches are generally more efficient for data combinations and complex calculations [carr2007], particularly when integrating climate data, which is often distributed in netCDF format and best processed using raster routines. Converting such data to a vector format for analysis is technically feasible but suboptimal, leading to increased computational overhead and memory usage. Consequently, raster-based methods are essential for large-scale analyses, such as land suitability assessments using climate projections.

Beyond these technical constraints, the lack of user flexibility in existing software introduces

43 further limitations. First, when programs offer a restricted set of functionalities, they hinder the
44 reproducibility of scientific results. For instance, ALSE aggregates criteria suitability using the
45 maximum limitation method (Elsheikh et al., 2013), whereas ALUES provides only minimum,
46 maximum, and average aggregation options (Asaad et al., 2022). This discrepancy renders the
47 reproduction of cross-software results impossible. Second, tools designed for specific use cases
48 (e.g., agriculture) limit broader adoption and, in the case of free and open-source software
49 (FOSS), may impede community engagement and growth. Finally, rigid frameworks restrict
50 the ability to explore edge cases or out-of-the-box analyses, which are increasingly important
51 in a rapidly changing environment.

52 The limitations of existing software programs motivated the development of LSAPy (Land
53 Suitability Analysis in Python), a new, highly customizable Python package that supports
54 raster-like data. Thanks to its custom-based approach, frameworks used in most software
55 programs mentioned here can be implemented in LSAPy.

56 Key features

57 LSAPy provides three core objects that operate together to perform the land suitability analysis
58 (LSA) according to user-defined frameworks \autoref{fig:lsapy}.




Figure 1: Overview of LSAPy's object structures and their associated properties and methods. * is used as an abbreviation of .label{fig:lsapy}

59 Suitability Function

60 SuitabilityFunction is built around a function that transforms input data into suitability
61 values \autoref{fig:lsapy}. LSAPy includes built-in functions for both discrete and continuous
62 data. For the latter, the package follows a fuzzy-logic approach, implementing Gaussian-like
63 and sigmoid-like membership functions previously used in LSA studies. If the provided functions
64 do not meet user requirements, custom functions can be defined.

65 Suitability Criteria

66 The SuitabilityCriteria defines an individual criteria used in LSA. Its indicator prop-
67 erty refers to the input data, while func specifies the associated SuitabilityFunction
68 \autoref{fig:lsapy}. The weight and category properties allow users to customise how

each criteria is aggregated with others in the analysis. The `compute()` method applies the `SuitabilityFunction` to the given indicator to calculate the suitability score.

Land Suitability Analysis

`LandSuitabilityAnalysis` is the top-level class in LSAPy, defining the LSA framework. All criteria for the analysis are stored in the `criteria` property. The `run()` method executes the LSA, with parameters specifying the level of suitability to compute (i.e., criteria, category, or overall land suitability) and the aggregation method to use. Currently, supported aggregation methods include median, mean, weighted mean, geometric mean, weighted geometric mean, and limiting factor.

Additional Features

- The `stats` module offers functions for computing spatial (e.g., national, regional...) summary statistics of land use suitability.
- LSAPy's `open_data` function provides access to sample datasets, including soil/land data from NZGLID (Hamon, 2025) and climate data from NEX-GDDP-CMIP6 (Thrasher et al., 2022) datasets for tutorial and training purposes.

Research Applications

LSAPy has been used to assess the impact of climate change on apples, cherries, maize and wheat in New Zealand (Hamon, Quénol, et al., 2025a; Hamon, Quénol, et al., 2025b). Additionally, ongoing research investigating the influence of land suitability analysis criteria on land use suitability also utilises LSAPy (Hamon, Le Roux, et al., 2025).

Acknowledgements

The development of LSAPy began as part of a PhD funded by the Food Transition 2050 Joint Postgraduate School.

References

- Asaad, A.-A. B., Salvacion, A. R., & Yen, B. T. (2022). ALUES: R package for agricultural land use evaluation system. *Journal of Open Source Software*, 7(73), 4228. <https://doi.org/10.21105/joss.04228>
- Chen, C., Judge, J., & Hulse, D. (2022). PyLUSAT: An open-source python toolkit for GIS-based land use suitability analysis. *Environmental Modelling & Software*, 151, 105362. <https://doi.org/10.1016/j.envsoft.2022.105362>
- De La Rosa, D., Mayol, F., Diaz-Pereira, E., Fernandez, M., & De La Rosa, D. (2004). A land evaluation decision support system (MicroLEIS DSS) for agricultural soil protection. *Environmental Modelling & Software*, 19(10), 929–942. <https://doi.org/10.1016/j.envsoft.2003.10.006>
- De La Rosa, D., Moreno, J. A., Garcia, L. V., & Almorza, J. (1992). MicroLEIS: A microcomputer-based mediterranean land evaluation information system. *Soil Use and Management*, 8(2), 89–96. <https://doi.org/10.1111/j.1475-2743.1992.tb00900.x>
- Elsheikh, R., Mohamed Shariff, A. R. B., Amiri, F., Ahmad, N. B., Balasundram, S. K., & Soom, M. A. M. (2013). Agriculture land suitability evaluator (ALSE): A decision and planning support tool for tropical and subtropical crops. *Computers and Electronics in Agriculture*, 93, 98–110. <https://doi.org/10.1016/j.compag.2013.02.003>

- 110 Hamon, B. (2025). *New zealand gridded land information dataset (NZGLID)* (Version v1.1).
111 Zenodo. <https://doi.org/10.5281/ZENODO.16249350>
- 112 Hamon, B., Le Roux, R., Aubry, M., Quénol, H., Vannier, C., & Cochrane, T. A. (2025).
113 Influence of climate thresholds and crop phenology on land suitability. In *[Manuscript in*
114 *preparation]*.
- 115 Hamon, B., Quénol, H., Vannier, C., & Cochrane, T. (2025a). *Bridging the gaps in the*
116 *future agroclimatic suitability of crops in new zealand*. EGU general assembly 2025, vienna,
117 austria, 27 apr-2 may 2025, EGU25-1157. <https://doi.org/10.5194/egusphere-egu25-1157>
- 118 Hamon, B., Quénol, H., Vannier, C., & Cochrane, T. A. (2025b). Regional impacts of climate
119 change on new zealand's agriculture: Land suitability analysis of key crops. In *[Manuscript*
120 *submitted for publication]*.
- 121 Hoyer, S., & Hamman, J. (2017). Xarray: N-d labeled arrays and datasets in python. *Journal*
122 *of Open Research Software*, 5(1). <https://doi.org/10.5334/jors.148>
- 123 Johnson, A. K. L., & Cramb, R. A. (1991). Development of a simulation based land evaluation
124 system using crop modelling, expert systems and risk analysis. *Soil Use and Management*,
125 7(4), 239–246. <https://doi.org/10.1111/j.1475-2743.1991.tb00881.x>
- 126 Kalogirou, S. (2002). Expert systems and GIS: An application of land suitability evaluation.
127 *Computers, Environment and Urban Systems*, 26(2), 89–112. [https://doi.org/10.1016/S0198-9715\(01\)00031-X](https://doi.org/10.1016/S0198-9715(01)00031-X)
128
- 129 Thrasher, B., Wang, W., Michaelis, A., Melton, F., Lee, T., & Nemani, R. (2022). NASA
130 global daily downscaled projections, CMIP6. *Scientific Data*, 9(1), 262. <https://doi.org/10.1038/s41597-022-01393-4>
131