



Equations in Hyperbolic(-esque) Groups

Barak Ohana

The Hebrew University of Jerusalem

9 Sep 2024



Equations in Groups

Equations in
Hyperbolic(-
esque)
Groups

Barak Ohana

First section

Sample subsection

Second
section

- Let Γ be a group, and $\Sigma(x_1, \dots, x_d)$ a system of equation on d variables. We define the **set of solution of Σ in Γ** to be the set

$$V_{\Gamma}(\Sigma) = \left\{ (g_1, \dots, g_d) \in \Gamma^d \mid \Sigma(g_1, \dots, g_d) =_{\Gamma} 1 \right\}$$



Equations in Groups

Equations in Hyperbolic(- esque) Groups

Barak Ohana

First section

Sample subsection

Second section

- Let Γ be a group, and $\Sigma(x_1, \dots, x_d)$ a system of equation on d variables. We define the **set of solution of Σ in Γ** to be the set

$$V_{\Gamma}(\Sigma) = \left\{ (g_1, \dots, g_d) \in \Gamma^d \mid \Sigma(g_1, \dots, g_d) =_{\Gamma} 1 \right\}$$

- fsadas



Sample Frame Title No. 2

Equations in
Hyperbolic(-
esque)
Groups

Barak Ohana

First section

Sample subsection

Second
section

- First item
- Second item
- Third item



Sample Frame Title No. 3

Equations in
Hyperbolic(-
esque)
Groups

Barak Ohana

First section
Sample subsection

Second
section

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua.

Something important

Einstein's formula

$$E = mc^2$$