

The case $n = 4$

In this note, we will apply the same technique in $n = 2$ and $n = 3$ cases to the matrix $A_4(x_1, x_2, x_3, x_4, x_5)$, simply A_4 . By definition, we have

$$A_4 = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_1 & x_2 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & x_1 & x_2 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_1 & x_2 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_1 & x_2 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_1 & 0 & x_3 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & x_1 & 0 & x_3 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_1 & 0 & x_3 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_1 & 0 & x_3 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_1 & 0 & 0 & x_4 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & x_1 & 0 & 0 & x_4 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_1 & 0 & 0 & x_4 & 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_1 & 0 & 0 & x_4 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_1 & 0 & 0 & 0 & x_5 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & x_1 & 0 & 0 & 0 & x_5 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_1 & 0 & 0 & 0 & x_5 & 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_1 & 0 & 0 & 0 & x_5 \end{bmatrix}$$

As in cases $n = 2$ and $n = 3$, we may consider $\det A_4$ as a function of the variables x_1, x_2, x_3, x_4 and x_5 , i.e. $f(x_1, x_2, x_3, x_4, x_5) := \det A_4$. Without loss of generality we may assume that x_2, x_3, x_4 and x_5 are constant distinct scalars then the determinant function can be expressed as a univariate polynomial depending on the variable x_1 . Simply we may denote it $g(x_1) = \det A_4$. Since x_1 occurs only four different columns, $\deg g(x_1) = 4$. When $x_1 = x_2$, first five rows are linearly dependent, thus $g(x_2) = 0$. Similarly when $x_1 = x_3, x_1 = x_4$ and $x_1 = x_5$ the second, third and last five rows, respectively, are linearly dependent, thus $g(x_3) = g(x_4) = g(x_5) = 0$, as well. Since $\deg g(x_1) = 4$, we have

$$g(x_1) = k(x_1 - x_2)(x_1 - x_3)(x_1 - x_4)(x_1 - x_5),$$

where $k \in \mathbb{F}_q$ is the coefficient of the monomial x_1^4 . To determine k , we follow the same steps given in the cases $n = 2$ and $n = 3$.

Step 1: Cancel the x_1 's except the first five rows. To do this, we subtract the second row from the seventh, twelfth and seventeenth rows, the third row from the eighth, thirteenth and eighteenth rows, the fourth row from the ninth, fourteenth and nineteenth rows, finally the fifth row from the tenth, fifteenth and twentieth

rows. All these operations don't change the determinant's value. Thus we get:

[illegible]

Step 2: Move all columns including x_1 as entry, to the first four columns. To do this, move the sixth column to the second column (four moves), the eleventh column to the third column (eight moves) and the sixteenth column to the fourth column (twelve moves) without changing the order of others. Note that we have twenty four operations. Then

[illegible]

Step 3: Collapse the rows including x_1 as entry, into the first four rows. So, move the first row into the

[illegible][illegible]

3

Thus

$$g(x_1) = (-1)^{43}$$

$$\begin{aligned}
& \times \begin{vmatrix} x_1 & 1 & 0 & 0 & x_2 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & x_1 & 1 & 0 & 0 & 0 & 0 & 0 & -x_2 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & x_1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -x_2 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & x_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -x_2 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x_2 & x_3 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_2 & x_3 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_2 & x_3 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_2 & x_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x_2 & 0 & x_4 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_2 & 0 & x_4 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_2 & 0 & x_4 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_2 & 0 & x_4 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x_2 & 0 & 0 & x_5 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_2 & 0 & 0 & x_5 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_2 & 0 & 0 & x_5 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x_2 & 0 & 0 & x_5 \end{vmatrix} \\
& = - \begin{vmatrix} x_1 & 1 & 0 & 0 & * & * \\ 0 & x_1 & 1 & 0 & * & * \\ 0 & 0 & x_1 & 1 & * & * \\ 0 & 0 & 0 & x_1 & * & * \\ * & * & * & * & I_4 & 0 \\ * & * & * & * & * & A_3(x_2, x_3, x_4, x_5) \end{vmatrix},
\end{aligned}$$

where I_4 and 0 are 4×4 identity and 4×12 zero matrices, respectively and $*$ are some convenient matrices. By considering the determinant of block matrices in the above equation, the coefficient of the x_1^4 in $g(x_1)$ is the negative of the determinant of the matrix $A_3(x_2, x_3, x_4, x_5)$. Then from the case $n = 3$ we get

$$[x_1^4]g(x_1) = k = (x_4 - x_5)(x_3 - x_4)(x_3 - x_5)(x_2 - x_3)(x_2 - x_4)(x_2 - x_5).$$

Consequently, we get

$$\det A_4 = f(x_1, x_2, x_3, x_4, x_5) = g(x_1) = \prod_{1 \leq i < j \leq 5} (x_i - x_j)$$

as desired.