

Manuscript

Kyra Baranick

Bayesian hierarchical regression analysis of timing and magnitude for free-flowing river extremes in the Intermountain West

Kyra R. Baranick^{1,2}, Robert O. Hall, Jr.¹, and Nanette M. Nelson¹

¹ University of Montana: Flathead Lake Biological Station Polson, MT, USA

² University of California Santa Barbara, Santa Barbara, CA, USA

Abstract

Introduction

In high elevation temperate regions, greater than 60% of peak discharge events are driven by warmer storms arriving over vulnerable snowpack, known as rain-on-snow floods (Surfleet and Tullos 2013). In temperate mountain ranges, snowpack is often close to its melting point and sensitive to minor changes in temperature, precipitation type, or timing. Rain-on-snow events are responsible for some of the most devastatingly large floods in the Intermountain West, most notably the 1964, 500-year flood in Glacier County, Flathead County and the Blackfeet Reservation, that claimed 31 lives and caused \$539,000,000 million in damages (reported in 2026 CPI inflation adjusted value) (Survey 1967). In the last 100 years, the Intermountain West has warmed by 0.5C-1.5C and this number is expected to increase to 2C and 5C by the end of the century (Chambers, Devoe, and Evenden 2008). Under a warmer future, rain-on-snow frequency and spring precipitation are projected to increase, while snowpack is expected to decrease in quantity and melt earlier (Surfleet and Tullos 2013; Pederson et al. 2011)

Probabilistic analyses of United States Geologic Survey (USGS) stream gage data can provide robust estimates and quantification of uncertainty surrounding river timing and magnitude, capturing how climatic drivers combine into the realized hydrograph. While hydrologic research often projects changes in snowpack and temperature, decision making relies on in-situ USGS stream gage data, where these factors and others combine into the realized stream

discharge. Changes in timing and magnitude of hydrologic pulses may have impacts on human safety, agricultural production, and ecosystems. As the Intermountain West is a region of economic and population growth, water and emergency managers will need probabilistic estimates of hydrologic changes to make informed decisions that minimize economic, societal, and environmental costs.

This study employed a Bayesian hierarchical model to answer the following (1) To what extent are peak discharge, timing, and magnitude shifting in response to evolving hydrologic regimes using probabilistic estimates?; (2) How do these probabilistic estimates of change compare to a non-hierarchical, river-by-river regression model? To answer these questions we undertook the following research objectives.

Hydrologic change objectives:

1. Quantified shifts in the peak discharge timing across rivers by modeling changes in the day of the water year (DWY) that the annual peakdischarge occurs.
2. Estimated changes to the timing of the water year center of mass (COM) to determine the degree to which 50% runoff timing may be changing through time.
3. Estimated the change in the magnitude of annual peak discharge, to assess whether extreme discharge events are intensifying, diminishing or stabilizing through time.

Model evaluation objective:

4. Assessed the extent to which the Bayesian Hierarchical model produces more precise and interpretable trend estimates than the river-by-river regression.

This study makes several contributions to the understanding of flood hydrology. A common way to analyze trends is to analyze each river independently, what we refer to here as the non-hierarchical, river-by-river analysis. However, stream discharge data are highly sensitive to noise, meaning that treating each river as its own independent data set may reflect random fluctuation rather than true change over time. The Bayesian hierarchical model partially pools this information and assumes rivers share the same distribution, as they come from similar climatic regions. This dramatically decreases the influence of noise from the observation level and allows us to evaluate the underlying pattern of hydrologic change over time.

A comprehensive review of Bayesian methods in hydrology highlights this model as the cornerstone of modern hydrological modeling due to the capacity to handle uncertainties in data (Haddad 2025). Evaluating a river-by river regression with the Bayesian hierarchical regression reveals whether the individual trends are adequate indicators of regional change or whether hierarchical pooling is necessary to extract the climate driven signal.

Methods

Study area

This study focuses on the Intermountain West, specifically Montana, Idaho and northern Wyoming. The major mountain ranges involved include the Abasroka-Beartooth Mountains, Gallatin Range, Bitterroot Range, Clearwater Mountains, and the Lewis and Livingston Ranges with gauge elevation at about 400-2000m above sea level. The sampled area was limited to mountains east of the Cascades, as the climate in the Northern U.S. Rocky Mountain region is more continental, with a larger annual average temperature range of about 5C larger then the coastal western ranges. (Price 1978). It is estimated that over 50% and as high as 80% of the annual stream discharge originates as snow(Zheng et al. 2018; Pederson et al. 2011).

Many U.S rivers are a part of a dammed – or in other ways altered system – to manage water supply and protect cities and ecosystems from extreme high and low flows. However, the mountainous western portion of free-flowing river reaches can provide insights into the effects of climate change, with minimal direct human influence on stream discharge timing and quantity.




Figure 1: Sampled area including western Montana, northern Wyoming and Idaho. The 16 blue points represent the sampled gages.

Data

Daily and Instantaneous stream gage data was retrieved from USGS and sites were hand selected using USGS Stream Stats, an information and interactive map service (Survey 2026). Rivers and streams were selected in free-flowing watersheds that drain mountainous regions of the northwestern United States. Watersheds that contained dams, diversions, discharge-altering infrastructure or agricultural land greater then 1% of drainage area, were not eligible

Gage ID	Site Name	Drainage Area (km ²)	Gage Elevation (m)
USGS-05014500	SWIFTCURRENT CREEK AT MANY GLACIER MT	80.8	1,486.4
USGS-06043500	GALLATIN RIVER NEAR GALLATIN GATEWAY, MT	2,121.2	1,575.1
USGS-06078500	NORTH FORK SUN RIVER NEAR AUGUSTA MT	670.8	1,458.1
USGS-06090500	BELT CREEK NEAR MONARCH MT	919.4	1,213.1
USGS-06188000	LAMAR RIVER NR TOWER RANGER STATION YNP	1,730.1	1,828.8
USGS-06200000	BOULDER RIVER AT BIG TIMBER MT	1,359.7	1,236.4
USGS-06207500	CLARKS FORK YELLOWSTONE RIVER NR BELFRY MT	2,986.3	1,215.0
USGS-09188500	GREEN RIVER AT WARREN BRIDGE, NEAR DANIEL, WY	1,194.0	2,276.3
USGS-09196500	PINE CREEK ABOVE FREMONT LAKE, WY	197.6	2,270.8
USGS-12332000	MIDDLE FORK ROCK CR NR PHILIPSBURG MT	313.4	1,659.4
USGS-12354000	ST. REGIS RIVER NEAR ST. REGIS, MT	787.4	806.2
USGS-12355500	N F FLATHEAD RIVER NR COLUMBIA FALLS MT	4,030.0	958.8
USGS-12358500	M F FLATHEAD RIVER NEAR WEST GLACIER MT	2,913.7	953.6
USGS-12414500	ST JOE RIVER AT CALDER, ID	2,654.7	663.1
USGS-13336500	SELWAY RIVER NR LOWELL ID	4,959.8	467.4
USGS-13337000	LOCHSA RIVER NR LOWELL ID	3,051.0	445.8

for this study. On each identified river, the most upstream stream gage that began operation before 1950 and is still in operation as of 2026 was selected. All selected sites began operation between 1888 and 1954. Years that contained fewer than 300 days were excluded from the analysis.

A complete list of gages with corresponding drainage area, regulation status, land cover, and period-of-record metadata is provided in Appendix A.

A complete list of gages with corresponding drainage area, regulation status, land cover, and period-of-record metadata is provided in Appendix A.

Estimating hydrologic change

Data manipulation before modeling

1. Day of water year containing annual peak discharge—(referred to from here on as DWY) (objective 1): After daily value gage data was obtained from the USGS NWIS database, annual maxima were extracted for each water year. The water year is 365 or 366 days, beginning October 1st and ending September 31st of the following year. Each annual maxima is assigned a corresponding digit, 1-366, representing the day of the water year that the annual peak discharge occurs.
2. Center of mass of annual discharge—(referred to from here on as COM) (objective 2): The center of mass was calculated using the ‘faastr’ package in R. Mathematically, it is calculated by summing the daily discharge values (per year and per river) and identifying the day in which the cumulative total reaches 50% of that years discharge.

3. Annual maximum discharge (m^3/s) (objective 3): Daily value gauge data was obtained from the USGS NWIS database, annual maxima were extracted for each water year and converted to cubic meters per second (m^3/s). Standardized z-scores were calculated across individual river means. This procedure centers each years annual maximum discharge around the mean of the river it belongs to, scaled by its standard deviation, preventing trends from being dominated by river size.

$$A_{\text{std},r,t} = \frac{\log(A_{r,t}) - \mu_{\log A,r}}{\sigma_{\log A,r}}$$

Model

Model for center of mass and annual maximum peak discharge timing

To evaluate long-term effects of climate change on the timing of annual maximum stream discharge and the center of mass of discharge, we fit a multilevel linear regression model with partial pooling using the “brms” package in R. The response variable is the day of the year that the peak discharge or center of mass occurred and the ‘water year’ was held as the predictor to estimate change over time. This model was selected because the annual maximum stream discharge timing and the day of the center of mass is expected to vary while sharing the same climatic drivers. Partial pooling allows this trend to be estimated using information and strength coming from the individual observations of the river, in conjunction with the collective behavior of the rivers in this region, as a response to changes over time.

Weakly informative normal (0,0.5) priors were specified for each hierarchical regression and because our observations came from 16 different free-flowing rivers we allowed both the intercept and slopes to vary using:

$$(D_{y,r}) = \beta_{0,r} + \beta_{1,r} * t + \varepsilon_{y,r}$$

where $\beta_{0,r}$ is the river specific intercept, $\beta_{1,r}$ represents the river specific slope, t is the time in years and $\varepsilon_{y,r}$ is the normally distributed error in the observations.

The rivers are assumed to come from a multivariate normal distribution of:

$$\begin{pmatrix} \beta_{0,r} \\ \beta_{1,r} \end{pmatrix} \sim \text{MVN} \left(\begin{pmatrix} \mu_{\beta_0} \\ \mu_{\beta_1} \end{pmatrix}, \Sigma \right)$$

With this model partial pooling is implied in that each individual river borrows strength from the full data set.

Model for annual maximum discharge

The change in DWY was also evaluated using the Bayesian hierarchical model however, because rivers differ substantially in drainage area, many of the discharge values vary in magnitude. To place all rivers on a comparable scale, the regression model for the change

in annual maximum discharge over time was log-transformed and then standardized using a z-score transformation.

To interpret the results all slope values were back transformed to percent change per year. Because the regression model was fit using z-score, the original output describes changes in standardized log units per year. This standardized values $A_{\text{std},r}$ were first multiplied by each rivers standard deviation $\sigma_{\log A,r}$ of log discharge and then exponentiated to represent a multiplicative factor for annual change in discharge $\exp(\Delta A_{\text{std},r} \cdot \sigma_{\log A,r}) - 1$. This change was multiplied by 100 to obtain the annual % change for each river, $\beta_{1,r}$.

$$\% \Delta A_r = (\exp(\Delta A_{\text{std},r} \cdot \sigma_{\log A,r}) - 1) \times 100$$

Model evaluation method

To evaluate whether the Bayesian hierarchical model produces more robust slope estimates and uncertainty quantification, we fit a simple linear regression to each watershed for comparison. We then compared slope and uncertainties across individual watersheds to determine whether the Bayesian model detects a coherent signal the non-hierarchical fit fails to detect. This comparison allows us to determine whether partial pooling reduces overfitting in small basins, and provides decision relevant uncertainty estimates.

Results




Figure 2: Posterior slope probability density histogram for center of mass (COM), day of water year containing annual peak discharge (DWY), and discharge, based on 1000 posterior draws. Horizontal segments across the x-axis represent the 90% confidence interval for each model and the dot represents the posterior mean slope. The red vertical line is the zero threshold, meaning no positive or negative slope. This visualization highlights the difference in slope and confidence among the modeled variables with discharge exhibiting a tightly constrained slope near zero and the DWY and COM showing a stronger negative slope.

Timing: DWY, COM

Across the 16 rivers, posterior estimates from the Bayesian Hierarchical model indicate a regional shift towards earlier COM and DWY timing. The model estimates that the day of the water year that contains the peak annual discharge is shifting between 0.10 and 0.21 days earlier per year, meaning peak annual discharge will occur roughly a week earlier every 50 years. The model estimates that the COM is shifting by 0.03 to .07 days per year, equivalent to 3 to 7 days every 100 years. Figure 1 shows the full posterior distribution for the regional COM and DWY trend. The distribution for both timing metrics is centered entirely below zero with a narrow confidence interval demonstrating high certainty in the estimate.

Magnitude: Annual Peak Discharge (m^3/s)

The 95% confidence interval for the slope of the magnitude of the annual maximum discharge over time remains tightly around zero, indicating the the model detected no regional change to discharge over time (Figure 1).

Justification for Pooling

Using the DWY variable as an example, we contrast the river by river non-hierarchical regression model with the Bayesian regression model.

When examining data from the DWY variable , it is clear that the observation level data is highly variable (Figure 2). When fitting a linear regression line to these points, it is unclear whether the slope is a product of data noise or true change over time. The linear model can also lead to overfitting especially when the record of data contain large gaps. In several rivers such as the St. Regis River, and the North Fork Sun River the data contains two period of large data gaps and several outliers that steepen the river specific slope estimate. In other rivers, such as the Green River and the Selway river, the data appear in a cloud shape around the regression line, making the slope estimate quite uncertain and difficult to interpret.




Figure 3: Time Series for day of the water year that the peak annual discharge occurs (DWY). The individual observations of DWY are plotted fit with simple linear regression lines (blue). All slopes are negative with the exception of the Boulder river.

Treating these rivers as a interconnected system draws meaning from this noise and makes the slope estimate interperetable. The river-by-river non-hierarchical model contains variable slope estimations and large confidence intervals, over half of which cross zero, indicating the directional trend is indistinguishable. In contrast, the Bayesian hierarchical model produces a narrower distribution of slopes and smaller uncertainty across rivers. This pattern reflects shrinkage, where rivers with gaps or noisy observations are pulled towards the regional mean.




Figure 4: Comparison of river level slope estimates for the day of water year (DWY) containing the annual peak discharge from the non-hierarchical linear model (red) and the Bayesian hierarchical model (blue). Points represent the slope estimate for each river and model and 95% confidence intervals are shown.

Discussion

Our Bayesian hierarchical analysis reveals a consistent decreasing timing (DWY and COM) across snowmelt dominated, free-flowing rivers in the Intermountain West but shows no change in magnitude of floods (annual max discharge). We also demonstrate that these statements are estimated with greater certainty using this model over the noisier river-by-river linear regression.

Numerous observational studies have proven earlier peak timing in snowmelt dominated basins across the globe (Dudley et al. 2017; “Changes in the Timing of High River Flows in New England over the 20th Century” 2003; Chen et al. 2016). Our study shows that the Intermountain West is not exception to this trend. It is noted that this trend will continue

due to warmer temperatures under a business-as-usual climate scenario (Stewart, Cayan, and Dettinger 2004).

It has been widely accepted that a warmer climate will accelerate the hydrologic cycle, increase atmospheric water holding capacity, precipitation extremes, and the occurrence of rain-on-snow events (“Climate Change Impacts on the Hydrological Cycle” 2008; Fischer and Knutti 2016; Surfleet and Tullos 2013). However, there has been disagreements about whether these factors directly translate into increased peak discharge over time (Zhang et al. 2022). Consistent with the results of this paper, many snowmelt dominated regions show low-to-no trend of increasing magnitude or peak discharge consistent with time or increase in CO (Hirsch and Ryberg 2012; Kundzewicz et al. 2014). We hypothesize that this is due to a spacial variability and multitude of warming driver factors that may suppress or accelerate runoff peaks. Theories include, changes in basin specific soil moisture, changes in flood drivers, (Zhang et al. 2022; “Streamflow Response to Snow Regime Shift Associated with Climate Variability in Four Mountain Watersheds in the US Great Basin” 2019) or competition between decreasing snow pack and accelerated runoff (Wang, Kumar, and Link 2016). Because our study is in the rain-snow transition zone, and runoff drivers are shifting, its is reasonable that our Bayesian Hierarchical model did not estimate a positive or negative trend in peak discharge magnitude. However, because we did not group by runoff drivers (rain-on-snow, snow pack, or precipitation type) it is possible that more specific change in peak discharge magnitude is occurring despite lack of complete regional change.

The type of analysis performed also drives interpretation of climate driven change. In our comparison of the non-hierarchical linear regression and the Bayesian regression model, we show the influence of partial pooling on the success of extracting the regional signal.

(not sure about this next part...)

We also noticed that many hydrologists use the Manning-Kendall (MK) test [Dudley et al. (2017);]. This evaluates each river independently and assigns a +1, -1, or 0 for each pair of years representing a positive, negative or no change from the last year to the next year. MK treat rivers independently

It is important to recognize that the Bayesian hierarchical model contains limitations as well but due to _____ is is often able to detect a credible signal while other tests may fail. The greatest advantage of the Bayesian Hierarchical Model is the computation of a full posterior distribution and credible intervals for the certainty of the regional estimate. This model can also be adapted to more complex applications and the model can be updated as new climate and hydrologic data is published (Haddad 2025).




Figure 5: all using freqpoly




Figure 6: Time Series for Day of water year for the Center of Mass of Peak Annual Discharge.

Comparison of LM vs Bayesian Hierarchical Slopes for Center of Mass




Figure 7: Comparison of LM vs Bayesian Hierarchical Slopes for COM.

Timing: Day of Year of Peak Annual Discharge

Volume: Annual Peak Discharge in Percent Change per Year




Figure 8: Time Series for DWY.




Figure 9: Comparison of LM vs Bayesian Hierarchical Slopes for DWY.

Supplementary Materials

Supplementary Materials

Supplementary Materials

Kyra Baranick

Supplementary Material A: Site Justification

Selection Method

- Rivers must be free-flowing. No upstream dams diversions or flow altering infrastructure.
- Significant agricultural withdraw upstream disqualifies the gage.
- Gage record must contain 70+ years worth of data
- Gage must continue to record data as of 2026
- No partial record gage (partial record gauge: only measures peak or low flows season)
- Most upstream age that satisfies the above measures is selected.

Site List and Justification

1. “USGS-06043500”, Gallatin River (At Gallatin Gateway, MT)

- Regulation: USGS Stream Stats regulation check returns a 1.33% regulation in the watershed upstream of the gage (Survey 2026).
- NWIS Discharge period of record : 08/01/1989 - 06/23/2026 (Present)
- Drainage Area: 2121km² (819 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the 5th gauge from the head waters. The previous 4 gages are decommissioned, recently implemented or have incomplete/partial periods of record. This basin drains part of Yellowstone and Gallatin National Forest.

2. “USGS-12332000”, Middle Fork Rock Creek (Near Philipsburg, MT)

- Regulation: USGS Stream Stats regulation check returns a 0.00% regulation in the watershed upstream of the gage (Survey 2026).
- NWIS Discharge period of record : 10/01/137 - 06/23/2026 (Present)
- Drainage area: 313 km² (121 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%

- This is the most upstream gage in the watershed. It does not contain any farmland or developed land. The drainage area drains Deerlodge National Forest.

3. **“USGS-09196500”, Pine Creek (Above Freemont Lake, WY)**

- Regulation: No identified regulation or farm areas (Google 2026)
- NWIS Discharge period of record : 10/01/1937 - 06/23/2026 (Present)
- Drainage area: 196km² (76 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the most upstream gage in the watershed. The watershed contains many natural lakes and mountainous terrain. The basin drains Bridger Wilderness and Bridger National Forest and is upstream of Pine Lake.

4. **“USGS-06090500”, Belt Creek (Near Monarch, MT)**

- Regulation: USGS Stream Stats regulation check returns a 0.00% regulation in the watershed upstream of the gage (Survey 2026).
- NWIS Discharge period of record: 15/01/1951 - 06/23/2026 (Present)
- Drainage area: 917km² (354 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the most upstream gage in the watershed. This basin drains Cascade County and Lewis and Clark National Forest.

5. **“USGS-12358500”, Middle Fork Flathead River(Near West Glacier, MT)**

- Regulation: USGS Stream Stats regulation check returns a 0.00% regulation in the watershed upstream of the gage (Survey 2026).
- NWIS Discharge period of record : 10/01/1939 - 06/23/2026 (Present) -
- Drainage area: 2194 km² (1125 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This gage is 10 gages from the most upstream gage, however, all previous 9 gages are decommissioned, recently implemented or have incomplete or partial periods of record. This is downstream of lake McDonald and downstream of the confluence between McDonald Creek and Middle Fork Flathead River. The drainage area includes part of Glacier National Forest, Flathead National Forest, Great Bear Wilderness and as far as the Sawtooth Mountain Range.

6. **“USGS-06188000”, Lamar River (Near Tower Ranger Station, WY)**

- Regulation: No identified regulation or farm areas (Google 2026).
- NWIS Discharge period of record : 05/01/1923- 06/23/2026 (Present)
- Drainage area: 1730 km² (668 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the most upstream age of the Lamar River. The drainage area is within Yellowstone National Park.

7. **“USGS-09188500”, Green River (At Warren Bridge, near Daniel, WY)**

- Regulation: No identified regulation or farm areas (Google 2026).
- NWIS Discharge period of record : 10/01/1931- 06/23/2026 (Present)
- Drainage area: 1193 km² (461 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the most upstream gage of the Green River. This drains part of the Bridger-Teton National Forest.

8. **“USGS-13336500”, Selway River (Near Lowell, ID)**

- Regulation: No identified regulation or farm areas (Google 2026).
- NWIS Discharge period of record : 04/11/1911- 06/23/2026 (Present)
- Drainage area: 4962 km² (1915 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the 5th gage from the head waters. The previous 4 gages are decommissioned, recently implemented or have incomplete or partial periods of record. This drainage area drains Selway-Bitterroot National Forest, Bitterroot Wilderness and Nezperce National Forest.

9. **“USGS-12355500”, North Fork Flathead River (Flathead British Colombia, MT)**

- Regulation: USGS Stream Stats regulation check returns a 0.00% regulation in the watershed upstream of the gage (Survey 2026).
- NWIS Discharge period of record : 04/01/1929- 06/23/2026 (Present)
- Drainage area: 4030 km² (1556 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the most upstream gage form the headwaters. This gage sits of the border of the U.S. and Canada and drains parts of Glacier National Park and Flathead National Forest.

10. **“USGS-12414500”, St. Joe River (at Calder, ID)**

- Regulation: One farm in Herrick, ID (Survey 2026).
- NWIS Discharge period of record : 05/01/1911 - 06/23/2026 (Present)
- Drainage area: 2654 km² (1025 miles²)
- Agricultural Land in Percentage of Drainage Area: .011%
- This is the 4th gage from the head waters. The previous 3 gages are decommissioned, recently implemented or have incomplete or partial periods of record. This drainage area drains the Saint Joe National Forest near the Bitterroot range.

11. **“USGS-06207500”, Clarks Fork Yellowstone River (Near Belfry, MT)**

- Regulation: USGS Stream Stats regulation check returns a 0.00% regulation in the watershed upstream of the gage (Survey 2026).

- NWIS Discharge period of record : 09/01/1921 - 06/23/2026 (Present)
- Drainage area: 2983 km² (1152 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the 6th gage from the head waters. The previous 5 gages are decommissioned, recently implemented or have incomplete or partial periods of record. This drainage area drains parts of Yellowstone National Park

12. **“USGS-13337000”, Locsha River (Near Lowell, ID)**

- Regulation: No identified regulation or farm areas (Survey 2026).
- NWIS Discharge period of record : 10/01/1910 - 06/23/2026 (Present)
- Drainage area: 3048 km² (1177 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the 5th gage from the head waters. The previous 4 gages are decommissioned, recently implemented or have incomplete or partial periods of record. This drainage area drains parts of Clearwater National Forest.

13. **“USGS-06078500”,North Fork Sun River (Near Augusta, MT)**

- Regulation: USGS Stream Stats regulation check returns a 0.00% regulation in the watershed upstream of the gage (Survey 2026).
- NWIS Discharge period of record : 05/01/1911- 06/23/2026 (Present)
- Drainage area: 670 km² (259 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the most upstream gage from the headwaters. This drains part of Lewis and Clark National Forest.

14. **“USGS-06079000”, South Fork Sun River (Near Augusta, MT)**

- Regulation: USGS Stream Stats regulation check returns a 0.00% regulation in the watershed upstream of the gage (Survey 2026).
- NWIS Discharge period of record : 05/01/1911- 06/23/2026 (Present)
- Drainage area: 650 km² (251 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the most upstream gage from the headwaters. This drains part of Lewis and Clark National Forest.

15. **“USGS-16102500”, Teton River (Below South Fork near Choteau, MT)**

- Regulation: USGS Stream Stats regulation check returns a 0.00% regulation in the watershed upstream of the gage (Survey 2026).
- NWIS Discharge period of record : 06/01/1947- 06/23/2026 (Present)
- Drainage area: 285 km² (110 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the most upstream gage from the headwaters. This drains part of Teton County.

16. **“USGS-05014500”, Swiftwater Creek (At Many Glacier, MT)**

- Regulation: USGS Stream Stats regulation check returns a 0.00% regulation in the watershed upstream of the gage (Survey 2026).
- NWIS Discharge period of record : 06/01/1912- 06/23/2026 (Present)
- Drainage area: 80 km² (31 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the 5th gage from the head waters. The previous 4 gages are decommissioned, recently implemented or have incomplete or partial periods of record. This drainage area drains parts of Glacier national Park. The gage sits downstream of Swiftcurrent Lake.

17. **“USGS-12354000”, St. Regis River (Near St. Regis, MT)**

- Regulation: USGS Stream Stats regulation check returns a 0.00% regulation in the watershed upstream of the gage (Survey 2026).
- NWIS Discharge period of record : 09/01/1910- 06/23/2026 (Present)
- Drainage area: 787 km² (304 miles²)
- Agricultural Land in Percentage of Drainage Area: 0%
- This is the most upstream gage from the headwaters. This basin drains part of Lolo National Forest.

18. **“USGS-06200000” Boulder River (at Big Creek, MT)**

- Regulation: USGS Stream Stats regulation check returns a 0.00% regulation in the watershed upstream of the gage (Survey 2026).
- NWIS Discharge period of record : 04/01/1947- 06/23/2026 (Present)
- Drainage area: 1357 km² (524 miles²)
- Agricultural Land in Percentage of Drainage Area: 1%
- This is the 5th gage from the head waters. The previous 4 gages are decommissioned, recently implemented or have incomplete or partial periods of record. This basin drains part of Gallatin National Forest and the Absaroka Mountains.

Google. 2026. “Google Earth Pro.” <https://earth.google.com/>.

Survey, U. S. Geological. 2026. “StreamStats: A u.s. Geological Survey Web Application.” <https://streamstats.usgs.gov/>.

- Chambers, Jeanne C., Nora Devoe, and Angela Evenden. 2008. "Collaborative Management and Research in the Great Basin: Examining the Issues and Developing a Framework for Action." RMRS-GTR-204. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. <https://doi.org/10.2737/RMRS-GTR-204>.
- "Changes in the Timing of High River Flows in New England over the 20th Century." 2003. *Journal of Hydrology* 278 (1): 244–52. [https://doi.org/https://doi.org/10.1016/S0022-1694\(03\)00155-0](https://doi.org/https://doi.org/10.1016/S0022-1694(03)00155-0).
- Chen, Xing, Mukesh Kumar, Rui Wang, Adam Winstral, and Danny Marks. 2016. "Assessment of the Timing of Daily Peak Streamflow During the Melt Season in a Snow-Dominated Watershed." *Journal of Hydrometeorology* 17 (8): 2225–44. <https://doi.org/10.1175/JHM-D-15-0152.1>.
- "Climate Change Impacts on the Hydrological Cycle." 2008. *Ecohydrology & Hydrobiology* 8 (2): 195–203. <https://doi.org/https://doi.org/10.2478/v10104-009-0015-y>.
- Dudley, Robert W, Glenn A Hodgkins, MR McHale, Michael J Kolian, and Benjamin Renard. 2017. "Trends in Snowmelt-Related Streamflow Timing in the Conterminous United States." *Journal of Hydrology* 547: 208–21.
- Fischer, Erich M, and Reto Knutti. 2016. "Observed Heavy Precipitation Increase Confirms Theory and Early Models." *Nature Climate Change* 6 (11): 986–91.
- Haddad, Khaled. 2025. "A Comprehensive Review and Application of Bayesian Methods in Hydrological Modelling: Past, Present, and Future Directions." *Water* 17 (7): 1095.
- Hirsch, R. M., and K. R. Ryberg. 2012. "Has the Magnitude of Floods Across the USA Changed with Global CO2 Levels?" *Hydrological Sciences Journal* 57 (1): 1–9. <https://doi.org/10.1080/02626667.2011.621895>.
- Kundzewicz, Zbigniew W., Shinjiro Kanae, Sonia I. Seneviratne, John Handmer, Neville Nicholls, Pascal Peduzzi, Reinhard Mechler, et al. 2014. "Flood Risk and Climate Change: Global and Regional Perspectives." *Hydrological Sciences Journal* 59 (1): 1–28. <https://doi.org/10.1080/02626667.2013.857411>.
- Pederson, Gregory T, Stephen T Gray, Toby Ault, Wendy Marsh, Daniel B Fagre, Andrew G Bunn, Connie A Woodhouse, and Lisa J Graumlich. 2011. "Climatic Controls on the Snowmelt Hydrology of the Northern Rocky Mountains." *Journal of Climate* 24 (6): 1666–87.
- Price, Larry W. 1978. "Mountains of the Pacific Northwest, USA: A Study in Contrasts*." *Arctic and Alpine Research* 10 (2): 465–78.
- Stewart, Iris T, Daniel R Cayan, and Michael D Dettinger. 2004. "Changes in Snowmelt Runoff Timing in Western North America Under Abusiness as Usual'climate Change Scenario." *Climatic Change* 62 (1): 217–32.
- "Streamflow Response to Snow Regime Shift Associated with Climate Variability in Four Mountain Watersheds in the US Great Basin." 2019. *Journal of Hydrology* 573: 255–66. <https://doi.org/https://doi.org/10.1016/j.jhydrol.2019.03.021>.
- Surfleet, Christopher G., and Desirée Tullos. 2013. "Variability in Effect of Climate Change on Rain-on-Snow Peak Flow Events in a Temperate Climate." *Journal of Hydrology* 479: 24–34. <https://doi.org/https://doi.org/10.1016/j.jhydrol.2012.11.021>.
- Survey, U. S. Geological. 1967. "Floods of December 1964 in the Far Western States." 1840-C. Water-Supply Paper. Washington, D.C.: U.S. Geological Survey. <https://pubs.usgs.gov/publication/wsp1840C>.

- . 2026. “StreamStats: A u.s. Geological Survey Web Application.” <https://streamstats.usgs.gov/>.
- Wang, Rui, Mukesh Kumar, and Timothy E Link. 2016. “Potential Trends in Snowmelt-Generated Peak Streamflows in a Warming Climate.” *Geophysical Research Letters* 43 (10): 5052–59.
- Zhang, Shulei, Liming Zhou, Lu Zhang, Yuting Yang, Zhongwang Wei, Sha Zhou, Dawen Yang, et al. 2022. “Reconciling Disagreement on Global River Flood Changes in a Warming Climate.” *Nature Climate Change* 12 (12): 1160–67.
- Zheng, Xiaohui, Qiguang Wang, Lihua Zhou, Qing Sun, and Qi Li. 2018. “Predictive Contributions of Snowmelt and Rainfall to Streamflow Variations in the Western United States.” *Advances in Meteorology* 2018 (1): 3765098.