

Heat, Health, and Harm: The Impact of Extreme Temperatures on Suicide

Yegor Baranovski *

February 20, 2026

Abstract

Climate change poses well-established risks to physical health, but little is known about its effects on mental health. I study the relationship between extreme temperature exposure during summer months and suicide risk in the United States from 1989 to 2019 using comprehensive county-level data on all deaths by suicide and high-resolution satellite-based temperature measurements. I identify the causal effects of extreme heat on suicide by relating year-over-year fluctuations in county-level summer monthly average temperatures to fluctuations in suicide rates. I compare these effects across urban and rural areas and various population subgroups. My results indicate that an increase of one defined extreme heat day in a summer month leads to approximately 0.091 additional suicide deaths per million residents for summer months, with effects concentrated among men and working-age adults, demographic groups already characterized by elevated baseline suicide risk. This paper provides large-scale empirical evidence that climate-related extreme heat significantly elevates suicide risk, underscoring the urgent need for targeted public health interventions to mitigate climate-induced mental health burdens.

*Baranovski: Honors Economics Undergraduate Thesis, UW-Madison, yegorbar01@gmail.com. Special thanks to my advisor Professor Ashley Swanson for overseeing my work and putting up with me, along with the countless people I complained and vented to.

Contents

1	Introduction	1
1.1	Motivation	1
1.2	Contribution	2
2	Data and Methodology	4
2.1	Population Data	4
2.2	Mortality Data	5
2.3	Temperature Data	5
2.4	Urbanization Classification	5
2.5	Empirical Approach	6
3	Results	7
4	Discussion	8
4.1	Limitations	9
5	Future Directions	10
6	Replication	10
7	Appendix	11
	References	16

1 Introduction

1.1 Motivation

Mental illness represents an enormous and expanding global health challenge, both economically and socially. In the United States, approximately 49,450 individuals die by suicide each year, contributing to the nearly 727,000 suicide deaths recorded worldwide annually [25], [27]. The prevalence of mental health issues is also high, with 23.1% of U.S. adults reporting a diagnosable mental, emotional, or behavioral disorder as of 2022 [24]. Financially, mental disorders impose substantial direct and indirect costs. Total direct societal burden was estimated to be \$280 billion and is expected to rise, making mental illness the costliest medical category in the country [1]. When considering indirect costs, including reduced productivity, lost wages, disability, and mortality, the total societal impact approaches \$450 billion annually in the U.S. and around \$2.5 trillion globally, figures projected to double by 2030 [36]. Beyond these economic impacts, mental health disorders also exacerbate physical health conditions, significantly increasing the overall health burden [31].

Climate conditions are increasingly recognized as influential on various aspects of human health, impacting outcomes ranging from vector-borne disease prevalence to rates of cardiovascular incidents [29], [38]. These health outcomes can be driven by direct physical stress (such as heatstroke or cyclone-related injuries), shifts in disease ecology (such as influenza or malaria dynamics), or socioeconomic disruptions such as drought-related famine [38]. Additionally, studies indicate that higher temperatures may increase interpersonal conflicts and violence, potentially through economic stress or physiological responses that heighten aggression [17].

Despite extensive research into physical health impacts, the connection between climate factors and mental health, specifically suicide rates, has received limited quantitative analysis [6], [9]. Understanding this link is critical, given that suicide deaths globally exceed those caused by interpersonal and group violence combined [19], rank among the top causes of death worldwide, and represent a significant portion of lost life-years, particularly in wealthier nations [20]. In the United States specifically, suicide is one of the leading causes of death among individuals aged 10–54 [15] and uniquely remains stable or increasing, unlike other major causes of death which are generally declining [35]. Consequently, even modest climate-related increases in suicide rates could significantly elevate global mental health burdens, particularly in developed


regions already experiencing high or rising suicide incidence.

1.2 Contribution

This paper contributes a unique approach in quantifying the relationship between suicide risk and climate change. Specifically, I analyze the effects of extreme and abnormal heat on mental health–related mortality, using a granular and wide–ranging dataset to compare impacts between urban and rural areas across multiple demographic subgroups. My approach offers four distinct advantages.


First, to more comprehensively capture mental health–related mortality, I define mental health–related deaths to include not only suicides but also deaths classified as injuries of undetermined intent and certain types of accidental deaths (such as poisoning, firearm, train, and drowning incidents), defined further in Section 2.2. This broader definition addresses the known problem of misclassification in mortality data [4], where some suicides may be recorded as accidents due to reporting practices or legal ambiguity. By including categories plausibly linked to underlying mental health crises, this approach provides a more complete measure of the population–level impact of extreme heat on mental health, improving power. I use suicide risk and mental health mortality in this paper interchangeably.

Figure 1: Average Monthly Suicide Rate(Deaths per Million) by County




Second, I define extreme heat exposure using both relative and absolute threshold criteria, rather than relying solely on simple temperature averages. Specifically, I calculate extreme heat days (EHDs) as the number of days in a month where the daily mean temperature exceeds either the 85th percentile of the 30-year historical monthly distribution for a given county or a daily mean temperature threshold of 30°C. This dual-definition approach contextualizes "extreme" relative to local climatological norms and human acclimation [30] while also accounting for objective human physiological tolerance levels, enabling more meaningful comparisons across regions and over time, and better capturing the types of heat events most relevant to health outcomes.

Figure 2: Average Summer Percentile Thresholds(°C) by County



Third, the estimated effects presented in this paper represent a lower bound on the true impact of temperature on mental health-related mortality. By defining extreme heat days using relatively high thresholds, either exceeding the 85th percentile of historical daily temperatures or surpassing 30°C, this approach focuses on only the most severe temperature exposures. However, milder but still abnormal heat events, as well as cumulative heat stress from sustained periods of moderately high temperatures, may also contribute to adverse mental health outcomes [26]. As such, the full burden of heat on mental health could be even larger than what is captured by the estimates reported here.

Figure 3: Trends in suicide rates and extreme heat days across urban and rural areas.



Fourth, the use of a granular, county-level panel dataset allows me to exploit both spatial and temporal variation in extreme heat exposure. By applying fixed effects and controlling for time trends at multiple levels, I am able to isolate the short-term causal effect of an extreme heat day on mental health mortality, while also exploring heterogeneity across urban and rural settings and demographic subgroups.

2 Data and Methodology

2.1 Population Data

County-level population data was used to calculate death rates and for regression weighting purposes. Subgroup county population data by sex, age, and race are obtained from the Surveillance, Epidemiology, and End Results Program [34]. I supplement this with county-level educational attainment data from the Economic Research Service branch of the US department of Agriculture that reports the number of adults 25+ in each educational attainment group (high school or less, some college or more) for 1980, 1990, 2000, and a 5 year average for 2016–2020 [37]. For each county, I linearly interpolate from 1980 to 2020 to construct annual estimates of county-level education attainment for 1989–2019. I then calculate the death rate as standard.

2.2 Mortality Data

Data on monthly health related deaths at the county level are derived from the restricted use Detailed Mortality–All County data; part of the Vital Statistics Data maintained by the NCHS and can become available via request and application from the Centers for Disease Control [12]. The raw data acquired includes every death reported in the United States between 1989 and 2019, along with the primary cause of each. Cause of death was used to categorize deaths as mental health related if they were caused by suicide, injuries of undetermined intent, or accidental poisoning, firearm, train, or drowning, explicitly detailed in table 1 [4]. For a given county in a given summer month each year, I counted the number of mental health related deaths, both overall and by subgroup. The set of subgroups includes sex (M,F), age (0–24, 25–64, 65+), race/origin (White and non-Hispanic, non-White or Hispanic), and educational attainment (high school diploma or less, some college credit or more). Analysis for educational attainment was limited to 25+ adults, as younger people may not have had time to get a college education. Group specific deaths rates are calculated in terms of number of deaths per million using group – specific population counts. All figures suppressed counties that observed fewer than 10 total or subgroup mental health related deaths for the study period but are still included in the analysis.

2.3 Temperature Data

Temperature data was obtained from the PRISM Climate Group [32], which reports high resolution gridded climate data. The PRISM data contains 4km by 4km gridded estimates of daily average temperature for the United States for from 1989–2019. I aggregated the daily average temperature data to the county level and then mapped the set of daily average county temperatures in a given month and year to the number of extreme heat days (described below in 2.5).

2.4 Urbanization Classification

Counties are classified as urban or rural according to the 2013 NCHS Urban–Rural Classification Scheme [23]. This system assigns each county to one of six categories based on its level of urbanization. For my analysis, I define urban counties as those falling into the top

three categories: large central metro, large fringe metro, and medium metro. Counties in the bottom three categories – small metro, micropolitan, and noncore – are considered rural.

2.5 Empirical Approach

I use a panel fixed effects regression to estimate the effect of smoke exposure on suicide rates:

$$Y_{cym} = \beta \cdot \text{EHD}_{cym} + \delta_{cm} + \delta_{cy} + \delta_{ym} + \epsilon_{cym},$$


$$\text{where } \text{EHD}_{cym} = \sum_{d=1}^{|m|_y} \mathbb{1}(T_{cymd} > P_{85,cm} \vee T_{cymd} > 30^\circ\text{C})$$

Y_{cym} is the defined suicide rate with deaths by suicide per million in county c , year y , and month m . EHD_{cym} denotes the number of days in the same county–year–month where the average daily temperature exceeded either the 85th percentile threshold for that county–month or 30°C . Percentile thresholds are calculated separately for each county and month, using historical daily temperature data from 1989 to 2019. δ_{cm} represents county–by–month fixed effects, which subtract the county–specific monthly average from both suicide rates and EHDs. This ensures that identification comes from within–county deviations, comparing county–months in relatively low–EHD years to the same county–months in relatively high–EHD years. My main specification also includes county–year fixed effects, δ_{cy} , which control for all county–specific factors that vary annually but not within a year (e.g., changes in economic conditions, demographic shifts). This prevents the need to separately control for county–level time–varying covariates. I also include year–month fixed effects, δ_{ym} which account for shocks common to all counties in a given month and year (such as the impacts of the 2008 financial crisis or coding changes like the ICD–10 transition). All regressions are weighted by 2000 Census county populations to account for differences in county size, and standard errors are clustered at the county level to allow for arbitrary serial correlation in the error terms within counties.

3 Results

I begin by presenting the estimated relationship between EHDs and suicide rates based on the baseline fixed effects regression model described in section 2.5.

Figure 4: Effect of one additional EHD on monthly suicide rates, by subgroup.



The markers indicate point estimates while the lines the 95% CI estimates of the indicated relationships. Estimates for each population subgroup and area type are taken from separate regressions based on population subgroup-specific suicide rates in the indicated sample of counties. “White/NH” refers to those who are White and non-Hispanic, and “NW/Hispanic” is the converse (non-White or Hispanic). “HS or less” refers to adults 25 years and older with a high school diploma or less, and “College+” refers to adults 25 years and older with some college credits or more.

For robustness, results were validated using Poisson regression models, where coefficients capture proportional changes rather than absolute changes in suicide rates. The Poisson models produced similar directional findings across subgroups, although the magnitudes of the effects differed somewhat between urban and rural counties. These differences likely reflect

the influence of lower baseline suicide rates in certain areas, which can affect proportional versus absolute change estimates. Overall, the consistency in the direction of effects across specifications reinforces the robustness of the main findings. Results are more unstable among rural areas due to count sizes. Regression estimates and mean suicide rates for each subgroup are reported in the Appendix, tables 2 and 3.

4 Discussion

My results demonstrate that exposure to EHDs increases suicide mortality in the United States. I estimate that one additional EHD in a summer month leads to an increase of 0.091 suicide deaths per million residents. This corresponds to a 0.39% rise relative to the baseline monthly suicide rate of 23.66 per million. While modest in absolute terms, this effect may accumulate meaningfully as extreme heat events become more frequent and intense under climate change, and with our definition of an EHD, this may serve as a possible lower bound. The economic implications are also significant, given the high direct and indirect costs associated with suicide and mental health-related deaths.

The effect is slightly more pronounced in rural populations and among demographic groups with elevated baseline suicide risk, including men and working-age adults. These findings suggest that extreme heat may exacerbate existing vulnerabilities, potentially widening existing mental health disparities.

Several mechanisms could explain the observed relationship [6]. At a physiological level, heat stress activates the hypothalamic-pituitary-adrenal axis, elevating cortisol and disrupting serotonergic pathways that regulate mood and impulsivity [33]. Elevated temperatures also impair sleep quality, and sleep disturbance is a well-documented risk factor for suicidal ideation and behavior [28]. More broadly, higher temperatures have been shown to increase aggression, substance use, and interpersonal conflict, each of which is independently associated with suicide risk [17]. These pathways may be particularly salient for individuals in lower socioeconomic groups, who are more likely to engage in outdoor labor or lack access to cooling resources, and for whom heat exposure may compound existing economic and psychological stressors [9].

The observed association between extreme heat and suicide highlights the often-overlooked

mental health dimension of climate change. If current trends in temperature extremes continue, the mental health burden could grow considerably. These findings underscore the need for targeted public health interventions—such as early warning systems, mental health outreach, and access to cooling infrastructure—especially in high-risk communities. Interestingly, although rural communities face higher baseline suicide risk, urban areas tend to experience more severe heat exposure. This contrast may help explain the similarity in urban and rural point estimates, as greater heat intensity in urban areas could offset lower baseline vulnerability.

4.1 Limitations

Several limitations of this study should be acknowledged. First, for the definition of EHD, the analysis was limited to summer months (June, July, and August). While this approach captures periods with the highest likelihood of extreme heat, it excludes potential effects of unseasonal heat spikes in spring or fall, potentially underestimating the overall impact of temperature on mental health, particularly as suicides typically peak in spring [7].

Additionally, due to time constraints and data processing limitations, I did not account for shifting county boundaries or weighing temperatures by population centers within counties. These refinements could improve precision, particularly in counties with large or unevenly distributed populations. Also, the fixed-effects design controls for unobserved, time-invariant county characteristics, but I did not incorporate models without fixed effects for comparison. Additionally, the analysis does not consider potential lagged effects of extreme heat on suicide risk, which could occur over multiple weeks or months.

A further limitation concerns the definition of EHDs. The 85th percentile threshold is computed over the full 30-year study period (1989–2019) for each county-month, meaning that by construction approximately 15% of days in the historical distribution exceed it. Because the threshold is static, the number of EHDs in a given county-month reflects year-to-year deviations from the long-run norm rather than secular warming trends, and EHD counts do not exhibit a strong upward trend over the study period even as average temperatures have risen. This property is a feature of the identification strategy: the causal estimates rely on idiosyncratic, within-county fluctuations in heat exposure that are plausibly exogenous to confounding trends in mental health, economic conditions, or health policy. However, it also

means the estimates cannot be straightforwardly extrapolated to project the mental health burden of long-run climate change, as an absolute or rolling-window threshold would better capture secular shifts in heat exposure. The 30°C absolute component of the EHD definition partially addresses this, but binds primarily in the hottest counties. Consequently, this measure may underestimate the cumulative burden of rising temperatures over the study period, and the true effects of climate-driven heat exposure on mental health are likely more severe than reported here.

Additionally, the EHD measure relies solely on dry-bulb temperature and does not incorporate humidity. Composite heat indices that combine temperature and relative humidity, such as the heat index or wet-bulb globe temperature, may better capture physiologically relevant heat stress, as high humidity impairs evaporative cooling and can amplify the health consequences of elevated temperatures [5].

5 Future Directions

Future research should expand upon this work by examining the effects of temperature extremes across all months of the year, potentially uncovering significant impacts during transitional seasons such as spring and fall. Additional studies incorporating dynamic models and lagged temperature effects would provide deeper insights into the temporal dynamics of heat-related mental health outcomes. Furthermore, refining temperature measures by weighting for population centers and accounting for shifting county boundaries could enhance precision and robustness. Adopting composite heat metrics that incorporate humidity, such as the heat index or wet-bulb globe temperature, could also yield a more physiologically meaningful measure of heat exposure. Lastly, exploring alternative temperature thresholds or extreme definitions and their varying health impacts would further elucidate how incremental climatic shifts influence mental health outcomes.

6 Replication

This analysis relies on the Restricted-Use Vital Statistics Data from the Centers for Disease Control and Prevention. Access to this data can be obtained by following the instructions at which can be found here: [12]. Other public data and my code can be found [here](#).

7 Appendix

Figure A.1: Average Monthly Suicide Rate(Deaths per Million) by County




Figure A.2: Average Summer Percentile Thresholds($^{\circ}\text{C}$) by County




Figure A.3: Trends in suicide rates and extreme heat days across urban and rural areas.




Figure A.4: Effect of one additional EHD on monthly suicide rates, by subgroup.




Table 1: ICD-9 and ICD-10 Codes Included in Mental Health-Related Mortality

Cause of Death	ICD-9 Codes	ICD-10 Codes
Suicides	E950-E959	Intentional self-harm: U03, X60-X84, Y87.0
Injuries of Undetermined Intent	Injury undetermined whether accidentally or purposely inflicted: E980-E989	Events of undetermined intent: Y10-Y34, Y87.2, Y89.9
Accidental Poisonings	Accidental poisoning: E850-E869	Accidental poisoning and exposure to noxious substances: X40-X49
Accidental Drownings	Accidental drowning and submersion: E910	Accidental drowning and submersion: W65-W74
Accidental Firearms	Accident caused by handgun: E922.0 Accident caused by other firearms: E922.1-E922.9	Accidental discharge of firearms: W32-W34
Train Accidents	Railway accidents: E800-E807 Motor vehicle-train collisions: E810	Railway accidents: V05, V15, V80.6, V81.2-V81.9 Motor vehicle-train collisions: V25, V35, V45, V55, V65, V75, V81.0-V81.1, V87.6, V88.6

Table 2: Linear Estimates

Region	Subgroup	Estimate (SE)	N
All	Full Population	0.091*** (0.013)	280639
	Male	0.154*** (0.021)	273493
	Female	0.029** (0.012)	270697
	Age 0–24	0.05*** (0.014)	118321
	Age 25–64	0.123*** (0.021)	248305
	Age 65+	0.068* (0.04)	277220
	White/NH	0.091*** (0.018)	268352
	NW/Hispanic	0.124*** (0.035)	168295
	HS or less	0.098** (0.044)	166512
	College+	0.064*** (0.027)	142230
Urban	Full Population	0.089*** (0.016)	73668
	Male	0.152*** (0.025)	73193
	Female	0.028* (0.015)	72902
	Age 0–24	0.048*** (0.015)	48897
	Age 25–64	0.123*** (0.024)	71476
	Age 65+	0.064 (0.049)	73324
	White/NH	0.09*** (0.022)	70964
	NW/Hispanic	0.121*** (0.035)	59547
	HS or less	0.127** (0.053)	44942
	College+	0.073** (0.03)	42168
Rural	Full Population	0.093*** (0.024)	206971
	Male	0.152*** (0.042)	200300
	Female	0.031 (0.025)	197795
	Age 0–24	0.074 (0.049)	69424
	Age 25–64	0.116*** (0.041)	176829
	Age 65+	0.075 (0.073)	203896
	White/NH	0.094*** (0.032)	197388
	NW/Hispanic	0.125 (0.12)	108748
	HS or less	— — 0.008 (0.069)	121570
	College+	0.092 (0.06)	100062

Notes: This table shows linear regression estimates of the effect of extreme heat days on suicide rates per million. Standard errors are in parentheses. All models include county-month, county-year, and year-month fixed effects, and are weighted by 2000 county population. Standard errors are clustered at the county level.

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Table 3: Poisson Estimates

Region	Subgroup	IRR (SE)	N
All	Full Population	1.004*** (0.001)	211090
	Male	1.004*** (0.001)	193568
	Female	1.002 (0.001)	123330
	Age 0–24	1.005** (0.002)	73148
	Age 25–64	1.004*** (0.001)	173273
	Age 65+	1.003 (0.002)	105106
	White/NH	1.003*** (0.001)	193367
	NW/Hispanic	1.005*** (0.002)	75249
	HS or less	1.003 (0.002)	103545
	College+	1.005* (0.003)	57557
Urban	Full Population	1.004*** (0.001)	67301
	Male	1.005*** (0.001)	64643
	Female	1.002 (0.002)	51888
	Age 0–24	1.005** (0.002)	38406
	Age 25–64	1.004*** (0.001)	62040
	Age 65+	1.003 (0.002)	44665
	White/NH	1.003*** (0.001)	63530
	NW/Hispanic	1.005*** (0.002)	37765
	HS or less	1.003 (0.002)	36532
	College+	1.005* (0.003)	27118
Rural	Full Population	1.003** (0.001)	143789
	Male	1.003** (0.001)	128925
	Female	1.002 (0.002)	71442
	Age 0–24	1.006* (0.003)	34742
	Age 25–64	1.003* (0.002)	111233
	Age 65+	1.002 (0.003)	60441
	White/NH	1.002 (0.001)	129837
	NW/Hispanic	1.006* (0.003)	37484
	HS or less	1.001 (0.002)	67013
	College+	0.999 (0.004)	30439

Notes: This table reports Poisson regression estimates of the effect of extreme heat days on suicide counts. Results are presented as incidence rate ratios (IRRs). Standard errors are in parentheses. All models include county-month, county-year, and year-month fixed effects, and are weighted by 2000 county population. Standard errors are clustered at the county level.

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

References

- [1] Boaz Abramson, Job Boerma, and Aleh Tsyvinski. Macroeconomics of mental health. *National Bureau of Economic Research*, 2024. doi:10.3386/w32354.
- [2] Environmental Protection Agency. Climate change indicators: Heat waves. <https://www.epa.gov/climate-indicators/climate-change-indicators-heat-waves>.
- [3] Environmental Protection Agency. Extreme heat. <https://www.epa.gov/climatechange-science/extreme-heat>.
- [4] Diane Alexander and Molly Schnell. Just what the nurse practitioner ordered: Independent prescriptive authority and population mental health. *Journal of Health Economics*, 2019. doi:10.1073/pnas.1518393112.
- [5] G. Brooke Anderson, Michelle L. Bell, and Roger D. Peng. Methods to calculate the heat index as an exposure metric in environmental health research. *Environmental Health Perspectives*, 2013. doi:10.1289/ehp.1206273.
- [6] Helen Berry, Kathryn Bowen, and Tord Kjellstrom. Climate change and mental health: a causal pathways framework. *International Journal of Public Health*, 2009. doi:10.1007/s00038-009-0112-0.
- [7] Stephen Bridges, Paul Yip, Krig Yang, and et al. Seasonal changes in suicide in the united states, 1971 to 2000. *Perceptual and Motor Skills*, 2005. doi:10.2466/pms.100.3c.920-924.
- [8] Lukas Brunner and Aiko Voigt. Pitfalls in diagnosing temperature extremes. *Nature Communications*, 2024. doi:10.1038/s41467-024-46349-x.
- [9] Marshall Burke, Felipe González, Patrick Baylis, Sam Heft-Neal, Ceren Baysan, Sanjay Basu, and Solomon Hsiang. Higher temperatures increase suicide rates in the united states and mexico. *Nature Climate Change*, 2018. doi:10.1038/s41558-018-0222-x.
- [10] Tamma A. Carleton and Solomon M. Hsiang. Social and economic impacts of climate. *Science*, 2016. doi:10.1126/science.aad9837.
- [11] Anne Case and Angus Deaton. Rising morbidity and mortality in midlife among white non-hispanic americans in the 21st century. *Proceedings of the National Academy of Sciences*, 2015. doi:10.1073/pnas.1518393112.
- [12] Centers for Disease Control and Prevention - National Center for Health Statistics. National vital statistics system, restricted-use vital statistics data: Deaths (mortality)—multiple cause of death, states and all counties, 2021. <https://www.cdc.gov/nchs/nvss/nvss-restricted-data.htm>.
- [13] C. Christodoulou, A. Douzenis, F. C. Papadopoulos, A. Papadopoulou, G. Bouras, R. Gournellis, and L. Lykouras. Suicide and seasonality. *Acta Psychiatrica Scandinavica*, 2012. doi:10.1111/j.1600-0447.2011.01750.x.

- [14] Centers for Disease Control and Prevention. Suicide mortality in the united states, 2002–2022, 2014. <https://www.cdc.gov/nchs/products/databriefs/db398.htm>.
- [15] Centers for Disease Control and Prevention. Web-based injury statistics query and reporting system: Leading cause of death visualization tool, 2020, 2020. <https://wisqars.cdc.gov/data/lcd/home>.
- [16] Andrew Gelman and Jesse Auerbach. Age-aggregation bias in mortality trends. *Proceedings of the National Academy of Sciences*, 2016. doi:10.1073/pnas.1523465113.
- [17] Solomon Hsiang, Marshall Burke, and Edward Miguel. Quantifying the influence of climate on human conflict. *Science*, 2013. doi:10.1126/science.1235367.
- [18] Asha Ivey-Stephenson, Alex Crosby, Shane Jack, Tadesse Haileyesus, and Marcie-Jo Kresnow-Sedacca. Suicide trends among and within urbanization levels by sex, race/ethnicity, age group, and mechanism of death—united states, 2001–2015. *MMWR Surveillance Summaries*, 2017. doi:10.15585/mmwr.ss6618a1.
- [19] Stephan Lim, Theo Vos, Abraham Flaxman, and et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the global burden of disease study 2010. *The Lancet*, 2013. doi:10.1016/S0140-6736(12)61766-8.
- [20] Rafael Lozano, Mohsen Naghavi, Kyle Foreman, and et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the global burden of disease study 2010. *The Lancet*, 2013. doi:10.1016/S0140-6736(12)61728-0.
- [21] David Molitor, Jamie Mullins, and Corey White. Air pollution and suicide in rural and urban america: Evidence from wildfiresmoke. *Proceedings of the National Academy of Sciences*, 2023. doi:10.1073/pnas.2221621120.
- [22] Jamie T. Mullins and Corey White. Temperature and mental health: Evidence from the spectrum of mental health outcomes. *Journal of Health Economics*, 2019. doi:10.1016/j.jhealeco.2019.102240.
- [23] National Center for Health Statistics. 2013 national center for health statistics urban-rural classification scheme, 2013. <https://www.cdc.gov/nchs/dataaccess/urbanrural.htm>.
- [24] National Institute of Mental Health. Mental illness. <https://www.nimh.nih.gov/health/statistics/mental-illness>.
- [25] National Institute of Mental Health. Suicide. <https://www.nimh.nih.gov/health/statistics/suicide>.
- [26] World Health Organization. Heat and health. <https://www.who.int/news-room/fact-sheets/detail/climate-change-heat-and-health>.

- [27] World Health Organization. Suicide. <https://www.nimh.nih.gov/health/statistics/suicide>.
- [28] Stephanie Pappas. How heat affects the mind. *American Psychological Association*, 2024.
- [29] Johnathan Patz, Howard Frumpkin, and Tracey Holloway. Climate change challenges and opportunities for global health. *Journal of the American Medical Association*, 2014. doi:10.1001/jama.2014.13186.
- [30] Julien Periard, Gavin Travers, Sebastien Racinais, and Michael Sawka. Cardiovascular adaptations supporting human exercise-heat acclimation. *Autonomic Neuroscience*, 2016. doi:10.1016/j.autneu.2016.02.002.
- [31] Martin Prince, Vikram Patel, Shekhar Saxena, Mario Maj, Joanna Maselko, Michael Phillips, and Atif Rahman. No health without mental health. *The Lancet*, 2016. doi:10.1016/S0140-6736(07)61238-0.
- [32] PRISM Climate Group. Prism climate data, 2022. <https://prism.oregonstate.edu/>.
- [33] Moustaq Rony and Hasnat Alamgir. High temperatures on mental health: Recognizing the association and the need for proactive strategies—a perspective. *Health Science Reports*, 2023. doi:10.1002/hsr2.1729.
- [34] SEER Program, National Cancer Institute. Us county population data, 1969–2020, 2022. <https://seer.cancer.gov/popdata/index.html>.
- [35] Jefferey Swanson, Richard Bonnie, and Paul Appelbaum. Getting serious about reducing suicide more “how” and less “why”. *Journal of the American Medical Association*, 2015. doi:10.1001/jama.2015.15566.
- [36] Sebastian Trautmann, Jurgen Rehm, and Hans-Ulrich Wittchen. The economic costs of mental disorders: Do our societies react appropriately to the burden of mental disorders? *European Molecular Biology Organization*, 2016. doi:10.15252/embr.201642951.
- [37] USDA Economic Research Service. Educational attainment for the u.s., states, and counties, 1970–2020, 2022. <https://www.ers.usda.gov/data-products/county-level-data-sets/download-data/>.
- [38] Nick Watts, Neil Adger, Paolo Agnolucci, and et al. Health and climate change: policy responses to protect public health. *The Lancet*, 2015. doi:10.1016/S0140-6736(15)60854-6.