

# **A Space and Solar Physics Data Model**

**from the SPASE Consortium**

**Version: 1.0.1**

Release Date: 2006-01-03

## **Consortium Members:**

### **Augsburg College**

Mark Engebretson, <engebret@augsborg.edu>  
Noel Petit, <petit@augsborg.edu>

### **Centre de Données de la Physique des Plasmas (CDPP)**

Michel Gangloff, <gangloff@cesr.fr>  
Christopher Harvey, <christopher.harvey@cesr.fr>  
Claude Huc, <claudio.huc@cnes.fr>  
Thierry Levoir, <thierry.levoir@cnes.fr>

### **Istituto Nazionale di Astrofisica (INAF)**

Kevin Reardon, <kreardon@arcetri.astro.it>

### **Japan Aerospace eXploration Agency (JAXA) - STP/Ehime**

Yasumasa Kasaba, <kasaba@stp.isas.jaxa.jp>  
Ken T. Murata, STP/Ehime, <murata@cite.ehime-u.ac.jp>

### **Jet Propulsion Laboratory (JPL)**

Dan Crichton, <dan.crichton@jpl.nasa.gov>  
Steven Hughes, <j.steven.hughes@jpl.nasa.gov>

### **John Hopkins University/Applied Physics Laboratory (JHU/APL)**

Rose Daley, <rose.daley@jhuapl.edu>  
Brand Fortner, <brand.fortner@jhuapl.edu>  
Stu Nylund, <stu.nylund@jhuapl.edu>  
Jon Vandergriff, <jon.vandergriff@jhuapl.edu>  
Michele Weiss, <michele.weiss@jhuapl.edu>

### **Goddard Space Flight Center (GSFC)**

Dieter Bilitza (RSTX), <bilitza@mail630.gsfc.nasa.gov>  
Bobby Candey, <candey@mail630.gsfc.nasa.gov>  
Carl Cornwell (Aquilent), <carl.cornwell@aquilent.com>  
Joe Gurman, <gurman@grace.nascom.nasa.gov>  
Joe Hourcle (EITI), <oneiros@grace.nascom.nasa.gov>  
Mona Kessel, <kessel@ndadsb-f.gsfc.nasa.gov>  
Joe King, <jking@mail630.gsfc.nasa.gov>  
Terry Kucera, <kucera@stars.gsfc.nasa.gov>  
Bob McGuire, <rmcguire@pop600.gsfc.nasa.gov>  
Lou Reich (CSC), <lreich@pop500.gsfc.nasa.gov>  
Aaron Roberts, <roberts@vayu.gsfc.nasa.gov>  
Don Sawyer, <donald.sawyer@gsfc.nasa.gov>  
Dave Sibeck <dsibeck@pop600.gsfc.nasa.gov>  
Adam Szabo, <aszabo@pop600.gsfc.nasa.gov>  
Jim Thieman (NSSDC), <james.r.thieman@nasa.gov>  
Karen North, <Karen.C.North@nasa.gov>  
Aaron Smith (Aquilent), <aaron.smith@aquilent.com>  
Isaac Verghese (Aquilent), <Isaac.Verghese@aquilent.com>  
Vasili Rezapkin (Aquilent), <vasili.rezapkin@aquilent.com>

**National Aeronautics and Space Administration (NASA) HQ**

Joe Bredekamp, <jbredeka@mail.hq.nasa.gov>

Chuck Holmes, <cholmes@mail.hq.nasa.gov>

**National Oceanic and Atmospheric Administration (NOAA)**

Eric Kihn, <eric.a.kihn@noaa.gov>

**QSS Group, Inc.**

Ed Bell, <ed.bell@gsfc.nasa.gov>

**Rutherford Appleton Laboratory (RAL)**

Chris Perry, <c.h.perry@rl.ac.uk>

Phil Richards, <P.J.Richards@rl.ac.uk>

**Stanford University**

Rick Bogart, <rbogart@stanford.edu>

**Southwest Research Institute (SwRI)**

Joey Mukherjee, <jmukherjee@swri.org>

Dave Winningham, <david@cluster.space.swri.edu>

**University of California, Los Angeles (UCLA)**

Steven Joy, <sjoy@igpp.ucla.edu>

Todd King, <tking@igpp.ucla.edu>

Ray Walker, <rwalker@igpp.ucla.edu>

**Table of Contents**

1. Executive Summary .....	1
2. Introduction .....	2
2.1. What is a data model? .....	2
2.2. What is its use? .....	2
2.3. What process led to the current Data Model? .....	3
2.4. The Data Model in a Virtual Observatory context .....	3
3. Data Model overview and general concepts .....	5
3.1. Resources .....	5
3.2. Numerical Data Resources .....	6
3.3. Display Data Resources .....	8
3.4. Catalogue Resources .....	8
3.5. Other Resource Types .....	8
3.6. Future Direction .....	8
3.7. General Considerations .....	8
4. The Data Model presented hierarchically .....	10
5. Guidelines for Metadata Descriptions of Products .....	15
6. Examples .....	16
7. Definitions of the Data Model Terms .....	18
8. Enumeration of Selected Quantities .....	46
9. Appendix A - Graphical Presentation of the SPASE Schema .....	62
10. Bibliography .....	70
11. Index .....	71
12. Change History .....	72

## 1. Executive Summary

The Solar and Space Physics communities need a unified data environment to facilitate finding, retrieving, formatting, and obtaining basic information about data essential for their research. With the increasing requirement for data from multiple sources, this need has become acute. A unified method to describe data and other resources is the key to achieving this unified environment. The SPASE (Space Physics Archive Search and Extract) Data Model provides a basic set of terms and values organised in a simple and homogeneous way, to facilitate access to Solar and Space Physics resources. This initial version of the Model is primarily concerned with numerical data and the display data based on it, but the model will be extended to include descriptive terms for other resources including data from empirical or physics based models and software for models and data interpretation. The SPASE Data Model is comparable to the data models developed by the Planetary Data System (PDS) and the International Virtual Observatory Alliance (IVOA) for planetary and astronomical data, respectively. The SPASE Model will provide the detailed information at the parameter level required for Solar and Space Physics applications.

The SPASE consortium is an international team of space and solar physicists and information scientists. It first examined many existing data models, but found none to be adequate. A set of terms based on a half-dozen or so of the most complete of such models was refined based on applying the model at various levels of detail to a large number of existing products to arrive at the current version. The major creators of SPASE-based product descriptions are expected to be domain-based Virtual Observatories ("VxOs"), data centers, and individual data and model providers. The SPASE Data Model will continue to evolve in a controlled way as data and service providers and benefiting researchers suggest improvements to extend its framework of common standards. Success of the model will be measured by the extent of community support and use.

The present Data Model provides enough detail to allow a scientist to understand the content of Data Products (e.g., a set of files for 3 second resolution Geotail magnetic field data for 1992 to 2005), together with essential retrieval and contact information. A typical use would be to have a collection of descriptions stored in one or more related internet-based registries of products; these could be queried with specifically designed search engines which link users to the data they need. Initially, this Data Model stops short of providing the means to access specific "granules" (individual files) or subsets of them, but this extension is foreseen. Other extensions of the model will include the description of software tools and services useful to read and analyse the data, and of physically- or empirically-based models to interpret it.

This document provides potential users of SPASE with the Data Model for review and use. Sections 2 and 3 provide an overview of the origins and the concepts of the data model. Section 4 presents the set of elements in a hierarchy that shows the natural relationships among them. This is followed by usage suggestion and pedagogic examples in Section 5 and 6, and by the complete set of definitions of terms and enumerated lists in Section 7.

## 2. Introduction

The Space and Solar Physics community is now addressing fundamental questions concerning the plasma and magnetic environment of the Sun, Heliosphere, and planets. We seek to understand everything from "microscopic" phenomena such as magnetic reconnection and turbulent energy dissipation to global issues such as how solar events are related to potentially damaging electric currents and energetic particles in the vicinity of the Earth. Multispacecraft and ground observatory investigations are becoming the norm, but there has been little corresponding unification of data access, formats and tools, resulting in a great deal of time being spent finding, retrieving and reformatting data. The key to reducing this inefficiency is a uniform way to describe adequately what exists. This is the purpose of the SPASE Data Model, which is intended to do for Space and Solar Physics what the Planetary Data System (PDS) is doing for Planetary Science, and what the International Virtual Observatory Alliance (IVOA) is doing for Astronomy and Astrophysics.

Astronomy, Planetary Science, and Space and Solar Physics have developed their data models with different objectives and constraints. Astronomy has objects defined by their direction, and makes much use of standard formats; PDS was developed for long-term archiving; while SPASE is oriented towards data searching and exploitation. PDS often does a better job of describing technical information such as how the data were processed, the form of the data etc., while SPASE includes a better physical description, which facilitates rapid data retrieval and exploitation. Over time, the Astronomy, PDS, and SPASE models may converge. The increasing use of, for example, time series in Astronomy and images in Space Physics may lead to stronger connections between the efforts in these areas. There will also be a need for SPASE to understand Earth Science data models, as Space Weather studies reveal the coupling of the larger plasma environment to terrestrial effects.

### 2.1. What is a data model?

A data model is a set of terms and their relationships that capture the essential concepts of a given domain. The Data Model presented here can be used to describe the scientific relevance of products resulting from observation and modeling in the domain of solar and space physics. These products typically consist of related collections of files that will be accessed, in whole or in part, by science users. Types of products include numerical datasets, display data plots, images, software, documentation, and event lists ("catalogues"). Products are a subset of a larger class of "Resources" that includes Spacecraft, Instruments, Repositories, and even People, that can be described by their own set of terms, and then referred to in product descriptions, rather than being repeatedly described in each product. This data model includes terms relevant to all "Sun-Earth Connection" domains, but it does not try to systematically include Earth Science terms. We also defer to later the question of a uniform description of "service resources" such as web-based format translators or display tools. We initially intend a level of description that will allow a scientist to use the data retrieved, and will add later a set of terms to describe specific access methods for direct data retrieval. At all stages, broad community input and feedback is essential.

### 2.2. What is its use?

The Data Model provides a set of terms that, ideally, can be mapped onto the terms used for specific products, thus providing a uniform means of access and description. The data model is intended to provide the cornerstone of one or more "Virtual Observatories" that will link broad ranges of solar and space physics in a natural way. More specifically, the data model should:

- (1) Provide a way of registering products using a standard set of terms that allow the products to be found with simple searches and described so that users can determine their utility for a specific purpose;
- (2) Allow searching for products containing particular physical quantities (e.g., magnetic field; spectral irradiance) that are variously represented in a diverse array of data products; and
- (3) Create a means of mapping comparable variables from many products onto a common set of terms so that visualization, analysis, and higher-order query tools and services can be used on all of them without regard to the origin of the data.

The accomplishment of these tasks requires "middleware" (either at the provider or in a VxO) that understands product registries and performs the translations needed to map the idiosyncratic product and parameter names of each repository onto the standard terms. This intermediate layer, which can take many forms for different purposes, will provide the links necessary to connect user applications and search-and-retrieval front ends to data repositories. Ultimately, the data environment centered on the data model will involve a number of software tools as well, linked together as internet-based services or other means. Specific software tools and documentation associated with products will be straightforwardly accessible. This "system" has the potential to provide capabilities that can aid even expert users of a particular dataset (e.g., on-the-fly coordinate transformations, the ability to merge datasets from different instruments, easy reference to related indices or other data), in addition to providing the broad access needed to solve global problems in Sun-Earth connection physics. Success will require a concerted cooperative effort across disciplines. Existing efforts in Space and Solar Physics as well as in other areas such as Earth and Planetary Sciences and Astronomy will guide the work.

### **2.3. What process led to the current Data Model?**

The data model presented here has grown from the efforts begun a number of years ago that became formalized in regular teleconferences of a group of interested data providers, including scientific and technical representatives of some of the largest data holdings in the US, Europe, and Japan. As the effort to provide seamless access to distributed data proceeded, it became clear that the data model efforts were central. Thus, in March of 2003 a meeting of many of the people in the Contributors list at the beginning of this document was convened to begin the data model construction in earnest. The initial effort involved collecting terms from CDPP, SWRI, NSSDC, ISTEP, and other sets to form a starting point. Two years of teleconferences, e-mailed revisions, and occasional face-to-face efforts, along with the application of the terms to specific cases, has led to the present document.

The general philosophy of the Data Model is to describe products using a natural taxonomy of data sources and of the physical world as represented in actual or potential datasets and models. The resulting Data Model has been put to many tests, but will have to evolve as new products are considered.


### **2.4. The Data Model in a Virtual Observatory context**

The current conceptual model is intended to serve as the basis for interoperability between independent data systems. It is an attempt to capture the various concepts that are used to represent the knowledge to be shared in the Space and Solar Physics domains. These concepts are used to convey queries and responses between the user and various remote data systems. Many such

systems, in diverse fields, are now functioning or planned. The paradigm for this interoperability is the "Virtual Observatory", originally coined by the astronomical community as a way to allow any researcher, anywhere in the world, to access all known observations of a chosen patch of the sky using the Internet to query repositories of information distributed around the world at data centers and observatories.

The concepts presented in this document are independent of any implementation, but we have in mind a likely map of the eventual data environment. Figure 1 illustrates a possible architecture; the details will depend on the outcome of a number of current efforts and their coordination. The web site <http://lwsde.gsfc.nasa.gov> gives a guide to many currently active projects and a great deal of background information. Of particular interest there is the document entitled, "A Framework for Space and Solar Physics Virtual Observatories."

Figure 1 is very schematic, but the "Access Points" could be discipline specific "VxOs" that aggregate the repositories they serve, making them all appear as one. This is now done by the VSO for solar data. The VSO can then be treated as a repository by other services (VSPO and EGSO currently do this). The "Gateways" in the figure would be places where large numbers of products are registered with pointers to Access Points, Repositories, and other services (not shown) as needed. With the right connections and a common language, the data environment becomes very rich and flexible. Note that more conventional access to any of the Repositories could continue as before.



A prototypical usage scenario is:

1. The scientist uses an application (e.g., a browser or IDL) to define his query in terms of



the SPASE model.

2. The application contacts a relevant participating access point or gateway that may generate a response (e.g. URLs of files or services, or metadata about these), or may pass the query to other access points or gateways to obtain further information before responding to the query.
3. The application uses the replies either to present metadata to the user or to request the resource. A request may go directly to a repository or an access point, or may be routed through a gateway for additional services.
4. The requested resources might be processed in various ways at the repository, the access point, the gateway, or by a separate service. For example, the system may assemble a collection of resources, subset the data, generate a graphic, or reformat the resource prior to delivering it to the application.

We envision that different control authorities will maintain different aspects of the Data Model. For example, the list of observatory names is maintained by NSSDC, the definition of prime meridian used in describing local latitude and longitude is set by IAU, and the accepted format for time representation is defined by ISO. There will need to be a community-approved group to be the central authority on the Data Model; SPASE is currently serving as a prototype for this role.

It remains to be determined the extent to which registries of products will be centralized such that one location will contain the latest list obtained from registered data providers. There should at least be a common format, if only keyword=value lists, for the exchange of product registries. XML schema are likely to provide the preferred framework.

### **3. Data Model overview and general concepts**

As noted in the introduction, this Data Model focuses on describing Products, which are a subset of Resources. In general, each Resource Type consists of a similar collection of things that share a common descriptive hierarchy. Data and Display Products (the latter being pre-rendered Plots and Images based on data) are typically defined as collections of files that differ only in the time of the observations to which they refer. Note that what is delivered to the user may be generated dynamically from underlying files, and thus can be more variable: for a particular product, modes may change with time of observation, and calibrations with time of processing. Thus, in general, the delineation of a product is up to the providers. All of the terms in the Model require a precise definition to be useful, and these are provided in the Dictionary. Note that in other data models what SPASE would call "files" or "granules" are sometimes referred to as "products" or "datasets," so it is important to keep in mind that these terms are used here to refer to complete collections of like data.

The SPASE Data Model describes Data Products by stating what was observed and where and when the data were obtained, as well as giving the source of the observations, and the location (repository), format and other technical aspects of the data.

#### **3.1. Resources**

At top level of the present Data Model is the Resource Type. These consist of the Product Types:

Numerical,

## Data Display, and Catalogue

and the Resource Types that support these:

Observatory,  
Repository,  
Instrument, and  
Person.

Each of these types has its own set of terms. In the case of the non-product resources, the present version of the dictionary provides only a basic set of terms that may be augmented if it proves useful to have more detail. Any resource may have a URL that will provide the user with more information to supplement the metadata.

A set of Elements (the general model term for what could roughly be termed a keyword) is common to all Resource descriptions; they are grouped for convenience of presentation in a Resource Header, consisting primarily of:

Resource Name  
Provider Resource Name  
Description  
Acknowledgement  
Contact  
Alias  
Information URL

For Products, a similar set is grouped as Access Information, which can be replicated with the appropriate changes to describe copies of the product which may differ only in format or encoding. The Access Information provides the:

Repository Name  
Availability  
Access URL  
Caveats  
Format  
Encoding

These general elements are often free text, although Instrument Name, Format, and various other terms are from a specific (often extensible) list. Having enumerated lists will allow searches to be more efficient, so that, for example, a simple query will isolate all registered products from a particular Region without worrying about many more-or-less equivalent terms.

### **3.2. Numerical Data Resources**

Since Numerical Data Resources are the key to most science investigations, our focus has been on providing a Data Model that will provide users with a means to exploit these data. It will be useful to agree on a uniform means for accessing the data (standard conventions for how to describe file naming and variable access methods), but initially our focus is on the scientific content of the data. We have initial sets of terms for the other Product Types, but these are less detailed; in many cases we expect much less detail will be needed even in the more final form.

Numerical Data Resources or Products (called "Datasets" in some systems) are described by the Headers plus other elements, including:

- Observatory ID (link to Resource description)
- Instrument ID (also a link)
- Measurement Type (the general category, such as Thermal Plasma)
- Temporal Description (time range available, resolution)
- Physical Parameter (optional, but very useful)

Each Physical Parameter is a description for a the physical quantity that the provider wishes to advertise in the product. This could consist of one entry describing image data as being a full-Sun image in white light, or it could contain many entries for the various parameters (density, temperature, velocity, variance in these, error bars, and quality flags) for a thermal plasma product. Each Physical Parameter would be described by a standard set of terms including:

- Name
- Description
- Caveats
- Cadence
- Units
- Units Conversion
- Coordinate System
- Dimension
  - Size
  - Description
- Measured
- Support

and then a choice of the parameter type (Support or Measured). Measured parameters are described by an Entity, and a Quantity with one or more Qualifiers. Note that the independent variables (e.g., energy and angular ranges for particle distribution functions) are specified in the current Data Model in the Description of the Dimensions of a Parameter. Later versions of the model should allow for a more direct numerical specification of independent variable ranges.

The list of Entities, which form the basic set of Measured parameters, is based on the natural taxonomy of the physical world, but also corresponds to entities in known products. The list of entities include: Photons, which are electromagnetic fields; Fields, distinguished from Photons by being measured as time series; Particles, which are forms of matter; and Mixed, which are composite or derived quantities. Photons form optical or X-ray images, or radio waves when measured as amplitudes of signals in frequency bands.

The Physical Quantities, specific to each entity, are the two classical fields (electric and magnetic) and the associated electric potential: number/energy fluxes to describe distribution functions of particles (differential by bins); the first four moments of a particle distribution function; intensity, counts, and various derived quantities. It remains to be determined if we want a list of all the derived quantities contained in all relevant products; what appears below covers the many cases we have examined, but modifications will be needed as new products are described. Wave fields are described either as time series of fields (waveform), or in terms of "spectral density," although some products use terms such as "peak electric field" whose precise meaning may not be immediately evident. Ideally the description will become more precise when cast in SPASE terms.

Qualifiers of Physical Quantities can be combined, e.g., the variance of the x-component. The field-aligned component of the proton heat flux vector would be described as the parallel component of the heat flux of protons in magnetic coordinates.

The essential information for a Numerical Data Product description consists of what measured it, where to get it, its basic Measurement Type, whom to ask about it, and a few other simple descriptors. Of course, the more detailed the description, the more likely it is that users will understand what a product is and be more able and likely to use it.

### **3.3. Display Data Resources**

Display products are very similar to Numerical Data, since they are based on the data, but are preprocessed images in, for example, JPEG, GIF or PNG format that show a picture of a graph or of an object or region. Generally these products will be summaries or browse-level images of data plots, so less detail will be needed in their description. The convention chosen here is that, for example, the FITS file containing the data for a solar H-alpha image is part of a Numerical Data Product, whereas a GIF or JPEG produced from that file, which is more difficult to use for quantitative purposes, is part of a Display Data Product.

### **3.4. Catalogue Resources**

Catalogues can include complete listings of files and the times they cover, but this is not the main intent here. Rather, a Catalogue will typically consist of a set of start and stop times for "events" which can be anything from Coronal Mass Ejection occurrences in SOHO images, to intervals when data are available in a particular product, to identifications of when a spacecraft is in a particular region such as the Earth's magnetosheath. These types of catalogues are already being used for assistance in data searches by VSO and EGSO.

### **3.5. Other Resource Types**

The "nonproduct" Resource types are generally self-explanatory. In a more complete Data Model, further information could be given about Observatories, Instruments, etc., but for now such details are left to the links provided in Information URLs.

### **3.6. Future Direction**

Future versions of the SPASE Data Model will include terms for Software (tools for use of data), Models (physics- and empirically-based predictive schemes, often in the form of programs), Documents (focusing on support documents for other resources rather than on scientific papers), and Services (web-based format translation, coordinate transformation, plotting, visualization, etc.) It was felt that the standards for these resource types were not mature enough to be included in this release.

### **3.7. General Considerations**

The SPASE Entity-Relationship Tree of Section 3 has been produced from information held in a data base. This database also holds all the element definitions which are explained in Section 5 and tabulated in Section 6. From this database it is also possible to generate an XSD schema, which is the default encoding that has been developed for evaluating the Data Model. Thus, Resources are most naturally described by XML files, but this is not essential. In fact, the examples in this

document are presented in a keyword=value format to be more human readable.

A PDF version of this document can be downloaded from the SPASE site,  
<http://www.igpp.ucla.edu/spase/>

#### 4. The Data Model presented hierarchically

The taxonomy tree shows the inter-relationship of elements in the data model. This provides a "big picture" view of the SPASE data model. This taxonomy is implementation neutral. Details for each element are contained in the data dictionary.

Notes: Occurrence specifications are enclosed in parenthesis: 0 = optional, 1 = required, \* = zero or more, + = 1 or more

```

+ Spase (1)
|   + Version (1)
|   + Catalog (*)
|       + Resource ID (1)
|       + Resource Header (1)
|           + Resource Name (1)
|           + Provider Resource Name (*)
|           + Description (1)
|           + Acknowledgement (0)
|           + Contact (1)
|               + Person ID (1)
|               + Role (+)
|           + Alias (*)
|           + Information URL (*)
|               + Name (0)
|               + URL (1)
|               + Description (0)
|       + Observatory ID (*)
|       + Instrument ID (*)
|       + Phenomenon Type (1)
|       + Timespan (0)
|           + Start Date (1)
|           + End Date (0)
|       + Keyword (*)
|       + Input Resource ID (+)
|       + Access Information (1)
|           + Repository Name (1)
|           + Availability (0)
|               + Offline (0)
|               + Medium (0)
|               + Online (0)
|           + Access URL (+)
|               + Name (0)
|               + URL (1)
|               + Description (0)
|           + Caveats (*)
|           + Format (1)
|           + Encoding (0)
|           + Acknowledgement (1)
|   + Display Data (*)
|       + Resource ID (1)
|       + Resource Header (1)

```

			+ Resource Name (1)
			+ Provider Resource Name (*)
			+ Description (1)
			+ Acknowledgement (0)
			+ Contact (1)
			+ Person ID (1)
			+ Role (+)
			+ Alias (*)
			+ Information URL (*)
			+ Name (0)
			+ URL (1)
			+ Description (0)
			+ Provider Processing Level (0)
			+ Provider Version (0)
			+ Observatory ID (1)
			+ Instrument ID (1)
			+ Measurement Type (+)
			+ Temporal Description (0)
			+ Timespan (1)
			+ Start Date (1)
			+ End Date (0)
			+ Cadence (0)
			+ Exposure (0)
			+ Spectral Range (*)
			+ Display Cadence (0)
			+ Instrument Region (*)
			+ Surface (0)
			+ Near Surface (0)
			+ Magnetosphere (0)
			+ Heliosphere (0)
			+ Sun (0)
			+ Observed Region (*)
			+ Surface (0)
			+ Near Surface (0)
			+ Magnetosphere (0)
			+ Heliosphere (0)
			+ Sun (0)
			+ Keyword (*)
			+ Input Resource ID (*)
			+ Access Rights (1)
			+ Access Information (1)
			+ Repository Name (1)
			+ Availability (0)
			+ Offline (0)
			+ Medium (0)
			+ Online (0)
			+ Access URL (+)
			+ Name (0)
			+ URL (1)
			+ Description (0)
			+ Caveats (*)

			+ Format (1)
			+ Encoding (0)
			+ Acknowledgement (1)
			+ Numerical Data (*)
			+ Resource ID (1)
			+ Resource Header (1)
			+ Resource Name (1)
			+ Provider Resource Name (*)
			+ Description (1)
			+ Acknowledgement (0)
			+ Contact (1)
			+ Person ID (1)
			+ Role (+)
			+ Alias (*)
			+ Information URL (*)
			+ Name (0)
			+ URL (1)
			+ Description (0)
			+ Provider Processing Level (0)
			+ Provider Version (0)
			+ Observatory ID (1)
			+ Instrument ID (1)
			+ Measurement Type (+)
			+ Temporal Description (0)
			+ Timespan (1)
			+ Start Date (1)
			+ End Date (0)
			+ Cadence (0)
			+ Exposure (0)
			+ Spectral Range (*)
			+ Instrument Region (*)
			+ Surface (0)
			+ Near Surface (0)
			+ Magnetosphere (0)
			+ Heliosphere (0)
			+ Sun (0)
			+ Observed Region (*)
			+ Surface (0)
			+ Near Surface (0)
			+ Magnetosphere (0)
			+ Heliosphere (0)
			+ Sun (0)
			+ Physical Parameter (+)
			+ Name (0)
			+ Description (0)
			+ Caveats (0)
			+ Cadence (0)
			+ Units (0)
			+ Units Conversion (0)
			+ Coordinate System (0)
			+ Coordinate Representation (0)



			+ Coordinate System Name (0)
			+ Dimension (0)
			+ Size (1)
			+ Description (1)
			+ Measured (0)
			+ Field (0)
			+ Field Qualifier (1)
			+ Field Physical Quantity (1)
			+ Particle (0)
			+ Particle Type (1)
			+ Particle Qualifier (0)
			+ Particle Physical Quantity (1)
			+ Photon (0)
			+ Photon Qualifier (1)
			+ Photon Physical Quantity (1)
			+ Mixed (0)
			+ Support (0)
			+ Keyword (*)
			+ Input Resource ID (*)
			+ Access Rights (1)
			+ Access Information (1)
			+ Repository Name (1)
			+ Availability (0)
			+ Offline (0)
			+ Medium (0)
			+ Online (0)
			+ Access URL (+)
			+ Name (0)
			+ URL (1)
			+ Description (0)
			+ Caveats (*)
			+ Format (1)
			+ Encoding (0)
			+ Acknowledgement (1)
			+ Instrument (*)
			+ Resource ID (1)
			+ Resource Header (1)
			+ Resource Name (1)
			+ Provider Resource Name (*)
			+ Description (1)
			+ Acknowledgement (0)
			+ Contact (1)
			+ Person ID (1)
			+ Role (+)
			+ Alias (*)
			+ Information URL (*)
			+ Name (0)
			+ URL (1)
			+ Description (0)
			+ Instrument Name (1)
			+ Instrument Type (0)

```

|      |      + Investigation Name (1)
|      |      + Acronym (1)
|      | + Observatory (*)
|      |      + Resource ID (1)
|      |      + Resource Header (0)
|      |      |      + Resource Name (1)
|      |      |      + Provider Resource Name (*)
|      |      |      + Description (1)
|      |      |      + Acknowledgement (0)
|      |      |      + Contact (1)
|      |      |      |      + Person ID (1)
|      |      |      |      + Role (+)
|      |      |      + Alias (*)
|      |      |      + Information URL (*)
|      |      |      |      + Name (0)
|      |      |      |      + URL (1)
|      |      |      |      + Description (0)
|      |      + Observatory Name (1)
|      |      + Observatory Group (0)
|      | + Person (*)
|      |      + Resource ID (1)
|      |      + Person Name (0)
|      |      + Organization Name (1)
|      |      + Address (0)
|      |      + Email (*)
|      |      + Phone Number (*)

```

## 5. Guidelines for Metadata Descriptions of Products

The following sections describe the details of the SPASE Data Model, especially the metadata used to describe data. There is a richness in the available metadata that allows very detailed descriptions of products. Many of the types of metadata may not apply in your case or you may not need much detail to adequately describe your data holdings. But it must be remembered that the better data are described, the easier they will be to use.

To determine what level of detail is needed, we recommend considering not only what the user needs to find the correct data, but also what is needed to know if the data will be useful for the requestor's purpose. The user might get this information by contacting you, but if the data were moved somewhere else and only the data description were available to determine the utility, consider if the user would have sufficient information to know if this is the right data set and what problems might be associated with the use of these data. Documentation may be added as a Resource Type to a future version of the SPASE Data Model, but for now we recommend using "Information URLs," available for each resource, to provide links to more detailed information about data products and their sources.

In summary, products need not be described in minute detail, but users will need, at minimum, information for assessing what the data products represent and where to find them. Of course it is also useful to include information on how the data can be applied and common pitfalls in their use, but the first need is to make the products usefully visible.

## 6. Examples

As an example let us consider a collection of numerical data from the magnetometer on the ACE spacecraft. This data set has been averaged to 1 minute intervals (cadence) and spans the beginning of the mission to the end of 2004 (1997-09-01 through 2004-12-31). The ACE spacecraft orbits the L1 point between the Earth and the Sun. The presented URLs are fictitious and will not direct you to the actual data. It's presented in the Object Definition Language (ODL) which is a Keyword=Value representation with a reserved keyword of "OBJECT" used to indicate groupings. In practice the preferred format will likely be XML because of its wide community support. The example is provided in ODL because it is more human readable.

```

VERSION                                = "1.0.0"
OBJECT                                = NUMERICAL_DATA
  RESOURCE_ID                          =
"UCLA::ACE-SW-MFI-4-RDR-PROC-V1.0:ACEMAG200301:1"
  MEASUREMENT_TYPE                    = "Magnetic Field"
  OBSERVATORY_ID                      = "ACE"
  INSTRUMENT_ID                      = "ACE_MFI"
  PROVIDER_ID                        = "UCLA"
  ACCESS_RIGHTS                      = "Open"
  OBJECT                              = RESOURCE_HEADER
  RESOURCE_NAME                      = "ACEMAG200301"
  ACKNOWLEDGEMENT                    = "
    User will acknowledge the data producer and instrument P.I. in any
    publication resulting from the use of these data."
  DESCRIPTION                        = "
    ACE MFI 1-minute averaged magnetic-field data in GSE coordinates
    from Jan 2003. These data have been linearly interpolated to the
    even minute from 16 second data."

OBJECT                                = CONTACT
  ROLE                                = "Principal Investigator"
  PERSON_ID                          = "nfness@bartol.udel.edu"
END_OBJECT                            = CONTACT

OBJECT                                = CONTACT
  ROLE                                = "Co-Investigator"
  PERSON_ID                          = "Charles.Smith@unh.edu"
END_OBJECT                            = CONTACT

OBJECT                                = CONTACT
  ROLE                                = "Data Producer"
  PERSON_ID                          = "jweygand@igpp.ucla.edu"
END_OBJECT                            = CONTACT

END_OBJECT                            = RESOURCE_HEADER

OBJECT                                = TEMPORAL_DESCRIPTION
  CADENCE                            = 00:01:00
  OBJECT                              = SPAN
    START_DATE                      = 1997-01-01T00:00
    STOP_DATE                       = 2004-01-31T23:59
  END_OBJECT                          = SPAN
END_OBJECT                            = TEMPORAL_DESCRIPTION

OBJECT                                = INSTRUMENT_REGION
  HELIOSPHERE                        = "Near L1"
END_OBJECT                            = INSTRUMENT_REGION

OBJECT                                = OBSERVED_REGION
  HELIOSPHERE                        = "Near L1"

```

```

END_OBJECT                                = OBSERVED_REGION

OBJECT                                    = PHYSICAL_PARAMETER
  NAME                                   = SAMPLE_TIME_UTC
  SUPPORT                                = TEMPORAL
  DESCRIPTION                            = "
    Sample UTC in the form DD MM YYYY hh mm ss where
    DD   = day of month (01-31)
    MM   = month of year (01-12)
    YYYY = Gregorian Year AD
    hh   = hour of day   (00:23)
    mm   = minute of hour (00-59)
    ss   = second of minute (00-60)."
END_OBJECT                                = PHYSICAL_PARAMETER

OBJECT                                    = PHYSICAL_PARAMETER
  NAME                                   = MAGNETIC_FIELD_VECTOR
  UNITS                                  = "nT"
  COORDINATE_SYSTEM                     = "GSE"
  DESCRIPTION                            = "
    Magnetic field vector in GSE Coordinates (Bx, By, Bz)."

OBJECT                                    = MEASURED

  OBJECT                                = FIELD
    FIELD_PHYSICAL_QUANTITY             = "Magnetic"
    FIELD_PHYSICAL_QUALIFIER             = "Resampled"
  END_OBJECT                            = FIELD

END_OBJECT                                = MEASURED

END_OBJECT                                = PHYSICAL_PARAMETER

OBJECT                                    = PHYSICAL_PARAMETER
  NAME                                   = SPACECRAFT_POSITION_VECTOR
  SUPPORT                                = "POSITIONAL"
  COORDINATE_SYSTEM                     = "GSE"
  UNITS                                  = "EARTH RADII"
  UNITS_CONVERSION                       = "6378.16 km"
  DESCRIPTION                            = "
    ACE spacecraft location in GSE coordinates (X,Y,Z)."
END_OBJECT                                = PHYSICAL_PARAMETER

OBJECT                                    = ACCESS_INFORMATION
  ACCESS_URL                             =
"http://www.igpp.ucla.edu/getdata?mission=ACE&instrument=mag&format=text"
  FORMAT                                 = "text"
  ENCODING                               = "gzip"
END_OBJECT                                = ACCESS_INFORMATION

OBJECT                                    = ACCESS_INFORMATION
  ACCESS_URL                             =
"http://www.igpp.ucla.edu/getdata?mission=ACE&instrument=mag&format=matlab"
  FORMAT                                 = "matlab"
  ENCODING                               = "none"
END_OBJECT                                = ACCESS_INFORMATION

END_OBJECT                                = NUMERICAL_DATA


END

```

## 7. Definitions of the Data Model Terms

### How to Read a Definition

Each element has certain attributes and context for use. The details for each element are presented in the following form:



The value associated with an element must be one of the following:

**Attribute** : Extra information regarding the element on which it appears. It is tightly coupled to the element.

**Container** : An element that is a container of other elements. If "Container" is specified the element must have sub-elements specified. When a container element is used no value is assigned to the element. All values are contained within the sub-elements.

**Numeric** : An element that has a value which is a real number expressed in base 10.

**Date** : An element that has a value which is a date. A date is given in the ISO 8601 recommended primary standard notation: YYYY-MM-DD, where YYYY is the year in the Gregorian calendar, MM is the month of the year between 01 (January) and 12 (December), and DD is the day of the month between 01 and 31. It may also have an optional time portion (See Time). The time portion must follow the date portion with both portions separated by a "T". For example, "2004-07-29" is July 29, 2004 and "2004-07-29T12:30:00" is precisely 12:30 on July 29, 2004.

**Enumeration** : An element that has a value selected from a list of values. The list to use is indicated in the definition. For example, "Enumerates - see Project List" indicates only values found in the "Project List" may be assigned to this element.

**Count** : An element that has a value which is a base 10 integer number.

**Item** : An element which indicates a state or existence of an attribute. An item is valueless. An item may not contain other elements, but could have attributes. An item may also be a member of an enumerated list.

**Text** : An element that has a value which is a sequence of characters. The number of characters may be limited and is indicated in the definition. A text may have a formation rule. If so this is indicated in the definition.

**URL** : An element or attribute that has a value of a Uniform Resource Locator, the global address of documents and other resources on the World Wide Web. The first part of the address indicates what protocol to use, and the second part specifies the IP address or the domain name where the resource is located, the third part is the path within the domain of the resource.

<b>Access Information</b>	<b>Container</b>
Attributes of the resource which pertain to how to acquire the resource, availability and storage format.	
Sub-elements:	
Access URL	
Acknowledgement	
Availability	
Caveats	
Encoding	
Format	
Repository Name	
<b>Access Rights</b>	<b>Enumeration</b>
Permissions granted or denied by the host of a product to allow other users to access and use the resource.	
Allowed Values:	
Open	
Restricted	
<b>Access URL</b>	<b>Text</b>
Attributes of the method of acquiring a resource including a URL, name and description.	
Sub-elements:	
Description	
Name	
URL	
<b>Acknowledgement</b>	<b>Text</b>
The individual, group or organization which should be acknowledged when the data is used in or contributes to a presentation or publication.	
<b>Acronym</b>	<b>Text</b>
An alternate or shortened name used to refer to a resource.	
<b>Activity Index</b>	<b>Item</b>
An indication, derived from one or more measurements, of the level of activity of an object or region, such as sunspot number, F10.7 flux, Dst, or the Polar Cap Indices.	
<b>Address</b>	<b>Text</b>
Directions for finding some location; written on letters or packages that are to be delivered to that location.	
<b>Aerosol</b>	<b>Item</b>
A suspension of fine solid or liquid particles in gas.	
<b>Alfven Mach Number</b>	<b>Item</b>
The ratio of the bulk flow speed to the Alfven speed.	
<b>Alias</b>	<b>Text</b>
An alternate or shortened name used to refer to a resource.	
<b>Alpha Particle</b>	<b>Item</b>
A positively charged nuclear particle that consists of two protons and two neutrons.	
<b>Antenna</b>	<b>Item</b>
A sensor used to measure electric potential.	
<b>ASCII</b>	<b>Item</b>
A sequence of characters that adheres to American Standard Code for Information Interchange (ASCII) which is an 7-bit character-coding scheme.	



<b>Aurora</b>	<b>Item</b>
A high latitude atmospheric optical phenomenon caused by electrically-charged particles bombarding the upper atmosphere of a planet in the presence of a magnetic field.	
<b>Availability</b>	<b>Enumeration</b>
An indication of the method or service which may be used to access the resource.	
Sub-elements:	
Offline	
Online	
Allowed Values:	
Offline	
Online	
<b>Average</b>	<b>Item</b>
The statistical mean; the sum of a set of values divided by the number of values in the set.	
<b>AVI</b>	<b>Item</b>
Audio Video Interleave (AVI) a digital format for movies that conforms to the Microsoft Windows Resource Interchange File Format (RIFF).	
<b>Azimuthal Angle</b>	<b>Item</b>
The horizontal angular distance from a reference direction. For a vector (X,Y,Z) in any coordinate system, the azimuthal angle is $\arctan(Y/X)$ .	
<b>Binary</b>	<b>Item</b>
A direct representation of the bits which may be stored in memory on a computer	
<b>Bow Shock Crossing</b>	<b>Item</b>
A crossing of the boundary between the undisturbed (except for foreshock effects) solar wind and the shocked, decelerated solar wind of the magnetosheath.	
<b>BZIP2</b>	<b>Item</b>
An open standard algorithm by Julian Seward using Burrows-Wheeler block sorting and Huffman coding. See < <a href="http://www.bzip.org/">http://www.bzip.org/</a> >	
<b>Cadence</b>	<b>Time</b>
The time interval between the start of successive measurements.	
<b>Calibrated</b>	<b>Value</b>
Data wherein sensor outputs have been convolved with instrument response function, often irreversibly, to yield physical parameter values.	
<b>Cartesian</b>	<b>Item</b>
A coordinate system in which the position of a point is determined by its distance from two or three mutually perpendicular axes.	
<b>Catalog</b>	<b>Container</b>
A tabular listing of events or observational notes, especially those that have utility in aiding a user in locating data. Catalogues include lists of events, files in a product, and data availability.	
Sub-elements:	
Access Information	
Input Resource ID	
Instrument ID	
Keyword	
Observatory ID	
Phenomenon Type	
Resource Header	
Resource ID	
Timespan	

<b>Caveats</b>	<b>Text</b>
Information which may be important in the avoidance of misuse of the resource. This includes things such as instrument maladies, corruption or contamination.	
<b>CDF</b>	<b>Item</b>
Common Data Format (CDF). A binary storage format developed at Goddard Space Flight Center (GSFC).	
<b>CEF</b>	<b>Item</b>
Cluster Exchange Format (CEF) is a self-documenting ASCII format designed for the exchange of data. There are two versions of CEF which are not totally compatible.	
<b>CEF 1</b>	<b>Item</b>
Cluster Exchange Format (CEF), version 1, is a self-documenting ASCII format designed for the exchange of data. The metadata contains information compatible with the ISTP recommendations for CDF.	
<b>CEF 2</b>	<b>Item</b>
Cluster Exchange Format (CEF), version 2, is a self-documenting ASCII format designed for the exchange of data and introduced for Cluster Active Archive. Compared to version 1, the metadata description of vectors and tensors is different.	
<b>CGM</b>	<b>Item</b>
Corrected Geomagnetic - A coordinate system from a spatial point with GEO radial distance and geomagnetic latitude and longitude, follow the epoch-appropriate IGRF/DGRF model field vector through to the point where the field line crosses the geomagnetic dipole equatorial plane. Then trace the dipole magnetic field vector Earthward from that point on the equatorial plane, in the same hemisphere as the original point, until the initial radial distance is reached. Designate the dipole latitude and longitude at that point as the CGM latitude and longitude of the original point. See < <a href="http://nssdc.gsfc.nasa.gov/space/cgm/cgmm_des.html">http://nssdc.gsfc.nasa.gov/space/cgm/cgmm_des.html</a> >	
<b>Channeltron</b>	<b>Item</b>
An instrument that detects electrons, ions, and UV-radiation, according to the principle of a secondary emission multiplier. It is typically used in electron spectroscopy and mass spectrometry.	
<b>Charged Particle Flux</b>	<b>Item</b>
Measurements of fluxes of charged or ionized particles at above thermal energies, including relativistic particles of solar and galactic origin. May give simple fluxes, but more complete distributions are sometimes possible. Composition measurements may also be made.	
<b>Chromosphere</b>	<b>Item</b>
The region of the Sun's (or a star's) atmosphere above the temperature minimum and below the Transition Region. The solar chromosphere is approximately 400 km to 2100 km above the photosphere, and characterized by temperatures from 4500 - 28000 K.	
<b>Circular</b>	<b>Item</b>
Relative to polarization, right-hand circularly polarized light is defined such that the electric field is rotating clockwise as seen by an observer towards whom the wave is moving. Left-hand circularly polarized light is defined such that the electric field is rotating counterclockwise as seen by an observer towards whom the wave is moving. The polarization of magnetohydrodynamic waves is specified with respect to the ambient mean magnetic field : right-hand polarized waves have a transverse electric field component which turns in a right-handed sense (that of the gyrating electrons) around the magnetic field.	
<b>Co-Investigator</b>	<b>Item</b>
An individual who is a scientific peer and major participant for an investigation.	
<b>Component</b>	<b>Item</b>
A part of a multi-part entity, e.g., the components of a vector.	

<b>Contact</b>	<b>Container</b>
The person or organization who may be able to provide special assistance or serve as a channel for communication for additional information about a resource.	
Sub-elements:	
Person ID	
Role	
<b>Coordinate Representation</b>	<b>Enumeration</b>
The method or form for specifying a given point in a given coordinate system	
Allowed Values:	
Cartesian	
Cylindrical	
Spherical	
<b>Coordinate System</b>	<b>Container</b>
Specification of the origin and orientation of axes against which the location of some point is given and the representative form of each point.	
Sub-elements:	
Coordinate Representation	
Coordinate System Name	
<b>Coordinate System Name</b>	<b>Enumeration</b>
Identifies the coordinate system in which the position, direction or observation has been expressed.	
Allowed Values:	
CGM	
DM	
GEI	
GEO	
GSE	
GSEQ	
GSM	
HAE	
HEE	
HEEQ	
HG	
HGI	
LGM	
MAG	
MFA	
RTN	
SC	
SE	
SM	
SR	
SR2	
SSE	
<b>Corona</b>	<b>Item</b>
The outermost atmospheric region of the Sun or a star, characterized by ionization temperatures above $10^5$ K. The solar corona starts at about 2100 km above the photosphere; there is no generally defined upper limit.	
<b>Coronal Mass Ejection</b>	<b>Item</b>
A solar event which involves a burst of plasma which is ejected from the Sun into the interplanetary medium.	
<b>Counts</b>	<b>Item</b>
An enumeration of the number of detection events occurring in a particle detector.	
<b>Cross Spectrum</b>	<b>Item</b>
The Fourier transform of the cross correlation of two physical or empirical observations.	

	Cylindrical	Item
	A system of curvilinear coordinates in which the position of a point in space is determined by its perpendicular distance from a given line, its distance from a selected reference plane perpendicular to this line, and its angular distance from a selected reference line when projected onto this plane.	
	Data Producer	Item
	An individual who generated the resource and is familiar with its provenance.	
	Dayside	Item
	Sunward of a dawn-dusk meridian, either on the surface of, or above, some reference body.	
	Description	Text
	A detailed description of the resource which should include discussions of the main quantities in the resource, possible uses and search terms. A description should also include whether any corrections (i.e, geometry, inertial) have been applied to it.	
	Deviation	Item
	The difference between an observed value and the expected value of a quantity.	
	Dimension	Container
	Attributes of an independent variable or axis associated with the data.	
	Sub-elements:	
Item	Description	
	Size	
	Display Cadence	Time
	The time interval between the successive display elements.	
	Display Data	Container
	A graphical representation of data wherein the underlying numeric values are not (readily) accessible for ana6q 0.. Examplues are lineplotes and xpecro grmsa.	

A dipole antenna of which the active (sensor) elements are small spheres located at the ends of two wires deployed in the equatorial plane, on opposite sides of a spinning spacecraft.

Dust	Item
------	------

Free microscopic particles of solid material.

Dynamic Spectra	Item
-----------------	------

A three-dimensional representation of successive spectra which allows time evolution to be clearly seen. Time is plotted along the abscissa, frequency (or particle energy) along the ordinate, and the spectral power density (or differential particle flux) is represented by different shades of grey, or color. This representation is also known as a spectrogram.

Electric	Item
----------	------

The physical attribute that exerts an electrical force.

Electric Field	Item
----------------	------

Measurements of electric field vectors (sometimes not all components) as a time series.

Electron	Item
----------	------

An elementary particle consisting of a charge of negative electricity equal to about  $1.602 \times 10^{19}$  Coulomb and having a mass when at rest of about  $9.109534 \times 10^{-28}$  gram.

Electron Drift Instrument	Item
---------------------------	------

An active experiment to measure the electron drift velocity based on sensing the displacement of a weak beam of electrons after one gyration in the ambient magnetic field.

Electrostatic Analyser	Item
------------------------	------

An instrument which uses charged plates to analyze the mass, charge and kinetic energies of charged particles which enter the instrument.

Email	Text
-------	------

The electronic address at which the individual may be contacted expressed in the form "local-part@domain".

Emissivity	Item
------------	------

The ratio of radiant energy from a material to that from a blackbody at the same kinetic temperature

Encoding	Enumeration
----------	-------------

A set of unambiguous rules that establishes the representation of information within a file.

Allowed Values:

ASCII

GZIP

None

Unicode

ZIP

End Date	Date
----------	------

Electrostatic Analyser: An instrument which uses charged plates to analyze the mass, charge and kinetic energies of charged particles which enter the instrument.

An enhancement of interplanetary fluxes of energetic ions accelerated by interplanetary shocks and/or solar flares.

<b>Energy</b>	<b>Item</b>
The capacity for doing work as measured by the capability of doing work (potential energy) or the conversion of this capability to motion (kinetic energy) Attributes: Units	
<b>Equivalent Width</b>	<b>Item</b>
The area of the spectral line profile divided by the peak height or depth.	
<b>Exposure</b>	<b>Time</b>
The time interval over which an individual measurement is taken.	
<b>Faraday Cup</b>	<b>Item</b>
An instrument consisting of an electrode from which electrical current is measured while a charged particle beam (electrons or ions) impinges on it. Used to determine energy spectrum and sometimes ion composition of the impinging particles.	
<b>Field</b>	<b>Container</b>
The space around a radiating body within which its electromagnetic attributes can exert force on another similar body that is not in direct contact. Sub-elements: Field Physical Quantity Field Qualifier	
<b>Field Physical Quantity</b>	<b>Enumeration</b>
The physical attribute of the field. Allowed Values: Cross Spectrum Electric Magnetic Potential Poynting Flux	
<b>Field Qualifier</b>	<b>Enumeration</b>
Characterizes the directional and statistical aspects of the field observation. Allowed Values: Average Component Deviation Magnitude Peak Perpendicular Variance Vector	
<b>Fit</b>	<b>Item</b>
Values that make an model agree with the data.	
<b>FITS</b>	<b>Item</b>
Flexible Image Transport System (FITS) is a digital format primarily designed to store scientific data sets consisting of multi-dimensional arrays (1-D spectra, 2-D images or 3-D data cubes) and 2-dimensional tables containing rows and columns of data.	
<b>Flux</b>	<b>Item</b>
In radiation studies, this refers to the amount of radiant energy passing through a unit area Attributes: Units	
<b>Flux Feedback</b>	<b>Item</b>
A search coil whose bandwidth and signal/noise ratio are increased by the application of	

negative feedback at the sensor (flux) level by driving a collocated coil with a signal from the preamplifier.

<b>Forbush Decrease</b>	<b>Item</b>
-------------------------	-------------

A rapid decrease in the observed galactic cosmic ray intensity following the passage of an outwardly convecting interplanetary magnetic field disturbance, such as those associated with large CME's, that sweep some galactic cosmic rays away from Earth.

<b>Format</b>	<b>Enumeration</b>
---------------	--------------------

The organization of data according to preset specifications. The value is selected from a list of accepted names for known, well documented formats.

Allowed Values:

AVI  
Binary  
CDF  
CEF  
CEF 1  
CEF 2  
FITS  
GIF  
HDF  
HDF 4  
HDF 5  
HTML  
IDFS  
IDL  
JPEG  
MPEG  
NetCDF  
PDF  
PNG  
QuickTime  
Text  
TIFF  
UDF  
XML

<b>Fourier Transform Spectrograph</b>	<b>Item</b>
---------------------------------------	-------------

An instrument that determines the spectra of a radiative source, using time-domain measurements and a Fourier transform.

<b>Frequency</b>	<b>Item</b>
------------------	-------------

(1) The number of occurrences within a given time period. (2) Vibrations per second of the photon field; may be given as a single number, multiple numbers, or as ranges.

<b>Gamma Rays</b>	<b>Item</b>
-------------------	-------------

Photons with a wavelength range: 0.00001 to 0.001 nm

<b>GEI</b>	<b>Item</b>
------------	-------------

Geocentric Equatorial Inertial - A coordinate system where the Z axis is along Earth's spin vector, positive northward. X axis points towards the first point of Aries (from the Earth towards the Sun at the vernal equinox). See Russell, 1971

<b>General Contact</b>	<b>Item</b>
------------------------	-------------

An individual who can provide information on a range of subjects or who can direct you to a domain expert.

<b>GEO</b>	<b>Item</b>
------------	-------------

Geographic - geocentric corotating - A coordinate system where the Z axis is along Earth's spin vector, positive northward. X axis lies in Greenwich meridian, positive towards Greenwich. See Russell, 1971.

<b>Geomagnetic Storm</b>	<b>Item</b>
A magnetospheric disturbance typically defined by variations in the horizontal component of the Earth's surface magnetic field. The variation typically starts with a field enhancement associated with a solar wind pressure pulse and continues with a field depression associated with an enhancement of the diamagnetic magnetospheric ring current.	
<b>GIF</b>	<b>Item</b>
Graphic Interchange Format (GIF) first introduced in 1987 by CompuServe. GIF uses LZW compression and images are limited to 256 colours.	
<b>GSE</b>	<b>Item</b>
Geocentric Solar Ecliptic - A coordinate system where the X axis is from Earth to Sun. Z axis is normal to the ecliptic, positive northward. See Russell, 1971.	
<b>GSEQ</b>	<b>Item</b>
Geocentric Solar Equatorial - A coordinate system where the X axis is from Earth to Sun. Y axis is parallel to solar equatorial plane. Z axis is positive northward. See Russell, 1971	
<b>GSM</b>	<b>Item</b>
Geocentric Solar Magnetospheric - A coordinate system where the X axis is from Earth to Sun, Z axis is northward in a plane containing the X axis and the geomagnetic dipole axis. See Russell, 1971	
<b>GZIP</b>	<b>Item</b>
An open standard algorithm distributed by GHU based on LZ77 and Huffman coding. See < <a href="http://www.gnu.org/software/gzip/gzip.html">http://www.gnu.org/software/gzip/gzip.html</a> > or < <a href="http://www.gzip.org/">http://www.gzip.org/</a> >	
<b>HAE</b>	<b>Item</b>
Heliocentric Aries Ecliptic - A coordinate system where the Z axis is normal to the ecliptic plane, positive northward. X axis is positive towards the first point of Aries (from Earth to Sun at vernal equinox). Same as SE below. See Hapgood, 1992.	
<b>Hard X-rays</b>	<b>Item</b>
Photons with a wavelength range: 0.001 to 0.1 nm	
<b>HDF</b>	<b>Item</b>
Hierarchical Data Format	
<b>HDF 4</b>	<b>Item</b>
Hierarchical Data Format, Version 4	
<b>HDF 5</b>	<b>Item</b>
Hierarchical Data Format, Version 5	
<b>Heat Flux</b>	<b>Item</b>
Flow of thermal energy through a gas or plasma; typically computed as third moment of a distribution function.	
<b>HEE</b>	<b>Item</b>
Heliocentric Earth Ecliptic - A coordinate system where the Z axis is normal to the ecliptic plane, positive northward. X axis points from Sun to Earth. See Hapgood, 1992	
<b>HEEQ</b>	<b>Item</b>
Heliocentric Earth Equatorial - A coordinate system where the Z axis is normal to the solar equatorial plane, positive northward. X axis is generally Earthward in the plane defined by the Z axis and the Sun-Earth direction. See Hapgood, 1992.	
<b>Heliosphere</b>	<b>Enumeration</b>
The solar atmosphere extending roughly from the outer corona to the edge of the solar plasma at the heliopause separating primarily solar plasma from interstellar plasma.	



## Allowed Values:

Inner  
Near 1AU  
Near L1  
Outer

HF radar	Item
An instrument which uses high frequency (HF) radar to obtain an image of an object.	
HG	Item
Heliographic - A heliocentric rotating coordinate system where the Z axis is normal to the solar equatorial plane, positive northward. X, Y axes rotate with a 25.38 day period. The zero longitude (X axis) is defined as the longitude that passed through the ascending node of the solar equator on the ecliptic plane on 1 January, 1854 at 12 UT. See < <a href="http://nssdc.gsfc.nasa.gov/space/helios/coord_des.html">http://nssdc.gsfc.nasa.gov/space/helios/coord_des.html</a> >	
HGI	Item
Heliographic Inertial - A heliocentric coordinate system where the Z axis is normal to the solar equatorial plane, positive northward. X axis is along the intersection line between solar equatorial and ecliptic planes. The X axis was positive at SE longitude of 74.367 deg on Jan 1, 1900. (See SE below.) See < <a href="http://nssdc.gsfc.nasa.gov/space/helios/coord_des.html">http://nssdc.gsfc.nasa.gov/space/helios/coord_des.html</a> >	
High Latitude	Item
The region located poleward of 60 degrees of latitude.	
HTML	Item
A text file containing structured information represented in the HyperText Mark-up Language (HTML). See < <a href="http://www.w3.org/MarkUp/">http://www.w3.org/MarkUp/</a> >	
IDFS	Item
Instrument Data File Set (IDFS) is a set of files written in a prescribed format which contain data, timing data, and meta-data. IDFS was developed at Southwest Research Institute (SwRI).	
IDL	Item
Interactive Data Language (IDL) save set.	
Image Intensity	Item
Measurements of the two-dimensional distribution of the intensity of photons from some region or object such as the Sun or the polar auroral regions; can be in any wavelength band, and polarized, etc.	
Imager	Item
An instrument which samples the radiation from an area at one or more spectral ranges emitted or reflected by an object.	
Information URL	Container
Attributes of the method of acquiring additional information.	
Sub-elements:	
Description	
Name	
URL	
Infrared	Item
Photons with a wavelength range: 760 to 1.00x10 <sup>6</sup> nm	
Inner	Item
The region of the heliosphere extending radially out from the "surface" of the Sun to 1 AU.	
Input Resource ID	Text
The resource identifier for a resource which was used to generate this resource.	

Instrument	Container
A device which is used to sense and parameterize a physical phenomenon.	
Sub-elements:	
Acronym	
Instrument Name	
Instrument Type	
Investigation Name	
Resource Header	
Resource ID	
Instrument ID	Text
The identifier of an Instrument resource.	
Instrument Name	Text
A programmatically assigned name for a set of one or more interrelated instruments selected as a unit to operate on an observatory.	
Instrument Region	Container
The portion of space occupied by the instrument at the time of an observation. A region is distinguished by certain natural features or physical characteristics.	
Sub-elements:	
Heliosphere	
Magnetosphere	
Near Surface	
Sun	
Surface	
Instrument Type	Enumeration
The technique of observation : the type of instrument used to make the measurement, including the sensor and other hardware or software features as appropriate.	
Allowed Values:	
Antenna	
Channeltron	
Double Sphere	
Electron Drift Instrument	
Electrostatic Analyser	
Energetic Particle Instrument	
Faraday Cup	
Flux Feedback	
Fourier Transform Spectrograph	
HF Radar	
Imager	
Langmuir Probe	
Long Wire	
Magnetometer	
Mass Spectrometer	
Microchannel Plate	
Monopole	
Particle Correlator	
Quadrispherical Analyser	
Resonance Sounder	
Search Coil	
Spacecraft Potential Control	
Spectral Power Receiver	
Spectrograph	
Spectrometer	
Waveform Receiver	
Intensity	Item
The amount of energy transmitted by electromagnetic radiation, for example, the number of photons arriving in a given time.	
Attributes: Units	

<b>Interior</b>	<b>Item</b>
The region inside the body which is not visible from outside the body.	
<b>Interplanetary Shock</b>	<b>Item</b>
A shock propagating generally antisunward through the slower solar wind, often seen in front of CME-associated plasma clouds.	
<b>Investigation Name</b>	<b>Text</b>
The name given to a related set of observation which may be obtained from more than one observatory.	
<b>Ion</b>	<b>Item</b>
An atom that has acquired a net electric charge by gaining or losing one or more electrons.(Note: $Z > 2$ )	
<b>Ion Composition</b>	<b>Item</b>
In situ measurements of the relative flux or density of electrically charged particles in the space environment. May give simple fluxes, but full distribution functions are sometimes measured.	
<b>Irradiance</b>	<b>Item</b>
Measurements of time series of the integrated intensity of photons in some wavelength band from some region or object, typically the Sun.	
<b>JPEG</b>	<b>Item</b>
A binary format for still images defined by the Joint Photographic Experts Group	
<b>Keyword</b>	<b>Text</b>
A word or phrase that is relevant to the resource but does not exist in other documentary information.	
<b>Langmuir Probe</b>	<b>Item</b>
A monopole antenna associated with an instrument. The instrument applies a potential to the antenna which is swept to determine the voltage/current characteristic. This provides information about the plasma surrounding the probe and spacecraft.	
<b>LGM</b>	<b>Item</b>
Local Geomagnetic - A coordinate system used mainly for Earth surface or near Earth surface magnetic field data. X axis northward from observation point in a geographic meridian. Z axis downward towards Earth's center. In this system, H (total horizontal component) = $\text{SQRT}(B_x^2 + B_y^2)$ and D (declination angle) = $\arctan(B_y/B_x)$	
<b>Line Depth</b>	<b>Item</b>
In spectra, a measure of the amount of absorption for a particular wavelength or frequency in the spectrum	
<b>Line of Sight</b>	<b>Item</b>
The line of sight is the line that connects the observer with the observed object. This expression is often used with measurements of Doppler velocity and magnetic field in magnetograms, where only the component of the vector field directed along the line of sight is measured.	
<b>Linear</b>	<b>Item</b>
Relative to polarization, confinement of the E-field vector to a given plane	
<b>Long Wire</b>	<b>Item</b>
A dipole antenna whose active (sensor) elements are two wires deployed in the equatorial plane on opposite sides of a spinning spacecraft, and whose length is several times greater than the spacecraft diameter.	
<b>Low Latitude</b>	<b>Item</b>
The region located at or anti-poleward of 60 degrees of latitude.	

<b>MAG</b>	Item
Geomagnetic - geocentric. Z axis is parallel to the geomagnetic dipole axis, positive north. X is in the plane defined by the Z axis and the Earth's rotation axis. If N is a unit vector from the Earth's center to the north geographic pole, the signs of the X and Y axes are given by $Y = N \times Z$ , $X = Y \times Z$ . See Russell, 1971, and < <a href="http://cdpp.cnes.fr/00428.pdf">http://cdpp.cnes.fr/00428.pdf</a> >	
<b>Magnetic</b>	Item
The physical attribute attributed to a magnet or its equivalent.	
<b>Magnetic Field</b>	Item
Measurements of magnetic field vectors (sometimes not all components) as time series; can be space- or ground-based. Also, [Zeeman splitting, etc. based]: A region of space near a magnetized body where magnetic forces can be detected [as measured by methods such as Zeeman splitting, etc.]	
<b>Magnetogram</b>	Item
Measurements of the vector or line-of-sight magnetic field determined from remote sensing measurements of the detailed structure of spectral lines, including their splitting and polarization. ("Magnetogram.")	
<b>Magnetometer</b>	Item
An instrument which measures the ambient magnetic field.	
<b>Magnetopause Crossing</b>	Item
A crossing of the interface between the shocked solar wind in the magnetosheath and the magnetic field and plasma in the magnetosphere.	
<b>Magnetosheath</b>	Item
The region between the bow shock and the magnetopause, characterized by very turbulent plasma.	
<b>Magnetosphere</b>	Enumeration
The region of space above the atmosphere or surface of the planet, and bounded by the magnetopause, that is under the direct influence of the planet's magnetic field.	
Allowed Values:	
Magnetosheath	
Magnetotail	
Main	
Polar	
<b>Magnetotail</b>	Item
The region on the night side of the body where the magnetic field is stretched backwards by the force of the solar wind. For Earth, the magnetotail begins at a night-side radial distance of 10 $R_E$ ( $X > -10R_E$ ).	
<b>Magnitude</b>	Item
A measure of the strength or size of a vector quantity.	
<b>Main</b>	Item
The region of the magnetosphere below 60 degrees in magnetic latitude and between the body's nearby region and the magnetopause. For Earth, the main magnetosphere is defined as the volume bounded by part of a sphere at 2000 km altitude, part of the magnetopause, the latitude = 60 deg surface and the $X = -10 R_E$ plane.	
<b>Mass</b>	Item
The property of a body that is a measure of its inertia, that is commonly taken as a measure of the amount of material it contains, that causes it to have weight in a gravitational field, and that along with length and time constitutes one of the fundamental quantities on which all physical measurements are based.	
<b>Mass Density</b>	Item

The mass of particles per unit volume.

Attributes: Units

<b>Mass Spectrometer</b>	<b>Item</b>
An instrument which distinguishes chemical species in terms of their different isotopic masses.	
<b>Measured</b>	<b>Container</b>
Attributes of observations obtained from an instrument or sensor.	
Sub-elements:	
Field	
Mixed	
Particle	
Photon	
<b>Measurement Type</b>	<b>Enumeration</b>
A high level characterization of the content of the data product.	
Allowed Values:	
Activity Index	
Charged Particle Flux	
Dopplergram	
Dynamic Spectra	
Electric Field	
Energetic Particles	
Image Intensity	
Ion Composition	
Irradiance	
Magnetic Field	
Magnetogram	
Neutral Atom Images	
Neutral Gas	
Profile	
Radiance	
Radio and Plasma Waves	
Radio Soundings	
Thermal Plasma	
<b>MFA</b>	<b>Item</b>
Magnetic Field Aligned - A coordinate system spacecraft-centered system with Z in the direction of the ambient magnetic field vector. X is in the plane defined by Z and the spacecraft-Sun line, positive sunward. See < <a href="http://cdpp.cnes.fr/00428.pdf">http://cdpp.cnes.fr/00428.pdf</a> >	
<b>Microchannel Plate</b>	<b>Item</b>
An instrument used for the detection of elementary particles, ions, ultraviolet rays and soft X-rays constructed from very thin conductive glass capillaries.	
<b>Microwave</b>	<b>Item</b>
Photons with a wavelength range: $1.00 \times 10^6$ to $1.50 \times 10^7$ nm	
<b>Mixed</b>	<b>Item</b>
A measured observation which is derived from a combination of two or more individual measurements.	
<b>Mode Amplitude</b>	<b>Item</b>
In helioseismology the magnitude of oscillation of waves of a particular geometry.	
<b>Molecule</b>	<b>Item</b>
A group of atoms so united and combined by chemical affinity that they form a complete, integrated whole, being the smallest portion of any particular compound that can exist in a free state	
<b>Moment</b>	<b>Item</b>

Parameters determined by integration over a distribution function convolved with a power of velocity.

<b>MPEG</b>	<b>Item</b>
A digital format for movies defined by the Motion Picture Experts Group	
<b>N</b>	<b>Count</b>
An integer that approximates the mass of an isotope and that designates the number of nucleons in the nucleus. Also called the mass number. Often equated with the atomic mass.	
<b>Name</b>	<b>Text</b>
A language unit by which a person or thing is known.	
<b>Near 1AU</b>	<b>Item</b>
A heliospheric region approximately as far from the Sun as is the Earth, but at a significant angular separation from the Earth-Sun line (i.e., not Near L1).	
<b>Near L1</b>	<b>Item</b>
A heliospheric region near the Earth that includes the location of L1 spacecraft as well as of spacecraft orbiting around or near the Earth that make excursions outside the bow shock. (The bow shock is where the solar wind undergoes a shock transition due to interaction with the Earth's magnetosphere.)	
<b>Near Surface</b>	<b>Enumeration</b>
The gaseous and possibly ionized environment of a body extending from the surface to some specified altitude. For the Earth, this altitude is 2000 km. Allowed Values: High Latitude Low Latitude	
<b>NetCDF</b>	<b>Item</b>
Unidata Program Center's Network Common Data Form (NetCDF). A self-describing data portable data format for array-oriented data access. See <a href="http://my.unidata.ucar.edu/content/software/netcdf">http://my.unidata.ucar.edu/content/software/netcdf</a>	
<b>Neutral</b>	<b>Item</b>
Either a particle, an object, or a system that has a net electric charge of zero	
<b>Neutral Atom Images</b>	<b>Item</b>
Measurements of neutral atom fluxes as a function of look direction; often related to remote energetic charged particles that lose their charge through charge-exchange and then reach the detector on a line.	
<b>Neutral Gas</b>	<b>Item</b>
Measurements of neutral atomic and molecular components of a body and its surrounding environments.	
<b>Nightside</b>	<b>Item</b>
Anti-Sunward of a dawn-dusk meridian, either on the surface of, or above, some reference body.	
<b>None</b>	<b>Item</b>
A lack or absence of anything.	
<b>Number Density</b>	<b>Item</b>
The number of particles per unit volume. Attributes: Units	
<b>Numerical Data</b>	<b>Container</b>
Data stored as numerical values in a specified format.	

## Sub-elements:

Access Information  
 Access Rights  
 Input Resource ID  
 Instrument ID  
 Instrument Region  
 Keyword  
 Measurement Type  
 Observatory ID  
 Observed Region  
 Physical Parameter  
 Provider Processing Level  
 Provider Version  
 Resource Header  
 Resource ID  
 Spectral Range  
 Temporal Description

<b>Observatory</b>	Container
The host (spacecraft, network, facility) for instruments making observations.	
Sub-elements:	
Observatory Group Observatory Name Resource Header Resource ID	
<b>Observatory Group</b>	Text
A set of programmatically related observatories. The value is taken from an approved list of observatory group names.	
<b>Observatory ID</b>	Text
The identifier of an Observatory resource.	
<b>Observatory Name</b>	Text
The language unit which identifies a location or platform. An observatory may be part of an observatory group. The value is taken from an approved list of observatory names.	
<b>Observed Region</b>	Container
The portion of space measured by the instrument at the time of an observation. A region is distinguished by certain natural features or physical characteristics. It is the location of the observatory for in situ data, the location or region sensed by remote sensing observatories and the location-of-relevance for parameters that are derived from observational data.	
Sub-elements:	
Heliosphere Magnetosphere Near Surface Sun Surface	
<b>Offline</b>	Text
Not directly accessible electronically. This includes resources which may to be moved to an online status in response to a given request.	
Sub-elements:	
Medium	
<b>Online</b>	Item
Directly accessible electronically.	
<b>Open</b>	Item
Access is granted to everyone.	
<b>Optical</b>	Item

Photons with a wavelength range: 380 to 760 nm

<b>Organization Name</b>	<b>Text</b>
A unit within a company or other entity (e.g., Government agency or branch of service) within which many projects are managed as a whole.	
<b>Other</b>	<b>Item</b>
Values, such as flags, that are not time tags, location data or measured or derived parameters.	
<b>Outer</b>	<b>Item</b>
The region of the heliosphere from 1 AU to the farthest extent of the heliosphere (heliopause).	
<b>Parallel</b>	<b>Item</b>
Having the same direction as a given direction	
<b>Particle</b>	<b>Container</b>
A description of the types of particles observed in the measurement. This includes both direct observations and inferred observations.	
Sub-elements:	
Particle Physical Quantity	
Particle Qualifier	
Particle Type	
<b>Particle Correlator</b>	<b>Item</b>
An instrument which correlates particle flux to help identify wave/particle interactions.	
<b>Particle Physical Quantity</b>	<b>Enumeration</b>
A characterization of the physical properties of the particle.	
Allowed Values:	
Alfven Mach Number	
Coordinate System	
Counts	
Differential Directional Parti	
Heat Flux	
Mass Density	
Number Density	
Phase-Space Density	
Plasma Beta	
Pressure	
Sonic Mach Number	
Temperature	
Thermal Speed	
Velocity	
<b>Particle Qualifier</b>	<b>Enumeration</b>
Characterizes the directional and statistical aspects of the particle observation.	
Allowed Values:	
Average	
Component	
Deviation	
Fit	
Magnitude	
Moment	
Parallel	
Peak	
Perpendicular	
Variance	
Vector	
<b>Particle Type</b>	<b>Enumeration</b>
A characterization of the kind of particle observed by the measurement.	



## Allowed Values:

Aerosol  
 Alpha Particle  
 Dust  
 Electron  
 Ion  
 Mass  
 Molecule  
 N  
 Neutral  
 Proton  
 Q  
 Size  
 Z

<b>PDF</b>	<u>Item</u>
A document expressed in the Portable Document Format (PDF) as defined by Adobe.	
<b>Peak</b>	<u>Item</u>
The maximum value for the quantity in question, over a period of time which is usually equal to the cadence.	
<b>Perpendicular</b>	<u>Item</u>
At right angles to a given direction.	
<b>Person</b>	<u>Container</u>
An individual human being.	
Sub-elements:	
Address Email Organization Name Person Name Phone Number Resource ID	
<b>Person ID</b>	<u>Text</u>
The identifier assigned to a Person description.	
<b>Person Name</b>	<u>Text</u>
The words used to address an individual.	
<b>Phase-Space Density</b>	<u>Item</u>
The number of particles per unit volume in the six-dimensional space of position and velocity.	
<b>Phenomenon Type</b>	<u>Enumeration</u>
The characteristics or categorization of an event type.	
Allowed Values:	
Aurora Bow Shock Crossing Coronal Mass Ejection Energetic Solar Particle Event Forbush Decrease Geomagnetic Storm Interplanetary Shock Magnetopause Crossing Solar Flare Solar Wind Extreme	
<b>Phone Number</b>	<u>Text</u>
The symbols and numerals required to contact an individual by telephone. The string may contain punctuation marks such as dash (-) or dot (.) to separate fields within the string.	

<b>Photon</b>	<b>Container</b>
Photon (radio through gamma-rays): the fundamental particle or quantum of electromagnetic radiation (radiant energy)	
Sub-elements:	
Photon Physical Quantity	
Photon Qualifier	
<b>Photon Physical Quantity</b>	<b>Enumeration</b>
A characterization of the physical properties of the photon.	
Allowed Values:	
Emissivity	
Equivalent Width	
Flux	
Intensity	
Line Depth	
Magnetic Field	
Mode Amplitude	
Photon Independent Variable	
Polarization	
Stoke's Parameters	
Velocity	
<b>Photon Qualifier</b>	<b>Enumeration</b>
Characterizes the directional and statistical aspects of the photon observation.	
Allowed Values:	
Average	
Circular	
Line of Sight	
Linear	
Peak	
Stoke's Parameter	
Variance	
Vector	
<b>Photosphere</b>	<b>Item</b>
The atmospheric layer of the Sun or a star from which continuum radiation, especially optical, is emitted to space. For the Sun, the photosphere is about 500 km thick.	
<b>Physical Parameter</b>	<b>Container</b>
A container of information regarding a parameter whose values are part of the product. Every product contains or can be related to one or more parameters.	
Sub-elements:	
Cadence	
Caveats	
Coordinate System	
Description	
Dimension	
Measured	
Name	
Support	
Units	
Units Conversion	
<b>Plasma Beta</b>	<b>Item</b>
The ratio of the plasma pressure to the magnetic pressure.	
<b>PNG</b>	<b>Item</b>
A digital format for still images. Portable Network Graphics (PNG)	
<b>Polar Angle</b>	<b>Item</b>
The angle between the Z axis and the given vector direction.	

<b>Polarization</b>	<b>Item</b>
Direction of the electric vector of an electromagnetic wave. The wave can be linearly polarized in any direction perpendicular to the direction of travel, circularly polarized (clockwise or counterclockwise), unpolarized, or mixtures of the above.	
<b>Positional</b>	<b>Item</b>
The specification of the location of an object or measurement within a reference coordinate system. The position is usually expressed as a set of values corresponding to the location along a set of orthogonal axes together with the date/time of the observation.	
<b>Potential</b>	<b>Item</b>
A field which obeys Laplace's Equation.	
<b>Poynting Flux</b>	<b>Item</b>
The rate of energy transport per unit area per steradian.	
<b>Pressure</b>	<b>Item</b>
A characterization of the random energy related to the velocity and distribution of particles.	
<b>Principal Investigator</b>	<b>Item</b>
An individual who is the administrative and scientific lead for an investigation.	
<b>Profile</b>	<b>Item</b>
Measurements of a quantity as a function of height above an object such as the limb of a body.	
<b>Proton</b>	<b>Item</b>
An elementary particle that is a constituent of all atomic nuclei, that carries a positive charge numerically equal to the charge of an electron, and that has a mass of $1.673 \times 10^{-24}$ gram.	
<b>Provider ID</b>	<b>Item</b>
The identifier for a Contact resource for the person or organization who provided the resource.	
<b>Provider Processing Level</b>	<b>Text</b>
The provider specific classification of the processing performed on the product.	
<b>Provider Release Date</b>	<b>Date</b>
The date the product was made available by the provider. The Provider Release Date is relevant only to the product life-cycle of the provider.	
<b>Provider Resource Name</b>	<b>Text</b>
A short textual description of a resource used by the provider which may be used to identify a resource.	
<b>Provider Version</b>	<b>Text</b>
Describes the release or edition of the product used by the provider. The formation rule may vary between providers. It is intended to aid in queries to the provider regarding the product.	
<b>Q</b>	<b>Numeric</b>
The charge state of an ion, generally expressed in multiples of the coulomb charge of an electron.	
<b>Quadrispherical Analyser</b>	<b>Item</b>
An instrument used for the 3-D detection of plasma, energetic electrons and ions, and for positive-ion composition measurements.	
<b>QuickTime</b>	<b>Item</b>
A format for digital movies, as defined by Apple Computer. See <a href="http://developer.apple.com/quicktime/">http://developer.apple.com/quicktime/</a>	
<b>Radiance</b>	<b>Item</b>

A measure of the amount of electromagnetic radiation leaving a point on the surface in a specified time period in a unit area from a given direction.

<b>Radio and Plasma Waves</b>	<b>Item</b>
Measurements of electric and/or magnetic fields using electric or magnetic antennas at frequencies anywhere between the spacecraft spin frequency and the characteristic frequencies of the ambient plasma. The output can be waveform, power spectral density, or other statistical parameters.	
<b>Radio Frequency</b>	<b>Item</b>
Photons with a wavelength range: 100,000 to 1.00x10 <sup>11</sup> nm	
<b>Radio Soundings</b>	<b>Item</b>
Measurements of plasma density, magnetic field and possibly other parameters of the space environment by active probing of the plasma by radio waves.	
<b>Repository Name</b>	<b>Text</b>
The name of the location or facility where the product is stored. The repository name is selected from a list of names for established repositories.	
Allowed Values:	
For a current list see Identifiers for the location or facility where the product is stored. The repository name is selected from a list of established repositories. See < <a href="http://www.igpp.ucla.edu/spase/">http://www.igpp.ucla.edu/spase/</a> > for the list.	
<b>Resonance Sounder</b>	<b>Item</b>
A combination of a radio receiver and a pulsed transmitter used to study the plasma surrounding a spacecraft by identifying resonances or cut-offs (of the wave dispersion relation), whose frequencies are related to the ambient plasma density and magnetic field. When the transmitter is off it is essentially a high frequency-resolution spectral power receiver.	
<b>Resource Header</b>	<b>Container</b>
Attributes of a resource which pertain to the provider of the resource and descriptive information about the resource.	
Sub-elements:	
Acknowledgement	
Alias	
Contact	
Description	
Information URL	
Provider Resource Name	
Resource Name	
<b>Resource ID</b>	<b>Text</b>
The unique identifier of an item. The resource ID must be unique within the context it is used.	
<b>Resource Name</b>	<b>Text</b>
A short textual description of a resource which may be useful when read by a person.	
<b>Restricted</b>	<b>Item</b>
Access to the product is regulated and requires some form of identification.	
<b>Role</b>	<b>Enumeration</b>
The assigned or assumed function or position of an individual.	
Allowed Values:	
Co-Investigator	
Data Producer	
General Contact	
Principal Investigator	
Scientist	
Team Leader	
Team Member	

## Technical Contact

<b>RTF</b>	<b>Item</b>
Rich Text Format (RTF). Structured information as defined by Microsoft.	
<b>RTN</b>	<b>Item</b>
Radial Tangential Normal. Typically centered at a spacecraft. Used for IMF and plasma V vectors. R (radial) axis is radially away from the Sun, T (tangential) axis is normal to the plane formed by R and the Sun's spin vector, positive in the direction of planetary motion. N (normal) is $R \times T$ .	
<b>SC</b>	<b>Item</b>
Spacecraft - A coordinate system defined by the spacecraft geometry and/or spin. Often has Z axis parallel to spacecraft spin vector. X and Y axes may or may not corotate with the spacecraft. See SR and SR2 below.	
<b>Scientist</b>	<b>Item</b>
An individual who is an expert in the phenomenon and related physics represented by the resource.	
<b>SE</b>	<b>Item</b>
Solar Ecliptic - A heliocentric coordinate system where the Z axis is normal to the ecliptic plane, positive northward. X axis is positive towards the first point of Aries (from Earth to Sun at vernal equinox). Same as HAE above. See < <a href="http://nssdc.gsfc.nasa.gov/space/helios/coor_des.html">http://nssdc.gsfc.nasa.gov/space/helios/coor_des.html</a> >	
<b>Search coil</b>	<b>Item</b>
A loop of wire used to determine the time variation of the magnetic flux threading the loop by measurement of the electric potential difference induced between the ends of the wire.	
<b>SGI</b>	<b>Item</b>
Binary data compatible with Silicon Graphic platforms.	
<b>Size</b>	<b>Count</b>
The physical dimensions, proportions, magnitude, or extent of an object..	
<b>SM</b>	<b>Item</b>
Solar Magnetic - A geocentric coordinate system where the Z axis is northward along Earth's dipole axis, X axis is in plane of z axis and Earth-Sun line, positive sunward. See Russell, 1971.	
<b>Soft X-rays</b>	<b>Item</b>
Range: $0.1 \leq x < 10$ nm; Conventional abbreviation: XUV	
<b>Solar Flare</b>	<b>Item</b>
An explosive event in the Sun's atmosphere which produces electromagnetic radiation across the electromagnetic spectrum at multiple wavelengths from long-wave radio to the shortest wavelength gamma rays.	
<b>Solar Wind Extreme</b>	<b>Item</b>
Intervals of unusually large or small values of solar wind attributes such as flow speed and ion density.	
<b>Sonic Mach Number</b>	<b>Item</b>
The ratio of the bulk flow speed to the speed of sound in the medium.	
<b>Spacecraft Potential Control</b>	<b>Item</b>
An instrument to control the electric potential of a spacecraft with respect to the ambient plasma by emitting a variable current of positive ions.	
<b>SPASE</b>	<b>Container</b>

Space Physics Archive Search and Extract (SPASE). The outermost container or envelope for SPASE metadata. This indicates the start of the SPASE metadata.

Sub-elements:

- Catalog
- Display Data
- Instrument
- Numerical Data
- Observatory
- Person
- Version

<b>Spatial Range</b>	<b>Item</b>
----------------------	-------------

A description, in an appropriate coordinate system, of the positions of the elements of an image; may be done using a reference and relative positions, or with bins giving the description of a 2-D grid.

<b>Spectral Power Receiver</b>	<b>Item</b>
--------------------------------	-------------

A radio receiver which determines the power spectral density of the electric or magnetic field, or both, at one or more frequencies.

<b>Spectral Range Name</b>	<b>Enumeration</b>
----------------------------	--------------------

The general term used to describe wavelengths or frequencies within a given span of values for those quantities.

Allowed Values:

- Gamma Rays
- Hard X-rays
- Infrared
- Microwave
- Optical
- Radio Frequency
- Ultraviolet
- X-Rays

<b>Spectrograph</b>	<b>Item</b>
---------------------	-------------

The image (intensity vs. wavelength) of the electromagnetic spectrum produced by a spectrometer.

<b>Spectrometer</b>	<b>Item</b>
---------------------	-------------

An instrument that measures the component wavelengths of light or other electromagnetic radiation into its component wavelengths.

<b>Spectrum</b>	<b>Item</b>
-----------------	-------------

Measurements of the intensity of radiation as a function of frequency or wavelength.

<b>Spherical</b>	<b>Item</b>
------------------	-------------

A system of curvilinear coordinates characterized by an azimuthal angle (longitude), a polar angle (latitude), and a distance (radius) from a point to the origin.

<b>SR</b>	<b>Item</b>
-----------	-------------

Spin Reference - A special case of a Spacecraft (SC) coordinate system for a spinning spacecraft. Z is parallel to the spacecraft spin vector. X and Y rotate with the spacecraft. See <<http://cdpp.cnes.fr/00428.pdf>>

<b>SR2</b>	<b>Item</b>
------------	-------------

Spin Reference 2 - A special case of a Spacecraft (SC) coordinate system for a spinning spacecraft. Z is parallel to the spacecraft spin vector. X is in the plane defined by Z and the spacecraft-Sun line, positive sunward. See <<http://cdpp.cnes.fr/00428.pdf>>

<b>SSE</b>	<b>Item</b>
------------	-------------

Spacecraft Solar Ecliptic - A coordinate system used for deep space spacecraft, for example Helios. - X axis from spacecraft to Sun. Z axis normal to ecliptic plane, positive northward.

Note: Angle between normals to ecliptic and to Helios orbit plane ~ 0.25 deg.

<b>Start Date</b>	<b>Date</b>
The specification of a starting point in time.	
<b>Stoke's Parameters</b>	<b>Item</b>
The four coordinates (usually called I, Q, U, and V) relative to a particular basis for the representation of the polarization state of an electromagnetic wave propagating through space.	
<b>Sun</b>	<b>Enumeration</b>
The star upon which our solar system is centered.	
Allowed Values:	
Chromosphere	
Corona	
Interior	
Photosphere	
Transition Region	
<b>Support</b>	<b>Enumeration</b>
Information useful in understanding the context of an observation, typically observed or measured coincidentally with a physical observation.	
Allowed Values:	
Other	
Positional	
Temporal	
<b>Surface</b>	<b>Item</b>
The outermost area of a solid object.	
<b>Team Leader</b>	<b>Item</b>
An individual who is the scientific and administrative lead for an investigation.	
<b>Team Member</b>	<b>Item</b>
An individual who is a major participant in an investigation.	
<b>Technical Contact</b>	<b>Item</b>
An individual who can provide specific information with regard to the resource or supporting software	
<b>Temperature</b>	<b>Item</b>
A measure of the kinetic energy of random motion with respect to the average. Temperature is properly defined only for an equilibrium particle distribution (Maxwellian distribution).	
<b>Temporal</b>	<b>Item</b>
Pertaining to time.	
<b>Temporal Description</b>	<b>Container</b>
A characterization of the time over which the measurement was taken.	
Sub-elements:	
Cadence	
Exposure	
Timespan	
<b>TeX</b>	<b>Item</b>
A document expressed in the typesetting language TeX originally defined by Donald Knuth.	
<b>Text</b>	<b>Item</b>
ASCII text	
<b>Thermal Plasma</b>	<b>Item</b>

Measurements of the plasma in the energy regime where the most of the plasma occurs. May be the basic fluxes in the form of distribution functions or the derived bulk parameters (density, flow velocity, etc.).

<b>Thermal Speed</b>	<b>Item</b>
----------------------	-------------

For a Maxwellian distribution, the difference between the mean speed and the speed within which ~69% (one sigma) of all the members of the speed distribution occur.

<b>TIFF</b>	<b>Item</b>
-------------	-------------

A binary format for still pictures. Tagged Image Format File (TIFF). Originally developed by Aldus and now controlled by Adobe.

<b>Timespan</b>	<b>Container</b>
-----------------	------------------

The duration of an interval in time.

Sub-elements:

End Date

Start Date

<b>Transition Region</b>	<b>Item</b>
--------------------------	-------------

A very narrow (<100 km) layer between the chromosphere and the corona where the temperature rises abruptly from about 8000 to about 500,000 K.

<b>UDF</b>	<b>Item</b>
------------	-------------

Universal Data Format (UDF). The Optical Technology Storage Association's Universal Disk Format, based on ISO 13346. See <<http://www.osta.org/specs/index.htm>>

<b>Ultraviolet</b>	<b>Item</b>
--------------------	-------------

Photons with a wavelength range: 10 to 400 nm

<b>Uncalibrated</b>	<b>Item</b>
---------------------	-------------

Duplicate data are removed from the data stream and data are time ordered. Values are not adjusted for any potential biases or external factors.

<b>Unicode</b>	<b>Item</b>
----------------	-------------

Text in multi-byte Unicode format.

<b>Units</b>	<b>Text</b>
--------------	-------------

A description of the standardized measurement increments in which a value is specified. The description is represented as a mathematical phrase. Individual units within the phrase must conform to the International System of Units (SI) which is maintained by BIPM (Bureau International des Poids et Mesures. See <<http://www.bipm.fr/>> ). The symbol associated with a unit should be used in the phrase. Within the phrase the circumflex (^) is used to indicate a power, a star (\*) is used to indicate multiplication and a slash (/) division. When symbols are not separated by a mathematical operator, multiplication is assumed. Symbols for base units can be found at: <[http://www.bipm.fr/en/si/si\\_brochure/chapter2/2-1/#symbols](http://www.bipm.fr/en/si/si_brochure/chapter2/2-1/#symbols)> and those for common derived units can be found at: <[http://www.bipm.fr/en/si/derived\\_units/2-2-2.html](http://www.bipm.fr/en/si/derived_units/2-2-2.html)>

<b>Units Conversion</b>	<b>Text</b>
-------------------------	-------------

The multiplicative factor for converting a unit into International System of Units (SI) units. The factor is expressed in the form "number > x", where "number" is a numerical value and "x" is the appropriate SI units. The basic SI units are Enumerated: m (meter), N (newton), kg (kilogram), Pa (pascal), s (second), Hz (hertz), A (ampere), V (volt), K (kelvin), W (watt), rad (radian), J (joule), sr (steradian), C (coulomb), T (tesla), ohm (ohm), mho (mho or seimens), H (henry), and F (farad). Two useful units which are not SI units are: degree (angle), and unitless (no units). An example is: "1.0E-5>T" which converts the units, presumable nT, to tesla. Another example is: "1.0e-1>km/s" which converts a velocity expressed in meters per second to kilometers per second.

<b>URL</b>	<b>Text</b>
------------	-------------

Uniform Resource Locator (URL) is the global address of documents and other resources on the World Wide Web. The first part of the address indicates what protocol to use, and the second



part specifies the IP address or the domain name where the resource is located followed by the pathname of the resource. A URL is specified in the form protocol://server.domain.name:port/pathname. Example protocols are HTTP or FTP, server domain name is the Internet name.

Variance	Item
A measure of dispersion of a set of data points around their mean value. The expectation value of the squared deviations from the mean.	
Vector	Item
A quantity having both magnitude and direction, e.g. displacement, velocity, acceleration and force.	
Velocity	Item
Rate of change of position. Also used for the average velocity of a collection of particles, also referred to as "bulk velocity".	
Version	Text
Indicates the release identifier. When used to indicate the release of the SPASE data model, it is a in the form Major.Minor.Fix where Major: A significant change in the architecture of the model or rewrite of the implementation. This includes major changes in design or implementation language. This number starts at 0 (zero). Minor: An addition of terms or features that require changes in documentation/external API. This number starts at 0 (zero). Fix: Any change that doesn't require documentation/external API changes. This number starts at 0 (zero).	
Waveform Receiver	Item
A radio receiver which outputs the value of one or more components of the electric and/or magnetic field as a function of time.	
Wavelength	Item
The distance between successive points of equal amplitude and phase on a wave (for example, crest to crest or trough to trough)	
Wavenumber	Item
A quantity that is inversely proportional to the wavelength of a wave.	
X-Rays	Item
Photons with a wavelength range: $0.001 \leq x < 10$ nm	
XDR	Item
Binary data in the eXternal Data Representation (XDR) format. See RFC 1014 < <a href="http://www.faqs.org/rfcs/rfc1014.html">http://www.faqs.org/rfcs/rfc1014.html</a> >	
XML	Item
eXtensible Mark-up Language (XML). A structured format for representing information. See < <a href="http://www.w3.org/XML/">http://www.w3.org/XML/</a> >	
Z	Count
The number of protons in an atomic nucleus. Also called the atomic number.	
ZIP	Item
An open standard for compression which is a variation of the LZW method and was originally used in the PKZIP utility.	

## 8. Enumeration of Selected Quantities

Lists are either "open" or "closed". The items in a "closed" list are determined by the SPASE model and definitions of each item is in the SPASE data dictionary. The items in an "open" list are determined by an external control authority. The URL for the control authority is indicated in the definition of each "open" list.

Access Rights List	Identifier
--------------------	------------

Term	Definition
Open	Access is granted to everyone.
Restricted	Access to the product is regulated and requires some form of identification.

Availability List	Identifier
-------------------	------------

Term	Definition
Offline	Not directly accessible electronically. This includes resources which may to be moved to an online status in response to a given request.
Online	Directly accessible electronically.

Coordinate Representation List	Closed
--------------------------------	--------

Term	Definition
Cartesian	A coordinate system in which the position of a point is determined by its distance from two or three mutually perpendicular axes.
Cylindrical	A system of curvilinear coordinates in which the position of a point in space is determined by its perpendicular distance from a given line, its distance from a selected reference plane perpendicular to this line, and its angular distance from a selected reference line when projected onto this plane.
Spherical	A system of curvilinear coordinates characterized by an azimuthal angle (longitude), a polar angle (latitude), and a distance (radius) from a point to the origin.

Coordinate System Name List	Closed
-----------------------------	--------

Identifiers for coordinate systems in which the position, direction or observation has been expressed.

Term	Definition
CGM	Corrected Geomagnetic - A coordinate system from a spatial point with GEO radial distance and geomagnetic latitude and longitude, follow the epoch-appropriate IGRF/DGRF model field vector through to the point where the field line crosses the geomagnetic dipole equatorial plane. Then trace the dipole magnetic field

	vector Earthward from that point on the equatorial plane, in the same hemisphere as the original point, until the initial radial distance is reached. Designate the dipole latitude and longitude at that point as the CGM latitude and longitude of the original point. See < <a href="http://nssdc.gsfc.nasa.gov/space/cgm/cgmm_des.html">http://nssdc.gsfc.nasa.gov/space/cgm/cgmm_des.html</a> >
DM	Dipole Meridian - A coordinate system centered at the observation point. Z axis is parallel to the Earth's dipole axis, positive northward. X is in the plane defined by Z and the line linking the observation point with the Earth's center. Y is positive eastward. See < <a href="http://cdpp.cnes.fr/00428.pdf">http://cdpp.cnes.fr/00428.pdf</a> >
GEI	Geocentric Equatorial Inertial - A coordinate system where the Z axis is along Earth's spin vector, positive northward. X axis points towards the first point of Aries (from the Earth towards the Sun at the vernal equinox). See Russell, 1971
GEO	Geographic - geocentric corotating - A coordinate system where the Z axis is along Earth's spin vector, positive northward. X axis lies in Greenwich meridian, positive towards Greenwich. See Russell, 1971.
GSE	Geocentric Solar Ecliptic - A coordinate system where the X axis is from Earth to Sun. Z axis is normal to the ecliptic, positive northward. See Russell, 1971.
GSEQ	Geocentric Solar Equatorial - A coordinate system where the X axis is from Earth to Sun. Y axis is parallel to solar equatorial plane. Z axis is positive northward. See Russell, 1971
GSM	Geocentric Solar Magnetospheric - A coordinate system where the X axis is from Earth to Sun, Z axis is northward in a plane containing the X axis and the geomagnetic dipole axis. See Russell, 1971
HAE	Heliocentric Aries Ecliptic - A coordinate system where the Z axis is normal to the ecliptic plane, positive northward. X axis is positive towards the first point of Aries (from Earth to Sun at vernal equinox). Same as SE below. See Hapgood, 1992.
HEE	Heliocentric Earth Ecliptic - A coordinate system where the Z axis is normal to the ecliptic plane, positive northward. X axis points from Sun to Earth. See Hapgood, 1992
HEEQ	Heliocentric Earth Equatorial - A coordinate system where the Z axis is normal to the solar equatorial plane, positive northward. X axis is generally Earthward in the plane defined by the Z axis and the Sun-Earth direction. See Hapgood, 1992.
HG	Heliographic - A heliocentric rotating coordinate system where the Z axis is normal to the solar equatorial plane, positive northward. X, Y axes rotate with a 25.38 day period. The zero longitude (X axis) is defined as the longitude that passed through the ascending node of the

	<p>solar equator on the ecliptic plane on 1 January, 1854 at 12 UT. See</p> <p>&lt;<a href="http://nssdc.gsfc.nasa.gov/space/helios/coord_des.html">http://nssdc.gsfc.nasa.gov/space/helios/coord_des.html</a>&gt;</p>
HGI	<p>Heliographic Inertial - A heliocentric coordinate system where the Z axis is normal to the solar equatorial plane, positive northward. X axis is along the intersection line between solar equatorial and ecliptic planes. The X axis was positive at SE longitude of 74.367 deg on Jan 1, 1900. (See SE below.) See</p> <p>&lt;<a href="http://nssdc.gsfc.nasa.gov/space/helios/coord_des.html">http://nssdc.gsfc.nasa.gov/space/helios/coord_des.html</a>&gt;</p>
LGM	<p>Local Geomagnetic - A coordinate system used mainly for Earth surface or near Earth surface magnetic field data. X axis northward from observation point in a geographic meridian. Z axis downward towards Earth's center. In this system, H (total horizontal component) = <math>\text{SQRT}(B_x^2 + B_y^2)</math> and D (declination angle) = <math>\arctan(B_y/B_x)</math></p>
MAG	<p>Geomagnetic - geocentric. Z axis is parallel to the geomagnetic dipole axis, positive north. X is in the plane defined by the Z axis and the Earth's rotation axis. If N is a unit vector from the Earth's center to the north geographic pole, the signs of the X and Y axes are given by <math>Y = N \times Z</math>, <math>X = Y \times Z</math>. See Russell, 1971, and</p> <p>&lt;<a href="http://cdpp.cnes.fr/00428.pdf">http://cdpp.cnes.fr/00428.pdf</a>&gt;</p>
MFA	<p>Magnetic Field Aligned - A coordinate system spacecraft-centered system with Z in the direction of the ambient magnetic field vector. X is in the plane defined by Z and the spacecraft-Sun line, positive sunward. See</p> <p>&lt;<a href="http://cdpp.cnes.fr/00428.pdf">http://cdpp.cnes.fr/00428.pdf</a>&gt;</p>
RTN	<p>Radial Tangential Normal. Typically centered at a spacecraft. Used for IMF and plasma V vectors. R (radial) axis is radially away from the Sun, T (tangential) axis is normal to the plane formed by R and the Sun's spin vector, positive in the direction of planetary motion. N (normal) is <math>R \times T</math>.</p>
SC	<p>Spacecraft - A coordinate system defined by the spacecraft geometry and/or spin. Often has Z axis parallel to spacecraft spin vector. X and Y axes may or may not corotate with the spacecraft. See SR and SR2 below.</p>
SE	<p>Solar Ecliptic - A heliocentric coordinate system where the Z axis is normal to the ecliptic plane, positive northward. X axis is positive towards the first point of Aries (from Earth to Sun at vernal equinox). Same as HAE above. See</p> <p>&lt;<a href="http://nssdc.gsfc.nasa.gov/space/helios/coord_des.html">http://nssdc.gsfc.nasa.gov/space/helios/coord_des.html</a>&gt;</p>
SM	<p>Solar Magnetic - A geocentric coordinate system where the Z axis is northward along Earth's dipole axis, X axis is in plane of z axis and Earth-Sun line, positive sunward. See Russell, 1971.</p>
SR	<p>Spin Reference - A special case of a Spacecraft (SC) coordinate system for a spinning spacecraft. Z is parallel</p>

SR2	to the spacecraft spin vector. X and Y rotate with the spacecraft. See < <a href="http://cdpp.cnes.fr/00428.pdf">http://cdpp.cnes.fr/00428.pdf</a> > Spin Reference 2 - A special case of a Spacecraft (SC) coordinate system for a spinning spacecraft. Z is parallel to the spacecraft spin vector. X is in the plane defined by Z and the spacecraft-Sun line, positive sunward. See < <a href="http://cdpp.cnes.fr/00428.pdf">http://cdpp.cnes.fr/00428.pdf</a> >
SSE	Spacecraft Solar Ecliptic - A coordinate system used for deep space spacecraft, for example Helios. - X axis from spacecraft to Sun. Z axis normal to ecliptic plane, positive northward. Note: Angle between normals to ecliptic and to Helios orbit plane ~ 0.25 deg.

**Encoding List**

Closed

Identifier for unambiguous rules that establishes the representation of information within a file.

Term	Definition
ASCII	A sequence of characters that adheres to American Standard Code for Information Interchange (ASCII) which is an 7-bit character-coding scheme.
BZIP2	An open standard algorithm by Julian Seward using Burrows-Wheeler block sorting and Huffman coding. See < <a href="http://www.bzip.org/">http://www.bzip.org/</a> >
GZIP	An open standard algorithm distributed by GHU based on LZ77 and Huffman coding. See < <a href="http://www.gnu.org/software/gzip/gzip.html">http://www.gnu.org/software/gzip/gzip.html</a> > or < <a href="http://www.gzip.org/">http://www.gzip.org/</a> >
None	A lack or absence of anything.
Unicode	Text in multi-byte Unicode format.
ZIP	An open standard for compression which is a variation of the LZW method and was originally used in the PKZIP utility.

**Field Physical Quantity List**

Closed

Identifiers for the physical attribute of the field.

Term	Definition
Cross Spectrum	The Fourier transform of the cross correlation of two physical or empirical observations.
Electric	The physical attribute that exerts an electrical force.
Magnetic	The physical attribute attributed to a magnet or its equivalent.
Potential	A field which obeys Laplace's Equation.
Poynting Flux	The rate of energy transport per unit area per steradian.

**Field Qualifier List**

Closed

Identifiers for terms which can be associated with a Field Quantity.

Term	Definition
Average	The statistical mean; the sum of a set of values divided by the number of values in the set.
Component	A part of a multi-part entity, e.g., the components of a

	vector.
Deviation	The difference between an observed value and the expected value of a quantity.
Magnitude	A measure of the strength or size of a vector quantity.
Peak	The maximum value for the quantity in question, over a period of time which is usually equal to the cadence.
Perpendicular	At right angles to a given direction.
Variance	A measure of dispersion of a set of data points around their mean value. The expectation value of the squared deviations from the mean.
Vector	A quantity having both magnitude and direction, e.g. displacement, velocity, acceleration and force.

**Format List**Closed

Identifiers for data organized according to preset specifications.

Term	Definition
AVI	Audio Video Interleave (AVI) a digital format for movies that conforms to the Microsoft Windows Resource Interchange File Format (RIFF).
Binary	A direct representation of the bits which may be stored in memory on a computer
CDF	Common Data Format (CDF). A binary storage format developed at Goddard Space Flight Center (GSFC).
CEF	Cluster Exchange Format (CEF) is a self-documenting ASCII format designed for the exchange of data. There are two versions of CEF which are not totally compatible.
CEF 1	Cluster Exchange Format (CEF), version 1, is a self-documenting ASCII format designed for the exchange of data. The metadata contains information compatible with the ISTP recommendations for CDF.
CEF 2	Cluster Exchange Format (CEF), version 2, is a self-documenting ASCII format designed for the exchange of data and introduced for Cluster Active Archive. Compared to version 1, the metadata description of vectors and tensors is different.
FITS	Flexible Image Transport System (FITS) is a digital format primarily designed to store scientific data sets consisting of multi-dimensional arrays (1-D spectra, 2-D images or 3-D data cubes) and 2-dimensional tables containing rows and columns of data.
GIF	Graphic Interchange Format (GIF) first introduced in 1987 by CompuServe. GIF uses LZW compression and images are limited to 256 colours.
HDF	Hierarchical Data Format
HDF 4	Hierarchical Data Format, Version 4
HDF 5	Hierarchical Data Format, Version 5
HTML	A text file containing structured information represented in the HyperText Mark-up Language (HTML). See <a href="http://www.w3.org/MarkUp/">http://www.w3.org/MarkUp/</a>

IDFS	Instrument Data File Set (IDFS) is a set of files written in a prescribed format which contain data, timing data, and meta-data. IDFS was developed at Southwest Research Institute (SwRI).
IDL	Interactive Data Language (IDL) save set.
JPEG	A binary format for still images defined by the Joint Photographic Experts Group
MPEG	A digital format for movies defined by the Motion Picture Experts Group
NetCDF	Unidata Program Center's Network Common Data Form (NetCDF). A self-describing data portable data format for array-oriented data access. See <a href="http://my.unidata.ucar.edu/content/software/netcdf/">http://my.unidata.ucar.edu/content/software/netcdf/</a>
PDF	A document expressed in the Portable Document Format (PDF) as defined by Adobe.
PNG	A digital format for still images. Portable Network Graphics (PNG)
QuickTime	A format for digital movies, as defined by Apple Computer. See <a href="http://developer.apple.com/quicktime/">http://developer.apple.com/quicktime/</a>
Text	ASCII text
TIFF	A binary format for still pictures. Tagged Image Format File (TIFF). Originally developed by Aldus and now controlled by Adobe.
UDF	Universal Data Format (UDF). The Optical Technology Storage Association's Universal Disk Format, based on ISO 13346. See <a href="http://www.osta.org/specs/index.htm">http://www.osta.org/specs/index.htm</a>
XML	eXtensible Mark-up Language (XML). A structured format for representing information. See <a href="http://www.w3.org/XML/">http://www.w3.org/XML/</a>

#### Heliosphere List

Closed

Identifiers for regions of the solar atmosphere which extends roughly from the inner corona to the edge of the solar plasma at the heliopause separating primarily solar plasma from interstellar plasma.

Term	Definition
Inner	The region of the heliosphere extending radially out from the "surface" of the Sun to 1 AU.
Near 1AU	A heliospheric region approximately as far from the Sun as is the Earth, but at a significant angular separation from the Earth-Sun line (i.e., not Near L1).
Near L1	A heliospheric region near the Earth that includes the location of L1 spacecraft as well as of spacecraft orbiting around or near the Earth that make excursions outside the bow shock. (The bow shock is where the solar wind undergoes a shock transition due to interaction with the Earth's magnetosphere.)
Outer	The region of the heliosphere from 1 AU to the farthest extent of the heliosphere (heliopause).

#### Instrument Type List

Closed

Identifiers for the type of experiment the instrument performs. This is the technique of

observation.

Term	Definition
Antenna	A sensor used to measure electric potential.
Channeltron	An instrument that detects electrons, ions, and UV-radiation, according to the principle of a secondary emission multiplier. It is typically used in electron spectroscopy and mass spectrometry.
Double Sphere	A dipole antenna of which the active (sensor) elements are small spheres located at the ends of two wires deployed in the equatorial plane, on opposite sides of a spinning spacecraft.
Electron Drift Instrument	An active experiment to measure the electron drift velocity based on sensing the displacement of a weak beam of electrons after one gyration in the ambient magnetic field.
Electrostatic Analyser	An instrument which uses charged plates to analyze the mass, charge and kinetic energies of charged particles which enter the instrument.
Energetic Particle Instrument	An instrument that measures fluxes of charged particles as a function of time, direction of motion, mass, charge and/or species
Faraday Cup	An instrument consisting of an electrode from which electrical current is measured while a charged particle beam (electrons or ions) impinges on it. Used to determine energy spectrum and sometimes ion composition of the impinging particles.
Flux Feedback	A search coil whose bandwidth and signal/noise ratio are increased by the application of negative feedback at the sensor (flux) level by driving a collocated coil with a signal from the preamplifier.
Fourier Transform Spectrograph	An instrument that determines the spectra of a radiative source, using time-domain measurements and a Fourier transform.
HF Radar	An instrument which uses high frequency (HF) radar to obtain an image of an object.
Imager	An instrument which samples the radiation from an area at one or more spectral ranges emitted or reflected by an object.
Langmuir Probe	A monopole antenna associated with an instrument. The instrument applies a potential to the antenna which is swept to determine the voltage/current characteristic. This provides information about the plasma surrounding the probe and spacecraft.
Long Wire	A dipole antenna whose active (sensor) elements are two wires deployed in the equatorial plane on opposite sides of a spinning spacecraft, and whose length is several times greater than the spacecraft diameter.
Magnetometer	An instrument which measures the ambient magnetic field.
Mass Spectrometer	An instrument which distinguishes chemical species in



	terms of their different isotopic masses.
Microchannel Plate	An instrument used for the detection of elementary particles, ions, ultraviolet rays and soft X-rays constructed from very thin conductive glass capillaries.
Particle Correlator	An instrument which correlates particle flux to help identify wave/particle interactions.
Quadrispherical Analyser	An instrument used for the 3-D detection of plasma, energetic electrons and ions, and for positive-ion composition measurements.
Resonance Sounder	A combination of a radio receiver and a pulsed transmitter used to study the plasma surrounding a spacecraft by identifying resonances or cut-offs (of the wave dispersion relation), whose frequencies are related to the ambient plasma density and magnetic field. When the transmitter is off it is essentially a high frequency-resolution spectral power receiver.
Search Coil	A loop of wire used to determine the time variation of the magnetic flux threading the loop by measurement of the electric potential difference induced between the ends of the wire.
Spacecraft Potential Control	An instrument to control the electric potential of a spacecraft with respect to the ambient plasma by emitting a variable current of positive ions.
Spectral Power Receiver	A radio receiver which determines the power spectral density of the electric or magnetic field, or both, at one or more frequencies.
Spectrograph	The image (intensity vs. wavelength) of the electromagnetic spectrum produced by a spectrometer.
Spectrometer	An instrument that measures the component wavelengths of light or other electromagnetic radiation into its component wavelengths.
Waveform Receiver	A radio receiver which outputs the value of one or more components of the electric and/or magnetic field as a function of time.

## Magnetosphere List

Closed

Term	Definition
Magnetosheath	The region between the bow shock and the magnetopause, characterized by very turbulent plasma.
Magnetotail	The region on the night side of the body where the magnetic field is stretched backwards by the force of the solar wind. For Earth, the magnetotail begins at a night-side radial distance of 10 Re ( $X > -10R_E$ ).
Main	The region of the magnetosphere below 60 degrees in magnetic latitude and between the body's nearby region and the magnetopause. For Earth, the main magnetosphere is defined as the volume bounded by part of a sphere at 2000 km altitude, part of the magnetopause, the latitude = 60 deg surface and the $X =$

-10 Re plane.

## Measurement Type List

Closed

Identifiers for the method of making an estimated value of a quantity that forms the basis of an observaton.

Term	Definition
Activity Index	An indication, derived from one or more measurements, of the level of activity of an object or region, such as sunspot number, F10.7 flux, Dst, or the Polar Cap Indices.
Charged Particle Flux	Measurements of fluxes of charged or ionized particles at above thermal energies, including relativistic particles of solar and galactic origin. May give simple fluxes, but more complete distributions are sometimes possible. Composition measurements may also be made.
Dopplergram	A map or image depicting the spatial distribution of line-of-sight velocities of the observed object.
Dynamic Spectra	A three-dimensional representation of successive spectra which allows time evolution to be clearly seen. Time is plotted along the abscissa, frequency (or particle energy) along the ordinate, and the spectral power density (or differential particle flux) is represented by different shades of grey, or color. This representation is also known as a spectrogram.
Electric Field	Measurements of electric field vectors (sometimes not all components) as a time series.
Energetic Particles	Pieces of matter that are moving very fast. Energetic particles include protons, electrons, neutrons, neutrinos, the nuclei of atoms, and other sub-atomic particles.
Image Intensity	Measurements of the two-dimensional distribution of the intensity of photons from some region or object such as the Sun or the polar auroral regions; can be in any wavelength band, and polarized, etc.
Ion Composition	In situ measurements of the relative flux or density of electrically charged particles in the space environment. May give simple fluxes, but full distribution functions are sometimes measured.
Irradiance	Measurements of time series of the integrated intensity of photons in some wavelength band from some region or object, typically the Sun.
Magnetic Field	Measurements of magnetic field vectors (sometimes not all components) as time series; can be space- or ground-based. Also, [Zeeman splitting, etc. based]: A region of space near a magnetized body where magnetic forces can be detected [as measured by methods such as Zeeman splitting, etc.]
Magnetogram	Measurements of the vector or line-of-sight magnetic field determined from remote sensing measurements of the detailed structure of spectral lines, including their splitting and polarization. ("Magnetogram.")

Neutral Atom Images	Measurements of neutral atom fluxes as a function of look direction; often related to remote energetic charged particles that lose their charge through charge-exchange and then reach the detector on a line.
Neutral Gas	Measurements of neutral atomic and molecular components of a body and its surrounding environments.
Profile	Measurements of a quantity as a function of height above an object such as the limb of a body.
Radiance	A measure of the amount of electromagnetic radiation leaving a point on the surface in a specified time period in a unit area from a given direction.
Radio and Plasma Waves	Measurements of electric and/or magnetic fields using electric or magnetic antennas at frequencies anywhere between the spacecraft spin frequency and the characteristic frequencies of the ambient plasma. The output can be waveform, power spectral density, or other statistical parameters.
Radio Soundings	Measurements of plasma density, magnetic field and possibly other parameters of the space environment by active probing of the plasma by radio waves.
Thermal Plasma	Measurements of the plasma in the energy regime where the most of the plasma occurs. May be the basic fluxes in the form of distribution functions or the derived bulk parameters (density, flow velocity, etc.).

## Near Surface List

Closed

Term	Definition
High Latitude	The region located poleward of 60 degrees of latitude.
Low Latitude	The region located at or anti-poleward of 60 degrees of latitude.

## Observatory Group List

Open

For a current list see Identifiers for programmatically related observatories. The value is taken from an approved list of observatory group names. See <http://www.igpp.ucla.edu/spase/> for the list.

## Observatory Name List

Open

For a current list see Identifiers for a location or platform. An observatory may be part of an observatory group. The value is taken from an approved list of observatory names. See <http://www.igpp.ucla.edu/spase/> for the list.

## Particle Physical Quantity List

Closed

Term	Definition
Alfven Mach Number	The ratio of the bulk flow speed to the Alfven speed.
Coordinate System	Specification of the origin and orientation of axes against which the location of some point is given and the

	representative form of each point.
Counts	An enumeration of the number of detection events occurring in a particle detector.
Heat Flux	Flow of thermal energy through a gas or plasma; typically computed as third moment of a distribution function.
Mass Density	The mass of particles per unit volume.
Number Density	The number of particles per unit volume.
Phase-Space Density	The number of particles per unit volume in the six-dimensional space of position and velocity.
Plasma Beta	The ratio of the plasma pressure to the magnetic pressure.
Pressure	A characterization of the random energy related to the velocity and distribution of particles.
Sonic Mach Number	The ratio of the bulk flow speed to the speed of sound in the medium.
Temperature	A measure of the kinetic energy of random motion with respect to the average. Temperature is properly defined only for an equilibrium particle distribution (Maxwellian distribution).
Thermal Speed	For a Maxwellian distribution, the difference between the mean speed and the speed within which ~69% (one sigma) of all the members of the speed distribution occur.
Velocity	Rate of change of position. Also used for the average velocity of a collection of particles, also referred to as "bulk velocity".

## Particle Qualifier List

Closed

Identifiers for terms which can be associated with a Particle Quantity.

Term	Definition
Average	The statistical mean; the sum of a set of values divided by the number of values in the set.
Component	A part of a multi-part entity, e.g., the components of a vector.
Deviation	The difference between an observed value and the expected value of a quantity.
Fit	Values that make an model agree with the data.
Magnitude	A measure of the strength or size of a vector quantity.
Moment	Parameters determined by integration over a distribution function convolved with a power of velocity.
Parallel	Having the same direction as a given direction
Peak	The maximum value for the quantity in question, over a period of time which is usually equal to the cadence.
Perpendicular	At right angles to a given direction.
Variance	A measure of dispersion of a set of data points around their mean value. The expectation value of the squared deviations from the mean.
Vector	A quantity having both magnitude and direction, e.g. displacement, velocity, acceleration and force.

## Particle Type List

Closed

Identifiers for the characterization of the kind of particle observed by the measurement.

Term	Definition
Aerosol	A suspension of fine solid or liquid particles in gas.
Alpha Particle	A positively charged nuclear particle that consists of two protons and two neutrons.
Dust	Free microscopic particles of solid material.
Electron	An elementary particle consisting of a charge of negative electricity equal to about $1.602 \times 10^{*-19}$ Coulomb and having a mass when at rest of about $9.109534 \times 10^{*-28}$ gram.
Ion	An atom that has acquired a net electric charge by gaining or losing one or more electrons.(Note: $Z > 2$ )
Mass	The property of a body that is a measure of its inertia, that is commonly taken as a measure of the amount of material it contains, that causes it to have weight in a gravitational field, and that along with length and time constitutes one of the fundamental quantities on which all physical measurements are based.
Molecule	A group of atoms so united and combined by chemical affinity that they form a complete, integrated whole, being the smallest portion of any particular compound that can exist in a free state
N	An integer that approximates the mass of an isotope and that designates the number of nucleons in the nucleus. Also called the mass number. Often equated with the atomic mass.
Neutral	Either a particle, an object, or a system that has a net electric charge of zero
Proton	An elementary particle that is a constituent of all atomic nuclei, that carries a positive charge numerically equal to the charge of an electron, and that has a mass of $1.673 \times 10^{*-24}$ gram.
Q	The charge state of an ion, generally expressed in multiples of the coulomb charge of an electron.
Size	The physical dimensions, proportions, magnitude, or extent of an object..
Z	The number of protons in an atomic nucleus. Also called the atomic number.

#### Phenomenon Type List

Closed

Identifiers for the characteristics or categorization of an observation. Note: Joe King to provide.

Term	Definition
Aurora	A high latitude atmospheric optical phenomenon caused by electrically-charged particles bombarding the upper atmosphere of a planet in the presence of a magnetic field.
Bow Shock Crossing	A crossing of the boundary between the undisturbed (except for foreshock effects) solar wind and the shocked, decelerated solar wind of the magnetosheath.
Coronal Mass Ejection	A solar event which involves a burst of plasma which is

Energetic Solar Particle Event	ejected from the Sun into the interplanetary medium. An enhancement of interplanetary fluxes of energetic ions accelerated by interplanetary shocks and/or solar flares.
Forbush Decrease	A rapid decrease in the observed galactic cosmic ray intensity following the passage of an outwardly convecting interplanetary magnetic field disturbance, such as those associated with large CME's, that sweep some galactic cosmic rays away from Earth.
Geomagnetic Storm	A magnetospheric disturbance typically defined by variations in the horizontal component of the Earth's surface magnetic field. The variation typically starts with a field enhancement associated with a solar wind pressure pulse and continues with a field depression associated with an enhancement of the diamagnetic magnetospheric ring current.
Interplanetary Shock	A shock propagating generally antisunward through the slower solar wind, often seen in front of CME-associated plasma clouds.
Magnetopause Crossing	A crossing of the interface between the shocked solar wind in the magnetosheath and the magnetic field and plasma in the magnetosphere.
Solar Flare	An explosive event in the Sun's atmosphere which produces electromagnetic radiation across the electromagnetic spectrum at multiple wavelengths from long-wave radio to the shortest wavelength gamma rays.
Solar Wind Extreme	Intervals of unusually large or small values of solar wind attributes such as flow speed and ion density.

## Photon Physical Quantity List

Closed

Identifiers for the characterization of the physical properties of the photon.

Term	Definition
Emissivity	The ratio of radiant energy from a material to that from a blackbody at the same kinetic temperature
Equivalent Width	The area of the spectral line profile divided by the peak height or depth.
Flux	In radiation studies, this refers to the amount of radiant energy passing through a unit area
Intensity	The amount of energy transmitted by electromagnetic radiation, for example, the number of photons arriving in a given time.
Line Depth	In spectra, a measure of the amount of absorption for a particular wavelength or frequency in the spectrum
Magnetic Field	Measurements of magnetic field vectors (sometimes not all components) as time series; can be space- or ground-based. Also, [Zeeman splitting, etc. based]: A region of space near a magnetized body where magnetic forces can be detected [as measured by methods such as Zeeman splitting, etc.]
Mode Amplitude	In helioseismology the magnitude of oscillation of waves

Polarization	of a particular geometry. Direction of the electric vector of an electromagnetic wave. The wave can be linearly polarized in any direction perpendicular to the direction of travel, circularly polarized (clockwise or counterclockwise), unpolarized, or mixtures of the above.
Stoke's Parameters	The four coordinates (usually called I, Q, U, and V) relative to a particular basis for the representation of the polarization state of an electromagnetic wave propagating through space.
Velocity	Rate of change of position. Also used for the average velocity of a collection of particles, also referred to as "bulk velocity".

**Photon Qualifier List**

Closed

Identifiers for terms which can be associated with a Photon Quantity.

Term	Definition
Average	The statistical mean; the sum of a set of values divided by the number of values in the set.
Circular	Relative to polarization, right-hand circularly polarized light is defined such that the electric field is rotating clockwise as seen by an observer towards whom the wave is moving. Left-hand circularly polarized light is defined such that the electric field is rotating counterclockwise as seen by an observer towards whom the wave is moving. The polarization of magnetohydrodynamic waves is specified with respect to the ambient mean magnetic field : right-hand polarized waves have a transverse electric field component which turns in a right-handed sense (that of the gyrating electrons) around the magnetic field.
Line of Sight	The line of sight is the line that connects the observer with the observed object. This expression is often used with measurements of Doppler velocity and magnetic field in magnetograms, where only the component of the vector field directed along the line of sight is measured.
Linear	Relative to polarization, confinement of the E-field vector to a given plane
Peak	The maximum value for the quantity in question, over a period of time which is usually equal to the cadence.
Variance	A measure of dispersion of a set of data points around their mean value. The expectation value of the squared deviations from the mean.
Vector	A quantity having both magnitude and direction, e.g. displacement, velocity, acceleration and force.

**Processing Level List**

Closed

Identifiers to characterize the amount and type of manipulation which has been applied to the sampled data.

Term	Definition
------	------------

Calibrated	Data wherein sensor outputs have been convolved with instrument response function, often irreversibly, to yield physical parameter values.
Uncalibrated	Duplicate data are removed from the data stream and data are time ordered. Values are not adjusted for any potential biases or external factors.

## Repository Name List

Open

For a current list see Identifiers for the location or facility where the product is stored. The repository name is selected from a list of established repositories. See <http://www.igpp.ucla.edu/spase/> for the list.

## Role List

Closed

Term	Definition
Co-Investigator	An individual who is a scientific peer and major participant for an investigation.
Data Producer	An individual who generated the resource and is familiar with its provenance.
General Contact	An individual who can provide information on a range of subjects or who can direct you to a domain expert.
Principal Investigator	An individual who is the administrative and scientific lead for an investigation.
Scientist	An individual who is an expert in the phenomenon and related physics represented by the resource.
Team Leader	An individual who is the scientific and administrative lead for an investigation.
Team Member	An individual who is a major participant in an investigation.
Technical Contact	An individual who can provide specific information with regard to the resource or supporting software

## Spectral Range List

Closed

Identifiers for names associated with wavelengths. Based on the ISO 21348 Solar Irradiance Standard. Additions have been made to extend the frequency ranges to include those used in space physics. Those additions are indicated in blue text. The "Total Solar Irradiance" category has not been included since it is a type of measurement and not a specific spectral range. See Appendix A - Comparison of Spectrum Domains for a comparison of the spectral ranges with other systems.

Term	Definition
Gamma Rays	Photons with a wavelength range: 0.00001 to 0.001 nm
Hard X-rays	Photons with a wavelength range: 0.001 to 0.1 nm
Infrared	Photons with a wavelength range: 760 to 1.00x10 <sup>6</sup> nm
Microwave	Photons with a wavelength range: 1.00x10 <sup>6</sup> to 1.50x10 <sup>7</sup> nm
Optical	Photons with a wavelength range: 380 to 760 nm
Radio Frequency	Photons with a wavelength range: 100,000 to 1.00x10 <sup>11</sup> nm
Ultraviolet	Photons with a wavelength range: 10 to 400 nm
X-Rays	Photons with a wavelength range: 0.001 ≤ x < 10 nm



## Sun List

Closed


Term	Definition
Chromosphere	The region of the Sun's (or a star's) atmosphere above the temperature minimum and below the Transition Region. The solar chromosphere is approximately 400 km to 2100 km above the photosphere, and characterized by temperatures from 4500 - 28000 K.
Corona	The outermost atmospheric region of the Sun or a star, characterized by ionization temperatures above $10^5$ K. The solar corona starts at about 2100 km above the photosphere; there is no generally defined upper limit.
Interior	The region inside the body which is not visible from outside the body.
Photosphere	The atmospheric layer of the Sun or a star from which continuum radiation, especially optical, is emitted to space. For the Sun, the photosphere is about 500 km thick.
Transition Region	A very narrow (<100 km) layer between the chromosphere and the corona where the temperature rises abruptly from about 8000 to about 500,000 K.

## Support List


Closed


Term	Definition
Other	Values, such as flags, that are not time tags, location data or measured or derived parameters.
Positional	The specification of the location of an object or measurement within a reference coordinate system. The position is usually expressed as a set of values corresponding to the location along a set of orthogonal axes together with the date/time of the observation.
Temporal	Pertaining to time.


## 9. Appendix A - Graphical Presentation of the SPASE Schema























## 10. Bibliography

National Solar Observatory Sacramento Peak

<http://www.sunspot.noao.edu/sunspot/pr/glossary.html>

Terms and Definitions

<http://www.pgd.hawaii.edu/eschool/glossary.htm>

International System of Units (SI)

<http://www.bipm.fr/en/si>

Base units: [http://www.bipm.fr/en/si/si\\_brochure/chapter2/2-1/#symbols](http://www.bipm.fr/en/si/si_brochure/chapter2/2-1/#symbols)

and those for Common derived units: [http://www.bipm.fr/en/si/derived\\_units/2-2-2.html](http://www.bipm.fr/en/si/derived_units/2-2-2.html)

ISO 8601:2004 - Date Format

[http://en.wikipedia.org/wiki/ISO\\_8601](http://en.wikipedia.org/wiki/ISO_8601)

- or -

<http://www.iso.ch/iso/en/CatalogueDetailPage.CatalogueDetail?CSNUMBER=40874>

- or -

<http://www.iso.org/iso/en/prods-services/popstds/datesandtime.html>

RFC 3339 - Date and Time on the Internet

The basis for the ISO 8601 standard. <http://www.ietf.org/rfc/rfc3339.txt>

RFC 1014 - XDR: External Data Representation standard

<http://www.faqs.org/rfcs/rfc1014.html>

## 11. Index

ASCII, 20  
 AVI, 21  
 Access Information, 20  
 Access Rights, 20  
 Access Rights List, 46  
 Access URL, 20  
 Acknowledgement, 20  
 Acronym, 20  
 Activity Index, 20  
 Address, 20  
 Aerosol, 20  
 Alfven Mach Number, 20  
 Alias, 20  
 Alpha Particle, 20  
 Antenna, 20  
 Aurora, 20  
 Availability, 21  
 Availability List, 46  
 Average, 21  
 Azimuthal Angle, 21  
 BZIP2, 21  
 Binary, 21  
 Bow Shock Crossing, 21  
 CDF, 22  
 CEF, 22  
 CEF 1, 22  
 CEF 2, 22  
 CGM, 22  
 Cadence, 21  
 Calibrated, 21  
 Cartesian, 21  
 Catalog, 21  
 Caveats, 21  
 Channeltron, 22  
 Charged Particle Flux, 22  
 Chromosphere, 22  
 Circular, 22  
 Co-Investigator, 22  
 Component, 22  
 Contact, 22  
 Coordinate Representation, 23  
 Coordinate Representation List, 46  
 Coordinate System, 23  
 Coordinate System Name, 23  
 Coordinate System Name List, 46  
 Corona, 23  
 Coronal Mass Ejection, 23  
 Counts, 23  
 Cross Spectrum, 23  
 Cylindrical, 23  
 DM, 24  
 Data Producer, 24  
 Dayside, 24  
 Description, 24  
 Deviation, 24  
 Dimension, 24  
 Display Cadence, 24  
 Display Data, 24  
 Dopplergram, 24  
 Double Sphere, 24  
 Dust, 25  
 Dynamic Spectra, 25  
 Electric, 25  
 Electric Field, 25  
 Electron, 25  
 Electron Drift Instrument, 25  
 Electrostatic Analyser, 25  
 Email, 25  
 Emissivity, 25  
 Encoding, 25  
 Encoding List, 49  
 End Date, 25  
 Energetic Particle Instrument, 25  
 Energetic Particles, 25  
 Energetic Solar Particle Event, 25  
 Energy, 26  
 Equivalent Width, 26  
 Exposure, 26  
 FITS, 26  
 Faraday Cup, 26  
 Field, 26  
 Field Physical Quantity, 26  
 Field Physical Quantity List, 49  
 Field Qualifier, 26  
 Field Qualifier List, 49  
 Fit, 26  
 Flux, 26  
 Flux Feedback, 26  
 Forbush Decrease, 27  
 Format, 27  
 Format List, 50  
 Fourier Transform Spectrograph, 27  
 Frequency, 27  
 GEI, 27  
 GEO, 27  
 GIF, 28  
 GSE, 28  
 GSEQ, 28

GSM, 28  
 GZIP, 28  
 Gamma Rays, 27  
 General Contact, 27  
 Geomagnetic Storm, 27  
 HAE, 28  
 HDF, 28  
 HDF 4, 28  
 HDF 5, 28  
 HEE, 28  
 HEEQ, 28  
 HF radar, 29  
 HG, 29  
 HGI, 29  
 HTML, 29  
 Hard X-rays, 28  
 Heat Flux, 28  
 Heliosphere, 28  
 Heliosphere List, 51  
 High Latitude, 29  
 IDFS, 29  
 IDL, 29  
 Image Intensity, 29  
 Imager, 29  
 Information URL, 29  
 Infrared, 29  
 Inner, 29  
 Input Resource ID, 29  
 Instrument, 29  
 Instrument ID, 30  
 Instrument Name, 30  
 Instrument Region, 30  
 Instrument Type, 30  
 Instrument Type List, 51  
 Intensity, 30  
 Interior, 30  
 Interplanetary Shock, 31  
 Investigation Name, 31  
 Ion, 31  
 Ion Composition, 31  
 Irradiance, 31  
 JPEG, 31  
 Keyword, 31  
 LGM, 31  
 Langmuir Probe, 31  
 Line Depth, 31  
 Line of Sight, 31  
 Linear, 31  
 Long Wire, 31  
 Low Latitude, 31  
 MAG, 31  
 MFA, 33  
 MPEG, 34  
 Magnetic, 32  
 Magnetic Field, 32  
 Magnetogram, 32  
 Magnetometer, 32  
 Magnetopause Crossing, 32  
 Magnetosheath, 32  
 Magnetosphere, 32  
 Magnetosphere List, 53  
 Magnetotail, 32  
 Magnitude, 32  
 Main, 32  
 Mass, 32  
 Mass Density, 32  
 Mass Spectrometer, 33  
 Measured, 33  
 Measurement Type, 33  
 Measurement Type List, 54  
 Microchannel Plate, 33  
 Microwave, 33  
 Mixed, 33  
 Mode Amplitude, 33  
 Molecule, 33  
 Moment, 33  
 N, 34  
 Name, 34  
 Near 1AU, 34  
 Near L1, 34  
 Near Surface, 34  
 Near Surface List, 55  
 NetCDF, 34  
 Neutral, 34  
 Neutral Atom Images, 34  
 Neutral Gas, 34  
 Nightside, 34  
 None, 34  
 Number Density, 34  
 Numerical Data, 34  
 Observatory, 35  
 Observatory Group, 35  
 Observatory Group List, 55  
 Observatory ID, 35  
 Observatory Name, 35  
 Observatory Name List, 55  
 Observed Region, 35  
 Offline, 35  
 Online, 35  
 Open, 35  
 Optical, 35  
 Organization Name, 36

Other, 36  
 Outer, 36  
 PDF, 37  
 PNG, 38  
 Parallel, 36  
 Particle, 36  
 Particle Correlator, 36  
 Particle Physical Quantity, 36  
 Particle Physical Quantity List, 55  
 Particle Qualifier, 36  
 Particle Qualifier List, 56  
 Particle Type, 36  
 Particle Type List, 56  
 Peak, 37  
 Perpendicular, 37  
 Person, 37  
 Person ID, 37  
 Person Name, 37  
 Phase-Space Density, 37  
 Phenomenon Type, 37  
 Phenomenon Type List, 57  
 Phone Number, 37  
 Photon, 37  
 Photon Physical Quantity, 38  
 Photon Physical Quantity List, 58  
 Photon Qualifier, 38  
 Photon Qualifier List, 59  
 Photosphere, 38  
 Physical Parameter, 38  
 Plasma Beta, 38  
 Polar Angle, 38  
 Polarization, 38  
 Positional, 39  
 Potential, 39  
 Poynting Flux, 39  
 Pressure, 39  
 Principal Investigator, 39  
 Processing Level List, 59  
 Profile, 39  
 Proton, 39  
 Provider ID, 39  
 Provider Processing Level, 39  
 Provider Release Date, 39  
 Provider Resource Name, 39  
 Provider Version, 39  
 Q, 39  
 Quadrispherical Analyser, 39  
 QuickTime, 39  
 RTF, 41  
 RTN, 41  
 Radiance, 39  
 Radio Frequency, 40  
 Radio Soundings, 40  
 Radio and Plasma Waves, 40  
 Repository Name, 40  
 Repository Name List, 60  
 Resonance Sounder, 40  
 Resource Header, 40  
 Resource ID, 40  
 Resource Name, 40  
 Restricted, 40  
 Role, 40  
 Role List, 60  
 SC, 41  
 SE, 41  
 SGI, 41  
 SM, 41  
 SPASE, 41  
 SR, 42  
 SR2, 42  
 SSE, 42  
 Scientist, 41  
 Search coil, 41  
 Size, 41  
 Soft X-rays, 41  
 Solar Flare, 41  
 Solar Wind Extreme, 41  
 Sonic Mach Number, 41  
 Spacecraft Potential Control, 41  
 Spatial Range, 42  
 Spectral Power Receiver, 42  
 Spectral Range List, 60  
 Spectral Range Name, 42  
 Spectrograph, 42  
 Spectrometer, 42  
 Spectrum, 42  
 Spherical, 42  
 Start Date, 43  
 Stoke's Parameters, 43  
 Sun, 43  
 Sun List, 60  
 Support, 43  
 Support List, 61  
 Surface, 43  
 TIFF, 44  
 TeX, 43  
 Team Leader, 43  
 Team Member, 43  
 Technical Contact, 43  
 Temperature, 43  
 Temporal, 43  
 Temporal Description, 43

Text, 43  
Thermal Plasma, 43  
Thermal Speed, 44  
Timespan, 44  
Transition Region, 44  
UDF, 44  
URL, 44  
Ultraviolet, 44  
Uncalibrated, 44  
Unicode, 44  
Units, 44  
Units Conversion, 44  
Variance, 45  
Vector, 45  
Velocity, 45  
Version, 45  
Waveform Receiver, 45  
Wavelength, 45  
Wavenumber, 45  
X-Rays, 45  
XDR, 45  
XML, 45  
Z, 45  
ZIP, 45

## 12. Change History

<b>0.99.1</b>		
2005-06-23	T.King	Removed duplicate entries; added Chris Harvey's definitions for Electron Drift; Particle Correlator and Spacecraft Potential Control
<b>0.99.2</b>		
2005-07-07	T.King	Corrected "Numerical Data" entry under Product
<b>0.99.3</b>		
2005-08-03	T.King	Added definitions supplied by J. Thieman, C. Harvey and T.King; Significant revision of document as suggested by Joe Hourcle
<b>0.99.4</b>		
2005-08-08	T.King	Restructured the taxonomy of elements to match the one suggested by A. Roberts; added definitions for new elements introduced in the new taxonomy
<b>0.99.5</b>		
2005-08-26	T.King	Clarified some definitions and corrected typographical errors based on comments from J. Thieman and J. Hourcle; changed data types of "Integer" to "Count" and "Double" to "Numeric"; added document elements to product resources; added catalog, display data to top list; included region descriptions from J. King with additions suggested by K. Reardon; and parameters loosely based on a model proposed by A.Roberts
<b>0.99.6</b>		
2005-09-07	T.King	Corrected the inclusion of Atmosphere-Ionosphere regions into the Magnetosphere; changed Surface to Ground; removed Body and references to it; added Spherical and Cartesian under Position; remove Ratio (Numerator and Denominator); change Upper Latitude to High Latitude, Lower to Low; introduced "Photon Context" and "Particle Context" as replacements for "Independent Variable"; removed "Provider" and "Manufacture" resources and replaced with ID pointers
<b>0.99.7</b>		
2005-09-08	T.King	Under Parameter add Description, Tensor Order; Change Photon Context and Particle Context to Independent Variable; Move Wavelength and Wave Number under Photon Independent Variable; Drop Speed from Particle Independent Variable; Move Polar Angle under Particle Independent Variable; Add Analysis Method under Field/Electric and Field/Magnetic; Add Wave Form, Spectra etc. under Analysis Method; Add Near 1AU under Heliosphere; Add Body under Atmosphere-Ionosphere, Magnetosphere and Ground; Add all planets + Moon under Body; Update definition of Magnetotail, etc. to be generic, add Earth examples; Change "Acceptable abbreviation" to "Conventional abbreviation" since abbreviations are not supported in the model
<b>0.99.8</b>		
2005-11-03	T.King	General clean-up and alignment with the schema agreed

upon at the APL meeting (Nov 2-4, 2005)

<b>0.99.9</b>		
2005-11-18	T.King, A. Robert	Incorporate comments from consortium members on the "final" draft before the release of version 1.0
<b>1.0.0</b>		
2005-11-22	T.King, and others	Incorporate comments from consortium members on the "final" draft before the release of version 1.0; Added Phenomenom Type list and defined terms in the list
<b>1.0.1</b>		
2006-01-03	T.King	Changes in value type for elements: Exposure, InputResourceID, RepositoryName, Size; Added elements: Pressure
<b>1.0.2</b>		
2006-03-07	T.King	Added "Project Scientist" to dictionary and "Role"; Added "Caveats" under "Instrument"; Added "Repository" resource class; Added "Registry" resource class
<b>1.0.3</b>		
2006-04-27	T.King	Added "Earth" as a enumeration with "Magnetosphere" as a member; changed "Observed Region" and "Instrument Region" to enumerations; changed definition of "Item" to indicate it is a value of an enumeration; Move "Access Rights" under "Access Information"; Made "Acknowledgement optional; change "HF Radar" to "Radar"; added "NCAR" as a "Format"; dropped N, Z, Q from dictionary; Moved Mass and Size under "Particle Physical Quantity" and changed to type item; added "Near Earth" under "Heliosphere" and added "Outside Bowshock" and "Orbital" under "Near Earth"; changed "Spectral Range Name" to "Spectral Range" for consistency; correct links to "Stoke's Parameters"
<b>1.1.0</b>		
2006-08-31	T.King	Removed "Orbital"; modified definition of "Near Earth"; changed "Instrument type" to allow multiple occurrences; made data type of "Mixed" text; added "Service" resource class; updated description of "Resource ID"; Added MAT_4, MAT_6, MAT_7 and VOTable as a Format; Added J2000 as a coordinate system; Added Base64 as an Encoding.; Added Parent ID, Energy Range, Frequency Range, Azimuthal Angle Range, Polar Angle Range, Atomic Number Range, Integral, Differential, Low and High.; Remove Coordinate System from Particle Physical Parameter; Updated Pressure definition; Add ObservatoryID under Instrument; Remove Observatory ID from Numerical Data and Display Data; Changed definition of Investigation Name; Remove Access Right from Display Data; Change Repository Name to Respoisotry ID under Access Information; Added Granule; Added Parameter Key under Physical Parameter; Add Release Date to Resource Header, Person, and Granule.; Changed "alias" to "alternate name".; Removed "Intrument Name" and "Observatory Name"; Add ChargeState to Particle Quantity; Add Field Component container; Add Statistics to Phenomenum Type.