

# Bicausal optimal transport for SDEs with irregular coefficients

---

**Benjamin A. Robinson** (University of Klagenfurt)

Austrian Stochastic Days — September 5, 2024

*Supported by Austrian Science Fund (FWF) projects (Y782-N25), (P35519), (P34743).*

# Bicausal optimal transport for SDEs with irregular coefficients

---

**Benjamin A. Robinson** (University of Klagenfurt)

Austrian Stochastic Days — September 5, 2024

*Supported by Austrian Science Fund (FWF) projects (Y782-N25), (P35519), (P34743).  
Joint work with*

**Michaela Szölgényi**  
University of Klagenfurt



# Bicausal optimal transport for SDEs with irregular coefficients

---

**Benjamin A. Robinson** (University of Klagenfurt)

Austrian Stochastic Days — September 5, 2024

*Supported by Austrian Science Fund (FWF) projects (Y782-N25), (P35519), (P34743).  
Joint work with*

**Michaela Szölgyenyi**  
University of Klagenfurt



**Julio Backhoff-Veraguas**  
University of Vienna



**Sigrid Källblad**  
KTH Stockholm



# Bicausal optimal transport for SDEs with irregular coefficients

---

**Benjamin A. Robinson** (University of Klagenfurt)

Austrian Stochastic Days — September 5, 2024

*Supported by Austrian Science Fund (FWF) projects (Y782-N25), (P35519), (P34743).  
Joint work with*

**Michaela Szölgényi**  
University of Klagenfurt

**Julio Backhoff-Veraguas**  
University of Vienna

**Sigrid Källblad**  
KTH Stockholm

*Adapted Wasserstein distance between the laws of SDEs  
(with J. Backhoff-Veraguas and S. Källblad) — arXiv:2209.03243, 2024*

*Bicausal optimal transport for SDEs with irregular coefficients  
(with M. Szölgényi) — arXiv:2403.09941, 2024*

# Comparing stochastic models

**Aim:** Compute a measure of model uncertainty

E.g. optimal stopping:

$$\mathbb{P} \mapsto v(\mathbb{P}) = \sup_{\tau \in \mathcal{T}} \mathbb{E}^{\mathbb{P}}[L(\tau, \omega)]$$

# Comparing stochastic models

**Aim:** Compute a measure of model uncertainty

E.g. optimal stopping:

$$\mathbb{P} \mapsto v(\mathbb{P}) = \sup_{\tau \in \mathcal{T}} \mathbb{E}^{\mathbb{P}}[L(\tau, \omega)]$$

**Want:**

- Appropriate topology on laws of stochastic processes
- Distance we can actually compute

# Comparing stochastic models

**Aim:** Compute a measure of model uncertainty

E.g. optimal stopping:

$$\mathbb{P} \mapsto v(\mathbb{P}) = \sup_{\tau \in \mathcal{T}} \mathbb{E}^{\mathbb{P}}[L(\tau, \omega)]$$

**Want:**

- Appropriate topology on laws of stochastic processes
- Distance we can actually compute

**SDEs:**

- Good computational methods available
- Rich class of models, beyond Lipschitz coefficients

## Main result

$$b, \bar{b}: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}, \quad \sigma, \bar{\sigma}: [0, T] \times \mathbb{R} \rightarrow [0, \infty),$$

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x_0,$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x_0.$$

$$\mu = \text{Law}(X), \quad \nu = \text{Law}(\bar{X})$$



# Main result

$$b, \bar{b}: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}, \sigma, \bar{\sigma}: [0, T] \times \mathbb{R} \rightarrow [0, \infty),$$

$$\begin{aligned} dX_t &= b_t(X_t)dt + \sigma_t(X_t)dB_t, & X_0 &= x_0, \\ d\bar{X}_t &= \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, & \bar{X}_0 &= x_0. \end{aligned}$$

$$\mu = \text{Law}(X), \nu = \text{Law}(\bar{X})$$

## Theorem [R., Szölgényi '24]

Under “weak assumptions” on the coefficients, we can compute an “appropriate distance”

# Main result

$$b, \bar{b}: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}, \sigma, \bar{\sigma}: [0, T] \times \mathbb{R} \rightarrow [0, \infty),$$

$$\begin{aligned} dX_t &= b_t(X_t)dt + \sigma_t(X_t)dB_t, & X_0 &= x_0, \\ d\bar{X}_t &= \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, & \bar{X}_0 &= x_0. \end{aligned}$$

$$\mu = \text{Law}(X), \nu = \text{Law}(\bar{X})$$

## Theorem [R., Szölgényi '24]

Under “weak assumptions” on the coefficients, we can compute an “appropriate distance” by

$$d_p(\mu, \nu)^p = \mathbb{E} \left[ \int_0^T |X_t - \bar{X}_t|^p dt \right], \quad \text{with } B = W.$$

# Application

E.g. optimal stopping:

$$\mathbb{P} \mapsto v(\mathbb{P}) = \sup_{\tau \in \mathcal{T}} \mathbb{E}^{\mathbb{P}}[L(\tau, \omega)]$$

E.g. optimal stopping:

$$\mathbb{P} \mapsto v(\mathbb{P}) = \sup_{\tau \in \mathcal{T}} \mathbb{E}^{\mathbb{P}}[L(\tau, \omega)]$$

## Theorem


[R., Szölgényi '24], [Acciaio, Backhoff-Veraguas, Zalashko '19]

$$\omega \mapsto L(t, \omega) \quad \text{Lipschitz on } (\Omega, \|\cdot\|_{L^p})$$


$\Rightarrow$

$$\mathbb{P} \mapsto v(\mathbb{P}) \quad \text{Lipschitz on } (\mathcal{P}_p(\Omega), d_p)$$

# Ingredients



# Ingredients




# Adapted Wasserstein distance

Define the Wasserstein distance w.r.t.  $L^p$  norm on  $\Omega := C([0, T], \mathbb{R})$  by

$$\mu, \nu \in \mathcal{P}_p(\Omega) \quad \rightsquigarrow \quad \mathcal{W}_p^p(\mu, \nu) := \inf_{\pi \in \text{Cpl}(\mu, \nu)} \mathbb{E}^\pi \left[ \int_0^T |\omega_t - \bar{\omega}_t|^p dt \right]$$

$$\text{Cpl}(\mu, \nu) = \{ \pi = \text{Law}(X, Y) : X \sim \mu, Y \sim \nu \}$$




# Adapted Wasserstein distance

Define the Wasserstein distance w.r.t.  $L^p$  norm on  $\Omega := C([0, T], \mathbb{R})$  by

$$\mu, \nu \in \mathcal{P}_p(\Omega) \quad \rightsquigarrow \quad \mathcal{W}_p^p(\mu, \nu) := \inf_{\pi \in \text{Cpl}(\mu, \nu)} \mathbb{E}^\pi \left[ \int_0^T |\omega_t - \bar{\omega}_t|^p dt \right]$$

$$\text{Cpl}(\mu, \nu) = \{ \pi = \text{Law}(X, Y) : X \sim \mu, Y \sim \nu \}$$






# Adapted Wasserstein distance

Define the **adapted** Wasserstein distance w.r.t.  $L^p$  norm on  $\Omega := C([0, T], \mathbb{R})$  by


$$\mu, \nu \in \mathcal{P}_p(\Omega) \quad \rightsquigarrow \quad \mathcal{AW}_p^p(\mu, \nu) := \inf_{\pi \in \mathbf{Cpl}_{bc}(\mu, \nu)} \mathbb{E}^\pi \left[ \int_0^T |\omega_t - \bar{\omega}_t|^p dt \right]$$

$$\mathbf{Cpl}_{bc}(\mu, \nu) = \{ \pi \in \mathbf{Cpl}(\mu, \nu) : \pi \text{ **bicausal** } \}$$

“ $\mathcal{F}_t^Y$  independent of  $\mathcal{F}_t^X$ , conditional on  $\mathcal{F}_t^X$ ” and vice-versa



# Ingredients



## Main result

$$b, \bar{b}: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}, \quad \sigma, \bar{\sigma}: [0, T] \times \mathbb{R} \rightarrow [0, \infty),$$

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x_0,$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x_0.$$

$$\mu = \text{Law}(X), \quad \nu = \text{Law}(\bar{X})$$

# Main result

$$b, \bar{b}: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}, \sigma, \bar{\sigma}: [0, T] \times \mathbb{R} \rightarrow [0, \infty),$$

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x_0,$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x_0.$$

$$\mu = \text{Law}(X), \nu = \text{Law}(\bar{X})$$

## Theorem [R., Szölgényi '24]

Under “weak assumptions” on the coefficients, we can compute an “appropriate distance”

# Main result

$$b, \bar{b}: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}, \sigma, \bar{\sigma}: [0, T] \times \mathbb{R} \rightarrow [0, \infty),$$

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x_0,$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x_0.$$

$$\mu = \text{Law}(X), \nu = \text{Law}(\bar{X})$$

## Theorem [R., Szölgényi '24]

Under “weak assumptions” on the coefficients, we can compute the adapted Wasserstein distance

# Main result

$$b, \bar{b}: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}, \quad \sigma, \bar{\sigma}: [0, T] \times \mathbb{R} \rightarrow [0, \infty),$$

$$\begin{aligned} dX_t &= b_t(X_t)dt + \sigma_t(X_t)dB_t, & X_0 &= x_0, \\ d\bar{X}_t &= \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, & \bar{X}_0 &= x_0. \end{aligned}$$

$$\mu = \text{Law}(X), \quad \nu = \text{Law}(\bar{X})$$

## Theorem [R., Szölgényi '24]

Under “weak assumptions” on the coefficients, we can compute the **adapted Wasserstein distance** by

$$\mathcal{AW}_p^p(\mu, \nu) = \mathbb{E} \left[ \int_0^T |X_t - \bar{X}_t|^p dt \right], \quad \text{with } B = W.$$

# Coupling SDEs

$$\begin{aligned}dX_t &= b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu \\d\bar{X}_t &= \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu\end{aligned}$$

**Lemma [Backhoff-Veraguas, Källblad, R. '24]**

Optimising over **bicausal couplings**  $\pi \in \text{Cpl}_{\text{bc}}(\mu, \nu)$

$\Leftrightarrow$

Optimising over **correlations** between  $B, W$

# Coupling SDEs

**Lemma [Backhoff-Veraguas, Källblad, R. '24]**


Optimising over bicausal couplings  $\pi \in \text{Cpl}_{\text{bc}}(\mu, \nu)$

$\Leftrightarrow$

Optimising over correlations between  $B, W$


**Product coupling**

$B, W$  independent




**Synchronous coupling**

Choose the same driving Brownian motion  $B = W$ .





# Ingredients



## Proof of main result

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

### Theorem [R., Szölgényi '24]

Under “weak assumptions” on the coefficients, we can compute the adapted Wasserstein distance by

$$\mathcal{AW}_p^p(\mu, \nu) = \mathbb{E} \left[ \int_0^T |X_t - \bar{X}_t|^p dt \right], \quad \text{with } B = W.$$

# Proof of main result

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

## Theorem [R., Szölgényi '24]

Under “weak assumptions” on the coefficients, we can compute the adapted Wasserstein distance by

$$\mathcal{AW}_p^p(\mu, \nu) = \mathbb{E} \left[ \int_0^T |X_t - \bar{X}_t|^p dt \right], \quad \text{with } B = W.$$

1. Discretise SDEs;

## Proof of main result

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

### Theorem [R., Szölgényi '24]

Under “weak assumptions” on the coefficients, we can compute the adapted Wasserstein distance by

$$\mathcal{AW}_p^p(\mu, \nu) = \mathbb{E} \left[ \int_0^T |X_t - \bar{X}_t|^p dt \right], \quad \text{with } B = W.$$

1. Discretise SDEs;
2. Solve discrete-time bicausal optimal transport problem;

## Proof of main result

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

### Theorem [R., Szölgényi '24]

Under “weak assumptions” on the coefficients, we can compute the adapted Wasserstein distance by

$$\mathcal{AW}_p^p(\mu, \nu) = \mathbb{E} \left[ \int_0^T |X_t - \bar{X}_t|^p dt \right], \quad \text{with } B = W.$$

1. Discretise SDEs;
2. Solve discrete-time bicausal optimal transport problem;
3. Pass to a limit.

## Proof of main result

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

### Theorem [R., Szölgényi '24]

Under “weak assumptions” on the coefficients, we can compute the adapted Wasserstein distance by

$$\mathcal{AW}_p^p(\mu, \nu) = \mathbb{E} \left[ \int_0^T |X_t - \bar{X}_t|^p dt \right], \quad \text{with } B = W.$$

1. Discretise SDEs;
2. Solve discrete-time bicausal optimal transport problem;
3. Pass to a limit.

## Key result in discrete time

$$\mu, \nu \in \mathcal{P}(\mathbb{R}^N) \quad \rightsquigarrow \quad \mathcal{AW}_p^p(\mu, \nu) := \inf_{\pi \in \text{Cpl}_{\text{bc}}(\mu, \nu)} \mathbb{E}^\pi \left[ \sum_{n=1}^N |X_n - Y_n|^p \right]$$

## Key result in discrete time

$$\mu, \nu \in \mathcal{P}(\mathbb{R}^N) \quad \rightsquigarrow \quad \mathcal{AW}_p^p(\mu, \nu) := \inf_{\pi \in \text{Cpl}_{\text{bc}}(\mu, \nu)} \mathbb{E}^\pi \left[ \sum_{n=1}^N |X_n - Y_n|^p \right]$$

### Knothe–Rosenblatt rearrangement

— generalisation of monotone rearrangement



## Key result in discrete time

$$\mu, \nu \in \mathcal{P}(\mathbb{R}^N) \quad \rightsquigarrow \quad \mathcal{AW}_p^p(\mu, \nu) := \inf_{\pi \in \text{Cpl}_{\text{bc}}(\mu, \nu)} \mathbb{E}^\pi \left[ \sum_{n=1}^N |X_n - Y_n|^p \right]$$

### Knothe–Rosenblatt rearrangement

— generalisation of monotone rearrangement

**Theorem** [Rüschendorf '85] [Beiglböck, Pammer, Posch '23]

For  $\mu, \nu$  **stochastically co-monotone**, the unique optimiser is the **Knothe–Rosenblatt rearrangement**.

This induces the **adapted weak topology**.

## Proof of main result

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

### Theorem [R., Szölgényi '24]

Under “weak assumptions” on the coefficients, we can compute the adapted Wasserstein distance by

$$\mathcal{AW}_p^p(\mu, \nu) = \mathbb{E} \left[ \int_0^T |X_t - \bar{X}_t|^p dt \right], \quad \text{with } B = W.$$

1. Discretise SDEs;
2. Solve discrete-time bicausal optimal transport problem;
3. Pass to a limit.

# A monotone numerical scheme

$$dX_t = b(X_t)dt \quad (\text{ODE})$$

## Euler scheme

$$X_0^h = X_0,$$

$$X_t^h = X_{kh}^h + b(X_{kh})(t - kh), \quad t \in (kh, (k+1)h].$$

## A monotone numerical scheme

$$dX_t = b(X_t)dt + dW_t \quad (\text{SDE})$$

### Euler–Maruyama scheme

$$X_0^h = X_0,$$

$$X_t^h = X_{kh}^h + b(X_{kh})(t - kh) + W_t - W_{kh}, \quad t \in (kh, (k+1)h].$$

## A monotone numerical scheme

$$dX_t = b(X_t)dt + dW_t \quad (\text{SDE})$$

### Euler–Maruyama scheme

$$X_0^h = X_0,$$

$$X_t^h = X_{kh}^h + b(X_{kh})(t - kh) + W_t - W_{kh}, \quad t \in (kh, (k+1)h].$$

Write  $X_k^h := X_{kh}^h$  and  $\mu^h = \text{Law}((X_k^h)_k)$ .

# A monotone numerical scheme

$$dX_t = b(X_t)dt + dW_t \quad (\text{SDE})$$

## Euler–Maruyama scheme

$$X_0^h = X_0,$$

$$X_t^h = X_{kh}^h + b(X_{kh})(t - kh) + W_t - W_{kh}, \quad t \in (kh, (k+1)h].$$

Write  $X_k^h := X_{kh}^h$  and  $\mu^h = \text{Law}((X_k^h)_k)$ .

## Remark

$X_k^h \mapsto X_{(k+1)}^h$  is **increasing** if  $b$  is Lipschitz,  $h \ll 1$

# A monotone numerical scheme

$$dX_t = b(X_t)dt + dW_t \quad (\text{SDE})$$

## Euler–Maruyama scheme

$$X_0^h = X_0,$$

$$X_t^h = X_{kh}^h + b(X_{kh})(t - kh) + W_t - W_{kh}, \quad t \in (kh, (k+1)h].$$

Write  $X_k^h := X_{kh}^h$  and  $\mu^h = \text{Law}((X_k^h)_k)$ .

## Remark

$X_k^h \mapsto X_{(k+1)}^h$  is **increasing** if  $b$  is Lipschitz,  $h \ll 1$

# A monotone numerical scheme

$$dX_t = b(X_t)dt + \sigma(X_t)dW_t \quad (\text{SDE})$$

## Monotone Euler–Maruyama scheme

$$X_0^h = X_0,$$

$$X_t^h = X_{kh}^h + b(X_{kh})(t - kh) + \sigma(X_{kh})(W_t^h - W_{kh}^h), \quad t \in (kh, (k+1)h].$$

$$W_t^h - W_{kh}^h = W_{t \wedge \tau_k^h} - W_{kh}, \quad \tau_k^h := \inf\{t > kh : |W_t - W_{kh}| > A_h\}$$



# A monotone numerical scheme

$$dX_t = b(X_t)dt + \sigma(X_t)dW_t \quad (\text{SDE})$$

## Monotone Euler–Maruyama scheme

$$X_0^h = X_0,$$

$$X_t^h = X_{kh}^h + b(X_{kh})(t - kh) + \sigma(X_{kh})(W_t^h - W_{kh}^h), \quad t \in (kh, (k+1)h].$$

$$W_t^h - W_{kh}^h = W_{t \wedge \tau_k^h} - W_{kh}, \quad \tau_k^h := \inf\{t > kh : |W_t - W_{kh}| > A_h\}$$

$$\text{Write } X_k^h := X_{kh}^h \quad \text{and} \quad \mu^h = \text{Law}((X_k^h)_k).$$

# A monotone numerical scheme

$$dX_t = b(X_t)dt + \sigma(X_t)dW_t \quad (\text{SDE})$$

## Monotone Euler–Maruyama scheme

$$X_0^h = X_0,$$

$$X_t^h = X_{kh}^h + b(X_{kh})(t - kh) + \sigma(X_{kh})(W_t^h - W_{kh}^h), \quad t \in (kh, (k+1)h].$$

$$W_t^h - W_{kh}^h = W_{t \wedge \tau_k^h} - W_{kh}, \quad \tau_k^h := \inf\{t > kh : |W_t - W_{kh}| > A_h\}$$

$$\text{Write } X_k^h := X_{kh}^h \quad \text{and} \quad \mu^h = \text{Law}((X_k^h)_k).$$

## Lemma [Backhoff-Veraguas, Källblad, R. '24]

For  $b, \sigma$  Lipschitz, the monotone Euler–Maruyama scheme is stochastically increasing.

Hence the Knothe–Rosenblatt rearrangement is optimal for  $\mu^h, \nu^h$ .

# Proof of main result

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

## Theorem [R., Szölgényi '24]

Under “**weak assumptions**” on the coefficients, we can compute the adapted Wasserstein distance by

$$\mathcal{AW}_p^p(\mu, \nu) = \mathbb{E} \left[ \int_0^T |X_t - \bar{X}_t|^p dt \right], \quad \text{with } B = W.$$

1. **Discretise SDEs**;
2. Solve discrete-time bicausal optimal transport problem;
3. Pass to a limit.

# Proof of main result

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

## Assumptions

- (A) discontinuous drift with exponential growth (time-homog.);
- (B) bounded measurable drift,  $\alpha$ -Hölder and uniformly elliptic  $\sigma$ ;
- (C) **continuous** coefficients, **linear** growth, **pathwise uniqueness**.

# Proof of main result

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

## Assumptions

- (A) discontinuous drift with exponential growth (time-homog.);
- (B) **bounded measurable** drift,  $\alpha$ -Hölder and **uniformly elliptic**  $\sigma$ ;
- (C) continuous coefficients, linear growth, pathwise uniqueness.

# Proof of main result

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

## Assumptions

- (A) **discontinuous** drift with **exponential** growth (time-homog.);
- (B) bounded measurable drift,  $\alpha$ -Hölder and uniformly elliptic  $\sigma$ ;
- (C) continuous coefficients, linear growth, pathwise uniqueness.

# Transformation-based semi-implicit Euler scheme

## Assumption (A)

Drift  $b: \mathbb{R} \rightarrow \mathbb{R}$  satisfies the following conditions **piecewise**:

- absolute continuity
- one-sided Lipschitz condition
- two-sided local Lipschitz condition
- exponential growth



# Transformation-based semi-implicit Euler scheme

## Assumption (A)

Drift  $b: \mathbb{R} \rightarrow \mathbb{R}$  satisfies the following conditions piecewise:

- absolute continuity
- one-sided Lipschitz condition
- two-sided local Lipschitz condition
- exponential growth





# Transformation-based semi-implicit Euler scheme

## Assumption (A)

Drift  $b: \mathbb{R} \rightarrow \mathbb{R}$  satisfies the following conditions piecewise:

- absolute continuity
- one-sided Lipschitz condition
- two-sided local Lipschitz condition
- exponential growth



Diffusion  $\sigma: \mathbb{R} \rightarrow [0, \infty)$  satisfies

- global Lipschitz condition
- $\sigma(\xi_k) \neq 0$ , for  $k \in \{1, \dots, m\}$  — no uniform ellipticity

# Transformation-based semi-implicit Euler scheme

Under Assumption (A) , the scheme is constructed as follows:

1. Apply the transformation  $G$  from [Leobacher, Szölgényi '17] to (SDE),

$$Z = G(X)$$

$$dZ_t = \tilde{b}(Z_t)dt + \tilde{\sigma}(Z_t)dW_t$$

- $\tilde{b}$  one-sided Lipschitz, locally Lip., a.c., exponential growth,
- $\tilde{\sigma}$  Lipschitz

# Transformation-based semi-implicit Euler scheme

Under Assumption (A) , the scheme is constructed as follows:

1. Apply the **transformation**  $G$  from [Leobacher, Szölgényi '17] to (SDE),
2. Apply a **semi-implicit Euler scheme** with **truncated Brownian increments** to the transformed SDE,

$$Z = G(X)$$

$$dZ_t = \tilde{b}(Z_t)dt + \tilde{\sigma}(Z_t)dW_t$$

$$Z_{(k+1)h}^h = Z_{kh}^h + \tilde{b}(Z_{(k+1)h}^h) \cdot h + \tilde{\sigma}(Z_{kh}^h)(W_{(k+1)h}^h - W_{kh}^h)$$

# Transformation-based semi-implicit Euler scheme

Under Assumption (A) , the scheme is constructed as follows:

1. Apply the **transformation**  $G$  from [Leobacher, Szölgényi '17] to (SDE),
2. Apply a **semi-implicit Euler scheme** with **truncated Brownian increments** to the transformed SDE,
3. **Transform back**.

$$Z = G(X)$$

$$dZ_t = \tilde{b}(Z_t)dt + \tilde{\sigma}(Z_t)dW_t$$

$$Z_{(k+1)h}^h = Z_{kh}^h + \tilde{b}(Z_{(k+1)h}^h) \cdot h + \tilde{\sigma}(Z_{kh}^h)(W_{(k+1)h}^h - W_{kh}^h)$$

$$X_{kh}^h = G^{-1}(Z_{kh}^h)$$

# Transformation-based semi-implicit Euler scheme

Under Assumption (A) , the scheme is constructed as follows:


1. Apply the **transformation**  $G$  from [Leobacher, Szölgényi '17] to (SDE),
2. Apply a **semi-implicit Euler scheme** with **truncated Brownian increments** to the transformed SDE,
3. **Transform back**.

## Theorem [R., Szölgényi '24]

Let  $(b, \sigma)$  satisfy Assumption (A). Then (SDE) admits a unique strong solution and, for all  $p \geq 1$ , there exists  $C_p \geq 0$  such that

$$\mathbb{E} \left[ |X_T - X_T^h|^p \right]^{\frac{1}{p}} \leq \begin{cases} C_p h^{\frac{1}{2}}, & p \in [1, 2], \\ C_p h^{\frac{1}{p(p-1)}}, & p \geq 2. \end{cases}$$

# Ingredients



# Main result

## Assumptions

(A) **discontinuous** drift with **exponential** growth (time-homog.);

# Main result

## Assumptions

- (A) discontinuous drift with exponential growth (time-homog.);
- (B) **bounded measurable** drift,  $\alpha$ -Hölder and **uniformly elliptic**  $\sigma$ ;



# Main result

## Assumptions

- (A) discontinuous drift with exponential growth (time-homog.);
- (B) bounded measurable drift,  $\alpha$ -Hölder and uniformly elliptic  $\sigma$ ;
- (C) continuous coefficients, linear growth, pathwise uniqueness.

# Main result

## Assumptions

- (A) discontinuous drift with exponential growth (time-homog.);
- (B) bounded measurable drift,  $\alpha$ -Hölder and uniformly elliptic  $\sigma$ ;
- (C) continuous coefficients, linear growth, pathwise uniqueness.

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

# Main result

## Assumptions

- (A) discontinuous drift with exponential growth (time-homog.);
- (B) bounded measurable drift,  $\alpha$ -Hölder and uniformly elliptic  $\sigma$ ;
- (C) continuous coefficients, linear growth, pathwise uniqueness.

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

## Main Theorem [R., Szölgényi '24]

Let  $(b, \sigma)$  and  $(\bar{b}, \bar{\sigma})$  each satisfy **one of assumptions** (A), (B), (C). Then, for  $p \in [1, \infty)$ , the adapted Wasserstein distance is given by

$$\mathcal{AW}_p^p(\mu, \nu) = \mathbb{E} \left[ \int_0^T |X_t - \bar{X}_t|^p dt \right], \text{ with } B = W$$

# Main result

## Assumptions

- (A) discontinuous drift with exponential growth (time-homog.);
- (B) bounded measurable drift,  $\alpha$ -Hölder and uniformly elliptic  $\sigma$ ;
- (C) continuous coefficients, linear growth, pathwise uniqueness.

$$dX_t = b_t(X_t)dt + \sigma_t(X_t)dB_t, \quad X_0 = x \rightsquigarrow \text{Law}(X) = \mu$$

$$d\bar{X}_t = \bar{b}_t(\bar{X}_t)dt + \bar{\sigma}_t(\bar{X}_t)dW_t, \quad \bar{X}_0 = x \rightsquigarrow \text{Law}(\bar{X}) = \nu$$

## Main Theorem [R., Szölgényi '24]

Let  $(b, \sigma)$  and  $(\bar{b}, \bar{\sigma})$  each satisfy **one of assumptions** (A), (B), (C). Then, for  $p \in [1, \infty)$ , the adapted Wasserstein distance is given by

$$\mathcal{AW}_p^p(\mu, \nu) = \mathbb{E} \left[ \int_0^T |X_t - \bar{X}_t|^p dt \right], \text{ with } B = W$$

**Synchronous coupling** solves general bicausal transport problem

# Summary

- We compute adapted Wasserstein distance between SDEs with irregular coefficients
- We prove strong convergence rates for a numerical scheme for SDEs with discontinuous and exponentially growing drift

References:

