

A regularized Kellerer theorem in arbitrary dimension

Benjamin A. Robinson

University of Vienna

November 20, 2023 — Vienna Probability Seminar

A regularized Kellner theorem in arbitrary dimension

Benjamin A. Robinson

University of Vienna

November 20, 2023 — Vienna Probability Seminar

Joint work with

Gudmund Pammer

ETH Zürich



Walter Schachermayer

University of Vienna



Motivating problem

Given probability measures μ, ν on \mathbb{R}^d do there exist random variables M_0, M_1 such that

- $\mathbb{E}[M_1 | M_0] = M_0$ (martingale property)

Motivating problem

Given probability measures μ, ν on \mathbb{R}^d do there exist random variables M_0, M_1 such that

- $\mathbb{E}[M_1 | M_0] = M_0$ (martingale property) and
- $\text{Law}(M_0) = \mu$ and $\text{Law}(M_1) = \nu$ (mimicking property)?

Motivating problem

Given probability measures μ, ν on \mathbb{R}^d do there exist random variables M_0, M_1 such that

- $\mathbb{E}[M_1 | M_0] = M_0$ (martingale property) and
- $\text{Law}(M_0) = \mu$ and $\text{Law}(M_1) = \nu$ (mimicking property)?

Necessary condition: $\mu \preceq \nu$ in convex order

Motivating problem

Given probability measures μ, ν on \mathbb{R}^d do there exist random variables M_0, M_1 such that

- $\mathbb{E}[M_1 | M_0] = M_0$ (martingale property) and
- $\text{Law}(M_0) = \mu$ and $\text{Law}(M_1) = \nu$ (mimicking property)?

Necessary condition: $\mu \preceq \nu$ in convex order

For any convex function $v : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$\int v d\mu = \mathbb{E}[v(M_0)] = \mathbb{E}[v(\mathbb{E}[M_1 | M_0])] \quad \text{martingale}$$

Motivating problem

Given probability measures μ, ν on \mathbb{R}^d do there exist random variables M_0, M_1 such that

- $\mathbb{E}[M_1 | M_0] = M_0$ (martingale property) and
- $\text{Law}(M_0) = \mu$ and $\text{Law}(M_1) = \nu$ (mimicking property)?

Necessary condition: $\mu \preceq \nu$ in convex order

For any convex function $v : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$\begin{aligned} \int v d\mu &= \mathbb{E}[v(M_0)] = \mathbb{E}[v(\mathbb{E}[M_1 | M_0])] \\ &\leq \mathbb{E}[\mathbb{E}[v(M_1) | M_0]] \quad \text{Jensen} \end{aligned}$$

Motivating problem

Given probability measures μ, ν on \mathbb{R}^d do there exist random variables M_0, M_1 such that

- $\mathbb{E}[M_1 | M_0] = M_0$ (martingale property) and
- $\text{Law}(M_0) = \mu$ and $\text{Law}(M_1) = \nu$ (mimicking property)?

Necessary condition: $\mu \preceq \nu$ in convex order

For any convex function $v : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$\begin{aligned}\int v d\mu &= \mathbb{E}[v(M_0)] = \mathbb{E}[v(\mathbb{E}[M_1 | M_0])] \\ &\leq \mathbb{E}[\mathbb{E}[v(M_1) | M_0]] \\ &= \mathbb{E}[v(M_1)] = \int v d\nu.\end{aligned}$$

Motivating problem

Given probability measures μ, ν on \mathbb{R}^d do there exist random variables M_0, M_1 such that

- $\mathbb{E}[M_1 | M_0] = M_0$ (martingale property) and
- $\text{Law}(M_0) = \mu$ and $\text{Law}(M_1) = \nu$ (mimicking property)?

Necessary condition: $\mu \preceq \nu$ in convex order

For any convex function $v : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$\int v d\mu \leq \int v d\nu.$$

Motivating problem

Given probability measures μ, ν on \mathbb{R}^d do there exist random variables M_0, M_1 such that

- $\mathbb{E}[M_1 | M_0] = M_0$ (martingale property) and
- $\text{Law}(M_0) = \mu$ and $\text{Law}(M_1) = \nu$ (mimicking property)?

Necessary condition: $\mu \preceq \nu$ in convex order

For any convex function $v : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$\int v d\mu \leq \int v d\nu.$$

... also sufficient [Strassen '65]

Problem statement

Given a family of probability measures $(\mu_t)_{t \in I}$ on \mathbb{R}^d , does there exist a **mimicking martingale** M such that

$$\text{Law}(M_t) = \mu_t, \quad \forall t \in I?$$

Problem statement

Given a family of probability measures $(\mu_t)_{t \in I}$ on \mathbb{R}^d , does there exist a **mimicking martingale** M such that

$$\text{Law}(M_t) = \mu_t, \quad \forall t \in I?$$

Necessary condition

For any convex function $v : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$\int v d\mu_s \leq \int v d\mu_t, \quad s \leq t.$$

Peacocks

Assume that μ is a **peacock**; i.e. for any convex function

$v : \mathbb{R}^d \rightarrow \mathbb{R}$,


$$\int v d\mu_s \leq \int v d\mu_t, \quad s \leq t.$$

Peacocks

Assume that μ is a **peacock**; i.e. for any convex function $v : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$\int v d\mu_s \leq \int v d\mu_t, \quad s \leq t.$$

Processus Croissant pour l'Ordre Convexe



[Hirsch, Profetta, Roynette, Yor '11]

Problem statement

Given a **peacock** $(\mu_t)_{t \in I}$ on \mathbb{R}^d , does there exist a **mimicking martingale** M such that

$$\text{Law}(M_t) = \mu_t, \quad \forall t \in I?$$

Problem statement

Given a **peacock** $(\mu_t)_{t \in I}$ on \mathbb{R}^d , does there exist a **mimicking martingale** M such that

$$\text{Law}(M_t) = \mu_t, \quad \forall t \in I?$$

Yes – [Strassen '65], [Doob '68], [Hirsch, Roynette '13]

Problem statement

Given a **peacock** $(\mu_t)_{t \in I}$ on \mathbb{R}^d , does there exist a **mimicking martingale** M such that

$$\text{Law}(M_t) = \mu_t, \quad \forall t \in I?$$

Desirable properties

Problem statement

Given a **peacock** $(\mu_t)_{t \in I}$ on \mathbb{R}^d , does there exist a **mimicking martingale** M such that

$$\text{Law}(M_t) = \mu_t, \quad \forall t \in I?$$

Desirable properties

- **Markovianity**

Problem statement

Given a **peacock** $(\mu_t)_{t \in I}$ on \mathbb{R}^d , does there exist a **mimicking martingale** M such that

$$\text{Law}(M_t) = \mu_t, \quad \forall t \in I?$$

Desirable properties

- **strong** Markovianity

Problem statement

Given a **peacock** $(\mu_t)_{t \in I}$ on \mathbb{R}^d , does there exist a **mimicking martingale** M such that

$$\text{Law}(M_t) = \mu_t, \quad \forall t \in I?$$

Desirable properties

- strong Markovianity
- **continuity** of paths

Problem statement

Given a **peacock** $(\mu_t)_{t \in I}$ on \mathbb{R}^d , does there exist a **mimicking martingale** M such that

$$\text{Law}(M_t) = \mu_t, \quad \forall t \in I?$$

Desirable properties

- strong Markovianity
- continuity of paths
- **uniqueness**

[Strassen '65]

Given a discrete-time peacock $(\mu_n)_{n \in N}$ on \mathbb{R}^d ,

[Strassen '65]

Given a discrete-time peacock $(\mu_n)_{n \in N}$ on \mathbb{R}^d , there exists a mimicking **Markov martingale** M .

[Strassen '65]

Given a discrete-time peacock $(\mu_n)_{n \in \mathbb{N}}$ on \mathbb{R}^d , there exists a mimicking Markov martingale M .

Volker Strassen *The Existence of Probability Measures with Given Marginals*, Ann. Math. Stat. 1965

Continuous time, $d = 1$

[Kellerer '72]

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R} ,

Continuous time, $d = 1$

[Kellerer '72]

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R} , there exists a mimicking **strong Markov martingale** M .

Continuous time, $d = 1$

[Kellerer '72]

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R} , there exists a mimicking strong Markov martingale M .

Hans G. Kellerer *Markov-Komposition und eine Anwendung auf Martingale*, Math. Ann. 1972

Continuous time, $d = 1$

[Kellerer '72]

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R} , there exists a mimicking strong Markov martingale M .

Hans G. Kellerer *Markov-Komposition und eine Anwendung auf Martingale*, Math. Ann. 1972

Subsequent contributions (**incomplete!**)

Albin, Baker, Beiglböck, Brücknerhoff, Boubel, Donati-Martin, Hamza, Hirsch, Huesmann, Juillet, Källblad, Klebaner, Lowther, Profetta, Roynette, Stebegg, Tan, Touzi, Yor, ...

Continuous time, $d = 1$

[Kellerer '72]

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R} , there exists a mimicking strong Markov martingale M .

[Lowther '08–10]

- M is the **unique** strong Markov mimicking martingale

Continuous time, $d = 1$

[Kellerer '72]

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R} , there exists a mimicking strong Markov martingale M .

[Lowther '08–10]

- M is the unique strong Markov mimicking martingale
- $t \mapsto M_t$ is **continuous**, if $t \mapsto \mu_t$ is weakly continuous with convex support.

Continuous time, $d = 1$

[Kellerer '72]

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R} , there exists a mimicking strong Markov martingale M .

[Lowther '08–10]

- M is the unique strong Markov mimicking martingale
- $t \mapsto M_t$ is continuous, if $t \mapsto \mu_t$ is weakly continuous with convex support.

Continuous time, $d \geq 2$

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^d , $d \geq 2$, does there exist a mimicking Markov martingale?

Continuous time, $d \geq 2$

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^d , $d \geq 2$, does there exist a mimicking Markov martingale?

Existing literature ...

Continuous time, $d \geq 2$

Given a **continuous-time peacock** $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^d , $d \geq 2$, does there exist a mimicking **Markov martingale**?

Existing literature ...

... without the Markov property

[Doob '68] (compact support)

Continuous time, $d \geq 2$

Given a **continuous-time peacock** $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^d , $d \geq 2$, does there exist a mimicking **Markov martingale**?

Existing literature ...

... without the Markov property

[Doob '68] (compact support), [Hirsch, Roynette '13] (\mathbb{R}^d)

Continuous time, $d \geq 2$

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^d , $d \geq 2$, does there exist a mimicking Markov martingale?

Existing literature ...

... with the Markov property

Continuous time, $d \geq 2$

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^d , $d \geq 2$, does there exist a mimicking Markov martingale?

Existing literature ...

... with the Markov property

no known results

Continuous time, $d \geq 2$

Given a continuous-time peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^d , $d \geq 2$, does there exist a mimicking Markov martingale?

Theorem 1 [Pammer, R., Schachermayer '23+]

There exists a strong Markov martingale diffusion mimicking a *regularized* continuous-time peacock on \mathbb{R}^d , $d \in \mathbb{N}$.

Main result

Weakly continuous \mathbb{R}^d -valued square-integrable peacock $(\mu_t)_{t \in [0,1]}$.

Regularize with a Gaussian $\mu_t^r := \mu_t * \gamma^{\varepsilon(t+\delta)}$

Main result

Weakly continuous \mathbb{R}^d -valued square-integrable peacock $(\mu_t)_{t \in [0,1]}$.
Regularize with a Gaussian $\mu_t^r := \mu_t * \gamma^{\varepsilon(t+\delta)}$

Theorem 1 [Pammer, R., Schachermayer '23+]

There exists a measurable $(t, x) \mapsto \sigma_t^r(x)$ that is **locally Lipschitz** in x and **non-degenerate**, uniformly in $t \in [0, 1]$, and a Brownian motion B such that $\text{Law}(M_t^r) = \mu_t^r$, for all $t \in [0, 1]$, where

$$dM_t^r = \sigma_t^r(M_t^r)dB_t.$$

Main result

Weakly continuous \mathbb{R}^d -valued square-integrable peacock $(\mu_t)_{t \in [0,1]}$.
Regularize with a Gaussian $\mu_t^r := \mu_t * \gamma^{\varepsilon(t+\delta)}$

Theorem 1 [Pammer, R., Schachermayer '23+]

There exists a measurable $(t, x) \mapsto \sigma_t^r(x)$ that is locally Lipschitz in x and non-degenerate, uniformly in $t \in [0, 1]$, and a Brownian motion B such that $\text{Law}(M_t^r) = \mu_t^r$, for all $t \in [0, 1]$, where

$$dM_t^r = \sigma_t^r(M_t^r)dB_t.$$

Moreover:

- M^r is a strong Markov martingale with continuous paths;

Main result

Weakly continuous \mathbb{R}^d -valued square-integrable peacock $(\mu_t)_{t \in [0,1]}$.
Regularize with a Gaussian $\mu_t^r := \mu_t * \gamma^{\varepsilon(t+\delta)}$

Theorem 1 [Pammer, R., Schachermayer '23+]

There exists a measurable $(t, x) \mapsto \sigma_t^r(x)$ that is locally Lipschitz in x and non-degenerate, uniformly in $t \in [0, 1]$, and a Brownian motion B such that $\text{Law}(M_t^r) = \mu_t^r$, for all $t \in [0, 1]$, where

$$dM_t^r = \sigma_t^r(M_t^r)dB_t.$$

Moreover:

- M^r is a strong Markov martingale with continuous paths;
- The result does not hold without regularization;

Main result

Weakly continuous \mathbb{R}^d -valued square-integrable peacock $(\mu_t)_{t \in [0,1]}$.
Regularize with a Gaussian $\mu_t^r := \mu_t * \gamma^{\varepsilon(t+\delta)}$

Theorem 1 [Pammer, R., Schachermayer '23+]

There exists a measurable $(t, x) \mapsto \sigma_t^r(x)$ that is locally Lipschitz in x and non-degenerate, uniformly in $t \in [0, 1]$, and a Brownian motion B such that $\text{Law}(M_t^r) = \mu_t^r$, for all $t \in [0, 1]$, where

$$dM_t^r = \sigma_t^r(M_t^r)dB_t.$$

Moreover:

- M^r is a strong Markov martingale with continuous paths;
- The result does not hold without regularization;
- There is no uniqueness for $d \geq 2$.

Proof idea

- Discretize and solve discrete-time problem;
- Pass to the continuous-time limit.

Proof idea

- Discretize and solve discrete-time problem;
- Pass to the continuous-time limit.

Major obstacle is that the following sets are not closed w.r.t. f.d.d. convergence:

- (strong) Markov processes,
- continuous martingales.

Proof idea

- Discretize and solve discrete-time problem;
- Pass to the continuous-time limit.

Major obstacle is that the following sets are not closed w.r.t. f.d.d. convergence:

- (strong) Markov processes,
- continuous martingales.

In **dimension one**, compactness w.r.t. f.d.d. convergence holds for the sets of processes that are

- càdlàg and **Lipschitz Markov** [Kellerer '72], [Lowther '09],
- **almost-continuous diffusions** [Lowther '09].

Proof idea

- Discretize and solve discrete-time problem;
- Pass to the continuous-time limit.

Major obstacle is that the following sets are not closed w.r.t. f.d.d. convergence:

- (strong) Markov processes,
- continuous martingales.

In dimension one, compactness w.r.t. f.d.d. convergence holds for the sets of processes that are

- càdlàg and Lipschitz Markov [Kellerer '72], [Lowther '09],
- almost-continuous diffusions [Lowther '09].

In \mathbb{R}^2 , these notions no longer help us [Lowther '09], [Juillet '16]!

Proof idea I

Discretize and take Bass martingales from μ_{t_k} to $\mu_{t_{k+1}}$ to get a diffusion process

Proof idea I

Discretize and take Bass martingales from μ_{t_k} to $\mu_{t_{k+1}}$ to get a diffusion process

Optimal Transport

$$\inf_{\substack{\pi \in \text{Cpl}(\mu_0, \mu_1) \\ (X, Y) \sim \pi}} \mathbb{E}[c(X, Y)]$$

$$\text{Cpl}(\mu_0, \mu_1) = \{\pi: \pi(A \times \mathbb{R}^d) = \mu_0(A), \pi(\mathbb{R}^d \times B) = \mu_1(B)\}$$

Proof idea I

Discretize and take Bass martingales from μ_{t_k} to $\mu_{t_{k+1}}$ to get a diffusion process

Martingale Optimal Transport

$$\inf_{\substack{\pi \in \mathcal{M}(\mu_0, \mu_1) \\ (X, Y) \sim \pi}} \mathbb{E}[c(X, Y)]$$

$$\mathcal{M}(\mu_0, \mu_1) = \{\pi \in \text{Cpl}(\mu_0, \mu_1) : (U, V) \sim \pi \Rightarrow \mathbb{E}[V \mid U] = U\}$$

[Hobson, Neuberger '12], [Beiglböck, Juillet '16]

Proof idea I

Discretize and take Bass martingales from μ_{t_k} to $\mu_{t_{k+1}}$ to get a diffusion process

Martingale Optimal Transport

$$\inf_{\substack{\pi \in \mathcal{M}(\mu_0, \mu_1) \\ (X, Y) \sim \pi}} \mathbb{E}[c(X, Y)] \text{ “=” } \mathbb{E}[c(M_0, M_1)]$$

$$(M_t)_{t \in [0, 1]} \quad \text{Bass martingale}$$


[Backhoff, Beiglböck, Huesmann, Källblad '19]

[Backhoff, Beiglböck, Schachermayer, Tschiderer '23]

Proof idea I

Discretize and take Bass martingales from μ_{t_k} to $\mu_{t_{k+1}}$ to get a diffusion process

Bass martingales




$$M_t = \mathbb{E}[\nabla v(B_1) \mid \mathcal{F}_t], \quad v: \mathbb{R}^d \rightarrow \mathbb{R} \text{ convex}, \quad B_0 \sim \alpha.$$

Proof idea I

Discretize and take Bass martingales from μ_{t_k} to $\mu_{t_{k+1}}$ to get a diffusion process

Bass martingales



$$M_t = \mathbb{E}[\nabla v(B_1) \mid \mathcal{F}_t] = \nabla v_t(B_t), \quad v_t: \mathbb{R}^d \rightarrow \mathbb{R} \text{ convex}, \quad B_0 \sim \alpha.$$

Proof idea I

Discretize and take Bass martingales from μ_{t_k} to $\mu_{t_{k+1}}$ to get a diffusion process

Bass martingales



Proof idea I

Discretize and take Bass martingales from μ_{t_k} to $\mu_{t_{k+1}}$ to get a diffusion process

Bass martingales



Proof idea II

Regularize with a Gaussian and make a **Markovian projection**

Proof idea II

Regularize with a Gaussian and make a **Markovian projection**

Mimicking Itô processes [Krylov '85], [Gyöngy '85], [Brunick, Shreve '13]

$$dX_t = \sigma_t dW_t$$

$$d\hat{X}_t = \hat{\sigma}_t(\hat{X}_t) dW_t, \quad \text{Law}(X_t) = \text{Law}(\hat{X}_t), \quad t \in [0, 1]$$

Proof idea II

Regularize with a Gaussian and make a **Markovian projection**

Mimicking Itô processes [Krylov '85], [Gyöngy '85], [Brunick, Shreve '13]

$$dX_t = \sigma_t dW_t$$

$$d\hat{X}_t = \hat{\sigma}_t(\hat{X}_t) dW_t, \quad \text{Law}(X_t) = \text{Law}(\hat{X}_t), \quad t \in [0, 1]$$

$$\text{Law}(\hat{X}_{t_k}) = \mu_{t_k}^r \quad \text{and} \quad \hat{\sigma} \text{ "nice"}$$

Proof idea III

Use **stability of diffusions** to pass to a limit

Proof idea III

Use **stability of diffusions** to pass to a limit

Theorem 2 [Pammer, R., Schachermayer '23+]

Proof idea III

Use **stability of diffusions** to pass to a limit

Theorem 2 [Pammer, R., Schachermayer '23+]

For $dX_t^k = \sigma_t^k(X_t^k)dB_t$ for “**nice**” σ^k , suppose for each (t, x)

$$\int_0^t \sigma_s^k(x)^2 ds \rightarrow \int_0^t \sigma_s(x)^2 ds.$$

Proof idea III

Use **stability of diffusions** to pass to a limit

Theorem 2 [Pammer, R., Schachermayer '23+]

For $dX_t^k = \sigma_t^k(X_t^k)dB_t$ for “nice” σ^k , suppose for each (t, x)

$$\int_0^t \sigma_s^k(x)^2 ds \rightarrow \int_0^t \sigma_s(x)^2 ds.$$

Then $X^k \rightarrow X$ in f.d.d., $dX_t = \sigma_t(X_t)dB_t$ and σ “nice”.

Main result

Weakly continuous \mathbb{R}^d -valued square-integrable peacock $(\mu_t)_{t \in [0,1]}$.
Regularize with a Gaussian $\mu_t^r := \mu_t * \gamma^{\varepsilon(t+\delta)}$

Theorem 1 [Pammer, R., Schachermayer '23+]

There exists a measurable $(t, x) \mapsto \sigma_t^r(x)$ that is locally Lipschitz in x and non-degenerate, uniformly in $t \in [0, 1]$, and a Brownian motion B such that $\text{Law}(M_t^r) = \mu_t^r$, for all $t \in [0, 1]$, where

$$dM_t^r = \sigma_t^r(M_t^r)dB_t.$$

Moreover:

- M^r is a strong Markov martingale with continuous paths;
- The result **does not hold** without regularization;
- There is no uniqueness for $d \geq 2$.

Counterexamples

Theorem 3 [Pammer, R., Schachermayer '23+]

There exists a weakly continuous square-integrable peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^4 such that, for the peacock $(\mu_t * \gamma^t)_{t \in [0,1]}$, there exists **no mimicking Markov martingale**.

Counterexamples

Theorem 3 [Pammer, R., Schachermayer '23+]


There exists a weakly continuous square-integrable peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^4 such that, for the peacock $(\mu_t * \gamma^t)_{t \in [0,1]}$, there exists no mimicking Markov martingale.

Circular Brownian Motion

[Émery, Schachermayer '99]

[Fernholz, Karatzas, Ruf '18]

[Larsson, Ruf '20]



[Cox, R. '23]

Counterexamples

Theorem 3 [Pammer, R., Schachermayer '23+]

There exists a weakly continuous square-integrable peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^4 such that, for the peacock $(\mu_t * \gamma^t)_{t \in [0,1]}$, there exists no mimicking Markov martingale.

Circular Brownian Motion

[Émery, Schachermayer '99]


[Fernholz, Karatzas, Ruf '18]

[Larsson, Ruf '20]

Theorem [Cox, R. '23]

There is a unique weak solution
but **no strong solution** of

$$dX_t = \frac{1}{|X_t|} \begin{bmatrix} -X_t^2 \\ X_t^1 \end{bmatrix} dW_t, \quad X_0 = 0.$$




[Cox, R. '23]

Counterexamples

Theorem 3 [Pammer, R., Schachermayer '23+]

There exists a weakly continuous square-integrable peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^4 such that, for the peacock $(\mu_t * \gamma^t)_{t \in [0,1]}$, there exists no mimicking Markov martingale.

1. No continuous Markov martingale mimicking μ ;




[Cox, R. '23]

Counterexamples

Theorem 3 [Pammer, R., Schachermayer '23+]

There exists a weakly continuous square-integrable peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^4 such that, for the peacock $(\mu_t * \gamma^t)_{t \in [0,1]}$, there exists no mimicking Markov martingale.

1. No continuous Markov martingale mimicking μ ;
2. No **Markov** martingale mimicking μ ;




[Cox, R. '23]

Counterexamples

Theorem 3 [Pammer, R., Schachermayer '23+]

There exists a weakly continuous square-integrable peacock $(\mu_t)_{t \in [0,1]}$ on \mathbb{R}^4 such that, for the peacock $(\mu_t * \gamma^t)_{t \in [0,1]}$, there exists no mimicking Markov martingale.

1. No continuous Markov martingale mimicking μ ;
2. No Markov martingale mimicking μ ;
3. No **Markov** martingale mimicking $(\mu * \gamma^t)_{t \in [0,1]}$.



[Cox, R. '23]

Main result

Weakly continuous \mathbb{R}^d -valued square-integrable peacock $(\mu_t)_{t \in [0,1]}$.
Regularize with a Gaussian $\mu_t^r := \mu_t * \gamma^{\varepsilon(t+\delta)}$

Theorem 1 [Pammer, R., Schachermayer '23+]

There exists a measurable $(t, x) \mapsto \sigma_t^r(x)$ that is locally Lipschitz in x and non-degenerate, uniformly in $t \in [0, 1]$, and a Brownian motion B such that $\text{Law}(M_t^r) = \mu_t^r$, for all $t \in [0, 1]$, where

$$dM_t^r = \sigma_t^r(M_t^r)dB_t.$$

Moreover:

- M^r is a strong Markov martingale with continuous paths;
- The result does not hold without regularization;
- There is **no uniqueness** for $d \geq 2$.

Faking Brownian motion

Do there exist stochastic processes with **Brownian marginals** that are not Brownian motion?

[Hamza, Klebaner '07]

There exists some **fake Brownian motion**.

Faking Brownian motion

Do there exist stochastic processes with Brownian marginals that are not Brownian motion?

[Hamza, Klebaner '07]

There exists some fake Brownian motion.

[Beiglböck, Lowther, Pammer, Schachermayer '21]

There exists a “very fake” Brownian motion in dimension $d = 1$.

Faking Brownian motion

Do there exist stochastic processes with Brownian marginals that are not Brownian motion?

[Hamza, Klebaner '07]

There exists some fake Brownian motion.

[Beiglböck, Lowther, Pammer, Schachermayer '21]

There exists a **Markov** process with **continuous paths** that mimics Brownian marginals in dimension $d = 1$.

Faking Brownian motion

Do there exist stochastic processes with Brownian marginals that are not Brownian motion?

[Hamza, Klebaner '07]

There exists some fake Brownian motion.

[Beiglböck, Lowther, Pammer, Schachermayer '21]

There exists a in dimension $d = 1$.

Recall: for $d = 1$, a strong Markov process is uniquely determined by its marginals [Lowther '09].

Non-uniqueness


Theorem 4 [Pammer, R., Schachermayer '23+]

There exists an \mathbb{R}^2 -valued strong Markov martingale diffusion with Brownian marginals, which is not a Brownian motion.


Non-uniqueness

Theorem 4 [Pammer, R., Schachermayer '23+]

There exists an \mathbb{R}^2 -valued strong Markov martingale diffusion with Brownian marginals, which is not a Brownian motion.



$$dX_t = |X_t|^{-1} X_t^\top dW_t$$



$$dX_t = |X_t|^{-1} X_t dW_t$$

Solutions of a stochastic control problem for martingales [Cox, R. '23]

Non-uniqueness

Theorem 4 [Pammer, R., Schachermayer '23+]


There exists an \mathbb{R}^2 -valued strong Markov martingale diffusion with Brownian marginals, which is not a Brownian motion.

Theorem [Cox, R. '23]

Let X be a weak solution of

$$dX_t = \frac{1}{|X_t|}(X_t + X_t^\perp)dW_t, \quad X_0 \sim \eta.$$

Then X is a **continuous strong Markov** fake Brownian motion.



[Pammer, R., Schachermayer '23+]

Main result

Weakly continuous \mathbb{R}^d -valued square-integrable peacock $(\mu_t)_{t \in [0,1]}$.
Regularize with a Gaussian $\mu_t^r := \mu_t * \gamma^{\varepsilon(t+\delta)}$

Theorem 1 [Pammer, R., Schachermayer '23+]




There exists a measurable $(t, x) \mapsto \sigma_t^r(x)$ that is locally Lipschitz in x and non-degenerate, uniformly in $t \in [0, 1]$, and a Brownian motion B such that $\text{Law}(M_t^r) = \mu_t^r$, for all $t \in [0, 1]$, where

$$dM_t^r = \sigma_t^r(M_t^r)dB_t.$$

Moreover:

- M^r is a strong Markov martingale with continuous paths;
- The result does not hold without regularization;
- There is no uniqueness for $d \geq 2$.

References

-  Alexander M. G. Cox and Benjamin A. Robinson, *Optimal control of martingales in a radially symmetric environment*, Stochastic Processes and their Applications **159** (2023), 149–198.
-  Alexander M. G. Cox and Benjamin A. Robinson, *SDEs with no strong solution arising from a problem of stochastic control*, Electronic Journal of Probability **28** (2023), 1–24.
-  Gudmund Pammer, Benjamin A. Robinson, and Walter Schachermayer, *A regularized Kellerer theorem in arbitrary dimension*, Annals of Applied Probability, to appear.

Summary

- We prove the first known Kellerer-type result in arbitrary dimension;
- In general, the result can fail without some regularization;
- In dimension $d \geq 2$, uniqueness fails.



G. Pammer, B. R., W. Schachermayer, *A regularized Kellerer theorem in arbitrary dimension*