

Applications of Mathematics in urban rainfall-runoff modelling.

James Fidal

Department of Architecture and Civil Engineering University of Bath

March 1, 2017

Introduction and Background

The world's population is increasing, with projections showing an increase per 1.2% per year with no signs of stopping.[worldbank, 2016].

Introduction and Background

The worlds population is increasing, with projections showing an increase per 1.2% per year with no signs of stopping.[worldbank, 2016].

As of 2014 54% of people are living in urban areas.

Introduction and Background

The worlds population is increasing, with projections showing an increase per 1.2% per year with no signs of stopping.[worldbank, 2016].

As of 2014 54% of people are living in urban areas.

Its debated that land use change such as urbanisation is linked to an increase in flooding events.

Introduction and Background


The worlds population is increasing, with projections showing an increase per 1.2% per year with no signs of stopping.[worldbank, 2016].

As of 2014 54% of people are living in urban areas.

Its debated that land use change such as urbanisation is linked to an increase in flooding events.


A method to capture these effects is rainfall-runoff modelling.

Range of Conceptual rainfall runoff models



A rainfall-runoff model is a mathematical model which describes the relationship between rainfall and river runoff for a catchment.


Range of Conceptual rainfall runoff models



A rainfall-runoff model is a mathematical model which describes the relationship between rainfall and river runoff for a catchment.

There is a wide spectrum of runoff models ranging from extremely detailed to generalised.

Range of Conceptual rainfall runoff models





A rainfall-runoff model is a mathematical model which describes the relationship between rainfall and river runoff for a catchment.


There is a wide spectrum of runoff models ranging from extremely detailed to generalised.

The applications of runoff models ranges from expanding stream climate and flow records, estimating river flow in ungauged catchments and prediction of the effects of land use change.


Difficulties of modelling




Difficulties of modelling




Sealing of previously pervious areas leads to more runoff generated.



Difficulties of modelling




Sealing of previously pervious areas leads to more runoff generated.




The ever changing environment of urban areas, due to construction or expansion etc can lead to conflicts with in modelling.

Difficulties of modelling



Sealing of previously pervious areas leads to more runoff generated.

The ever changing environment of urban areas, due to construction or expansion etc can lead to conflicts with in modelling.



The location of the urbanisation with relation to the river, can impact transportation times.

Develop lumped rainfall-runoff model to better represent urban effects on catchment hydrology; the URMOD model.

Develop lumped rainfall-runoff model to better represent urban effects on catchment hydrology; the URMOD model.

URMOD is a lumped, deterministic, continuous-time rainfall-runoff model.

Develop lumped rainfall-runoff model to better represent urban effects on catchment hydrology; the URMOD model.

URMOD is a lumped, deterministic, continuous-time rainfall-runoff model.

URMOD's urban runoff generation is linked to soil moisture as opposed to the more common approach of fixed percentage or no infiltration.


Develop lumped rainfall-runoff model to better represent urban effects on catchment hydrology; the URMOD model.

URMOD is a lumped, deterministic, continuous-time rainfall-runoff model.


URMOD's urban runoff generation is linked to soil moisture as opposed to the more common approach of fixed percentage or no infiltration.

A nested model approach was used, when no urbanisation is present URMOD defaults to a rural model.

Visual representation of URMOD



URMOD processes



Urban soil and runoff equation

How infiltration is treated

$$f = (1 - u)f_{rur} + uf_{ur}$$

Urban soil and runoff equation

How infiltration is treated

$$f = (1 - u)f_{rur} + uf_{ur}$$

$$f_{ur} = i(1 - \gamma) \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}}$$

$$f_{rur} = i \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}}$$

Urban soil and runoff equation

How infiltration is treated

$$f = (1 - u)f_{rur} + uf_{ur}$$

$$f_{ur} = i(1 - \gamma) \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}}$$

$$f_{rur} = i \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}}$$

$$f = (1 - u)i \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}} + ui(1 - \gamma) \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}}.$$

Urban soil and runoff equation

How infiltration is treated

$$f = (1 - u)f_{rur} + uf_{ur}$$

$$f_{ur} = i(1 - \gamma) \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}}$$

$$f_{rur} = i \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}}$$


$$f = (1 - u)i \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}} + ui(1 - \gamma) \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}}.$$

The urban moisture equation

$$\frac{dm}{dt} = \underbrace{\left((1 - u)i \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}} + ui(1 - \gamma) \left(1 - \frac{m}{SM}\right)^{\frac{1}{2}} \right)}_{\text{Infiltration}} - \underbrace{D(m - F)}_{\text{Drainage}} - \underbrace{E_p}_{\text{Evaporation}}$$


Gamma parameter

Infiltration vs Soil saturation



This is with a fixed scaling term γ . In the above example $\gamma = 0.7$

Case Study- Thames River



112 catchment's subsetting into 29 catchments in the Thames river.

Case Study- Thames River

Comparison of performance between URMOD and the default rural model.

Case Study- Thames River

Comparison of performance between URMOD and the default rural model.

Urbanisation levels from 8% up to 55%. Areas ranging from 8.835Km² up to 690.575 Km². Varying soil types from impervious (London clay) to pervious (chalk).

Case Study- Thames River

Comparison of performance between URMOD and the default rural model.

Urbanisation levels from 8% up to 55%. Areas ranging from 8.835Km² up to 690.575 Km². Varying soil types from impervious (London clay) to pervious (chalk).

The level of urbanisation criteria is URBEXT₂₀₀₀, which is a contribution of the urban and sub-urban areas of a catchment [Bayliss et al., 2006]

Case Study- Thames River

Comparison of performance between URMOD and the default rural model.

Urbanisation levels from 8% up to 55%. Areas ranging from 8.835Km² up to 690.575 Km². Varying soil types from impervious (London clay) to pervious (chalk).

The level of urbanisation criteria is URBEXT₂₀₀₀, which is a contribution of the urban and sub-urban areas of a catchment [Bayliss et al., 2006]

Rainfall data was available from 1961 to 2012 from the Gridded estimates of daily and monthly areal rainfall (GEAR).

Case Study- Thames River

Comparison of performance between URMOD and the default rural model.


Urbanisation levels from 8% up to 55%. Areas ranging from 8.835Km² up to 690.575 Km². Varying soil types from impervious (London clay) to pervious (chalk).

The level of urbanisation criteria is URBEXT₂₀₀₀, which is a contribution of the urban and sub-urban areas of a catchment [Bayliss et al., 2006]

Rainfall data was available from 1961 to 2012 from the Gridded estimates of daily and monthly areal rainfall (GEAR).

River runoff data used was in a 30 year period from 1980-2010.

Calibration and Validation period



The calibration period for the model was the black years, where as the Validation periods is the yellow years .

The Nash-Sutcliffe efficiency Statistic E

$$E = 1 - \frac{\sum_{n=1}^n (Q_{obs} - Q_{sim})^2}{\sum_{n=1}^n (Q_{obs} - \bar{Q}_m)^2}$$

E is bounded between 1 and $-\infty$, so will be transformed.

The Nash-Sutcliffe efficiency Statistic E

$$E = 1 - \frac{\sum_{n=1}^n (Q_{obs} - Q_{sim})^2}{\sum_{n=1}^n (Q_{obs} - \bar{Q}_m)^2}$$

E is bounded between 1 and $-\infty$, so will be transformed.

A *tanh* transformation was applied to the E statistic. Too constrict the lower bound of the E , bounds now are $(-1, 0.762)$.

Jack-knife experiment

The Jack-knife mean is obtained for each catchment.

$$\bar{Z}_N = N^{-1} \sum_{j=1}^N Z_{N-1,j}. \quad (1)$$

Jack-knife experiment

The Jack-knife mean is obtained for each catchment.

$$\bar{Z}_N = N^{-1} \sum_{j=1}^N Z_{N-1,j}. \quad (1)$$

The associated jackknife variance estimate can also be calculated, $\text{var}(Z_N) = V_N$.

Jack-knife experiment

The Jack-knife mean is obtained for each catchment.

$$\bar{Z}_N = N^{-1} \sum_{j=1}^N Z_{N-1,j}. \quad (1)$$


The associated jackknife variance estimate can also be calculated, $\text{var}(Z_N) = V_N$.

$$V_N = \frac{N-1}{N} \sum_{j=1}^N (Z_{N-1,j} - \bar{Z}_N)^2. \quad (2)$$

The 95% confidence intervals are calculated.

$$(\bar{Z}_N - 2 * \sqrt{V_N}, \bar{Z}_N + 2 * \sqrt{V_N}) \quad (3)$$

Comparison of urban and rural model



Comparison between Urban model and default rural. Circles indicate Urban model better performance and triangles indicate default rural model better performance.

Conclusion

Presented a new framework for generalised rainfall-runoff modelling in urbanised catchments.

Conclusion

Presented a new framework for generalised rainfall-runoff modelling in urbanised catchments.

Modelling urban catchments is difficult due to human influence.

Conclusion

Presented a new framework for generalised rainfall-runoff modelling in urbanised catchments.

Modelling urban catchments is difficult due to human influence.

Applications of this model, being used in thames water project.

Conclusion

Presented a new framework for generalised rainfall-runoff modelling in urbanised catchments.

Modelling urban catchments is difficult due to human influence.

Applications of this model, being used in thames water project.

Is URMOD a improvment ?

Further Reading



Bayliss, A., Black, K., Fava-Verde, A., and Kjeldsen, T. (2006).
URBEXT2000-a new FEH catchment descriptor. calculation,
dissemination and application.



worldbank (2016).
Population growth (annual %).