

On the constant terms of certain meromorphic modular forms

Barry Brent

draft 23h 29 May 2023

Abstract

In this report of computer experiments we study the divisibility properties of the constant terms of certain meromorphic modular forms for Hecke groups and relate those properties to several sequences, for example, to O.E.I.S. sequence A005148 [28], which was studied by Newman, Shanks and Zagier [27], [44], and several other sequences, the members of which appear in congruences of Ramanujan. At the end of the article, we construct from elementary arithmetic functions some meromorphic but not necessarily modular functions and study their constant terms.

1 Introduction

1.1 On constant terms.

Let $f(x) = 1/x + \sum_{n=0}^{\infty} a_{n+1}x^n$ (*sic.*) For $g(x)$ a Laurent series in x , let $C(g)$ denote its constant term. In this article, we study $C(f^k)$, mostly in settings where (after substituting one of several exponential functions for x) f is a meromorphic modular form for some matrix group.

The constant $C(f^k)$ is a function of the coefficients a_1, \dots, a_k . Furthermore the numbers $C(f), C(f^2), \dots$ determine a_1, a_2, \dots . To see this, let c_k be the coefficient of x^k in the polynomial $h(x) = (1 + \sum_{n=1}^k a_n x^n)^k$. It is clear that $c_k = C(f^k)$. We have $c_1 = a_1$, $c_2 = a_1^2 + 2a_2$, $c_3 = a_1^3 + 6a_1a_2 + 3a_3$, *etc.*

The occasion for our interest was a problem in the theory of quadratic forms, which led us to the empirical finding that equations (1) and (2) below and corresponding equations for other meromorphic modular forms are valid for $k \leq 50$ [9]. Here we test (1), (2) and several analogues for $k \leq 5000$.

The function $\Delta(z)$ occurring in equation (2) is defined as follows. For z in the upper half plane and $q = q(z) = \exp(2\pi iz)$, $\Delta(z)$ is the weight twelve normalized cusp form for $SL(2, \mathbb{Z})$ with Fourier expansion $\sum_{n=1}^{\infty} \tau(n)q^n$, where τ denotes Ramanujan's function. The reciprocal $1/\Delta(z)$ appears in expressions

for the dimensions of certain Lie algebras ([15], page 328; [16], page 45.) It also appears in string theory, for example, in the counting of black hole microstates ([26], equation (14).) We will study the constant terms $C(1/\Delta^k)$ (k a positive integer).

The Klein invariant appearing in equation (1), $j(z) = 1/q + \sum_{n=0}^{\infty} c(n)q^n$, defined on the upper half of the complex plane with $c(0) = C(j) = 744$, is central (for example) to the classical theory of modular forms and to the moonshine phenomenon. We will also study the constant terms $C(j^k)$.

Constant terms of meromorphic modular forms came into our work on quadratic forms as follows. Siegel studied the constant terms in the Fourier expansions of a particular family of meromorphic modular forms T_h for $SL(2, \mathbb{Z})$ (“level one modular forms”) in 1969 [34, 35]. Siegel demonstrated that these constant terms never vanish. He used this to establish a bound on the exponent of the first non-vanishing Fourier coefficient for a level one entire modular form f of weight h such that the constant term of f is itself non-vanishing. Theta functions fit this description, so Siegel was able to give an upper bound on the least positive integer represented by a positive-definite even unimodular quadratic form in $2h$ variables. While working on an extension of Siegel’s result on the non-vanishing of the T_h constant terms we came across the regularities described in equations (1) and (2).

One question we study is the nature of the special role of the primes $p = 2$ and 3 in equations (1) and (2): why these primes but not others (apparently)? We searched for regularities involving other primes among the modular forms for Hecke groups (section 7.) Along the way we made observations relevant to the classical situation as well (conjecture 3.)

1.2 The structure of constant terms.

Constant terms of meromorphic modular forms of certain kinds appear to have multiplicative structure. While seeking a level two version of Siegel’s result, the present writer found numerical evidence for divisibility properties of the constant terms for several kinds of modular form, including the T_h [9]; if these properties hold, the constant terms cannot vanish.¹ Let $d_b(n)$ be the sum of the digits in the base b expansion of n . Then (apparently)

$$\text{ord}_2(C(j^k)) = \text{ord}_2(C(1/\Delta^k)) = 3d_2(k) \quad (1)$$

and

$$\text{ord}_3(C(j^k)) = \text{ord}_3(C(1/\Delta^k)) = d_3(k). \quad (2)$$

We argue (based on numerical experiments) that the $C(j^k)$ inherit the stated properties from the OEIS sequence A005148 [28], which was originally studied

¹For example, see Serre [33], section 3.3, equation (22), or the Wikipedia page [42].

by Newman, Shanks and Zagier [27, 44] in an article on its use in series approximations to π .

We tried to find patterns in the p -orders of constant terms of j and other modular forms for $SL(2, \mathbb{Z})$ for p larger than three. Our search within $SL(2, \mathbb{Z})$ seemed to fail, so we searched among the Hecke groups $G(\lambda_n), n = 3, 4, \dots$. The matrix group $SL(2, \mathbb{Z})$ coincides with the Hecke group $G(\lambda_3)$, discussed below. It is isomorphic to the product of cyclic groups $C_2 * C_3$; while in general $G(\lambda_m) \cong C_2 * C_m$ for $m = 3, 4, \dots$. We will state some conjectures about the constant terms, for example, of meromorphic forms for Hecke groups isomorphic to $C_2 * C_{p^k}, p$ prime.

Recently we found apparent regularities for $p = 5, 7, 11$ in the original case of $SL(2, \mathbb{Z})$ (conjectures 2 and 13.) They are conditions equivalent to the statement that $\text{ord}_p(C(f))$ vanishes (for $p = 5, 7, 11$ when $f = j^k$, and for $p = 5$ and 7 when $f = 1/\Delta^k$.) These conditions are simple restrictions on the digits in the base p expansions of k . The author's thesis advisor² remarked that (1) and (2) might follow from congruences of Ramanujan. We report experiments that support this suggestion in the last section.

The present article states several conjectures based on extensive computations (mainly done with *SageMath*), but no theorems. The data is available in a GitHub repository [7].

2 Background

2.1 Ramanujan's congruences.

Here are the congruences of Ramanujan mentioned above ([38], page 290 and elsewhere.)^{3 4}

$$\tau(n) \equiv \sigma_{11}(n) \pmod{2^8} \tag{3}$$

for odd n .

$$\tau(n) \equiv n\sigma_1(n) \pmod{3}. \tag{4}$$

$$\tau(n) \equiv n^2\sigma_1(n) \pmod{3^2}. \tag{5}$$

$$\tau(n) \equiv n^2\sigma_7(n) \pmod{3^3}. \tag{6}$$

$$\tau(n) \equiv n\sigma_1(n) \pmod{5}. \tag{7}$$

$$\tau(n) \equiv n\sigma_9(n) \pmod{5^2}. \tag{8}$$

²Glenn Stevens

³The following congruences appear in Ramanujan's unpublished (before Berndt and Ono did publish it) manuscript [4]: (4) is Ramanujan's (11.8), (5) is Ramanujan's (12.3), (6) is also Ramanujan's (12.3), (7) is Ramanujan's (2.1), (8) is Ramanujan's (4.2), and (10) is Ramanujan's (12.7).

⁴It is well known that they have been strengthened; see the articles [4], [38], [39], [40], [31], [30], [43], [22], [21], and [2].

$$\tau(n) \equiv n\sigma_3(n) \pmod{7}. \quad (9)$$

$$\tau(n) \equiv \sigma_{11}(n) \pmod{691}. \quad (10)$$

$$\tau_r(n) \equiv n\sigma_{2r-1}(n) \pmod{11} \quad (11)$$

for $r = 2, 3$ and 4 .⁵

Remark 1. Equation (3) extends to all of the positive integers as follows: let $o = \text{ord}_2(n)$ and $g(n) = 8^o \cdot \sigma_{11}(n/2^o)$. Then

$$\tau(n) \equiv g(n) \pmod{2^8}.$$

To see this, recall Ramanujan's conjecture (proved by Mordell [25]) that, for $n \geq 1$ and p prime: $\tau(p)\tau(p^n) = \tau(p^{n+1}) + p^{11}\tau(p^{n-1})$.⁶ Setting $p = 2$, an easy induction argument shows that $\text{ord}_2(\tau(2^o)) = 3o$, and the claim follows from the multiplicativity of $\tau(n)$.

2.2 Modular forms for Hecke groups.

For $m = 3, 4, \dots$, let $\lambda_m = 2\cos \pi/m$ and let J_m be a certain meromorphic modular form for the Hecke group $G(\lambda_m)$, built from triangle functions, with Fourier expansion

$$J_m(z) = \sum_{n=-1}^{\infty} a_n(m) q_m^n,$$

where $q_m(z) = \exp 2\pi iz/\lambda_m$. (For further details, the reader is referred to the books by Carathéodory [13, 14] and by Berndt and Knopp [3], the articles of Lehner and Raleigh [23, 29], to the dissertation of Leo [24], and to a summary, including pertinent references to that material, in the 2021 article [8].)

Raleigh gave polynomials $P_n(x)$ such that $a_{-1}(m)^n q_m^{2n+2} a_n(m) = P_n(m)$ for $n = -1, 0, 1, 2$ and 3 . He conjectured that similar relations hold for all positive integers n [29].⁷ Akiyama proved Raleigh's conjectures in 1992 [1].

Using the weight-raising properties of differentiation and the J_m , Hecke constructed families \mathcal{H} comprising modular forms of positive weight for each $G(\lambda_m)$ sharing certain properties [18, 3]. It seems apparent that Akiyama's result can be extended: there should exist polynomials $Q_{\mathcal{H},n}(x)$ interpolating the coefficient of X_m^n in the Fourier expansions of the members of Hecke families \mathcal{H} .⁸

⁵The congruences in (11) are displayed in the table at the top of page SwD-32 (page 32 of the proceedings [39]), and, in the form shown here, as equation (13) on page Ran-6 (page 8 of the proceedings [32].)

⁶See equation (53) of proposition 14 in section 5.5 of Serre's book [33].

⁷For more on expansions over polynomial fields, see, for example, the book of Boas and Buck [5] and the articles by Buckholtz and Byrd ([11], [12].)

⁸See the paper [8].

In section 4 of the 2021 article, we made use of a certain uniformizing variable $X_m(z)$ for z in the upper half plane [8]. By Akiyama's theorem, we have a series of the form $\mathcal{J}_m(x) := \sum_{n=-1}^{\infty} \tilde{P}_n(x) X_m^n$ for polynomials $\tilde{P}_n(x)$ in $\mathbb{Q}[x]$ with the property that $J_m = \mathcal{J}_m(m)$. We will make use of the change of variables $X_m \mapsto 2^6 m^3 X_m$ for a $G(\lambda_m)$ -modular form (originally employed, apparently, by Leo ([24], page 31). It has the effect when $m = 3$ of recovering the Fourier series of a variety of standard modular forms. This is set up as a

Definition 1. For z in the upper half plane and $k_a \neq 0$, let

$$f(z) := \sum_{n=a}^{\infty} k_n X_m(z)^n$$

and

$$g(z) = \sum_{n=a}^{\infty} k_n 2^{6n} m^{3n} X_m(z)^n.$$

If the last expansion is written as $g(z) = \sum_{n=a}^{\infty} \tilde{k}_n X_m(z)^n$, then let

$$\bar{f}(z) := g(z)/\tilde{k}_a.$$

Also, for $m = 3, 4, \dots$, let $j_m(z) := \overline{J_m}(z)$.

The Fourier expansion of j_3 is ⁹

$$j_3(z) = 1/X_3(z) + 744 + 196884X_3(z) + 21493760X_3(z)^2 + \dots,$$

which matches the standard expansion $j(z) =$

$$1/\exp(2\pi iz) + 744 + 196884 \exp(2\pi i \cdot z) + 21493760 \exp(2\pi i \cdot 2 \cdot z) + \dots$$

Definition 2. Let $\mathcal{F} = \{f_3, \dots, f_m, \dots\}$ where f_m is modular for $G(\lambda_m)$. Then let the Fourier expansion of f_m^k in powers of X_m be written

$$f_m(z)^k = \sum_n a(f_m^k, n) X_m^n.$$

(Thus $C(f_m^k) = a(f_m^k, 0)$.)

Proposition 1. Let $\mathcal{K} = \{J_3, J_4, \dots\}$ and $\bar{\mathcal{K}} = \{j_3, j_4, \dots\}$. Then there exist polynomials $Q_{\mathcal{K},k,n}(x)$ and $Q_{\bar{\mathcal{K}},k,n}(x)$ in $\mathbb{Q}[x]$ such that $a(J_m^k, n) = Q_{\mathcal{K},k,n}(m)$ and $a(j_m^k, n) = Q_{\bar{\mathcal{K}},k,n}(m)$ for $k = 1, 2, \dots, m = 3, 4, \dots$, and $n = -k, 1 - k, \dots$

For k equal to one, the first claim is just Akiyama's theorem and the claim for k not equal to one is then obvious. The second statement follows immediately.

⁹See equation (23) of Serre's book [33], section 3, and the *SageMath* notebook "jpower constant term NewmanShanks 26oct22.ipynb" in [7].

2.3 Polynomial interpolation of Fourier coefficients.

When, given a sequence of functions f_m modular for $G(\lambda_m)$ in a family \mathcal{F} , we looked for polynomials $Q_{\mathcal{F},n}(x)$ such that each f_m with Fourier expansion

$$f_m(\tau) = \sum_n a_{m,n} X_m^n(\tau),$$

satisfied $Q_{\mathcal{F},n}(m) = a_{m,n}$. We evaluated finite sequences $\{a_{m,n}\}_{m=1,2,3,4,\dots,M_n}$ (with n held constant) and generated candidates $g_n(x)$ for $Q_{\mathcal{F},n}(x)$ by Lagrange interpolation. The bounds M_n were linear in n and chosen large enough that the degrees of the $g_n(x)$ produced in this way also appeared to be linear in n . Over the course of experiments described in the article [8], this linearity seemed to be associated with systematic behavior. For example, if a polynomial $g_n(x)$ was factored as $g_n(x) = r_n \cdot p_1(x) \cdot p_2(x) \dots \cdot p_a(x)$ where each of the p_i was monic, r_n was rational, and the degree of $g_n(x)$ was linear in n , then often the sequence $\{r_3, r_4, \dots\}$ was readily identifiable (sometimes after resorting to Sloane's encyclopedia.) We take such regularities as evidence that $g_n(m) = a_{m,n}$ for all m . Thus, when formulating conjectures about the $C(J_m^k)$ and $C(j_m^k)$ ¹⁰, we did not always use tables of the $C(J_m^k)$ and $C(j_m^k)$ directly. Instead (for example), we used Lagrange interpolation to identify polynomials $h_k(x)$ and $\bar{h}_k(x)$ such that $C(J_m^k) = h_k(m)$ and $C(j_m^k) = \bar{h}_k(m)$ by letting m run through a small set of values sufficient to produce the linearity behavior mentioned above; so we assumed (in this example) that $h_k(x) \equiv Q_{\mathcal{K},k,0}(x)$ and $\bar{h}_k(x) \equiv Q_{\bar{\mathcal{K}},k,0}(x)$ identically. We made tables of p orders of the $h_k(m)$ and the $\bar{h}_k(m)$. In this way we checked larger sets of m values than would have been practicable if we had checked the constant terms themselves.

Unlike the later conjectures, conjecture 1 is not a way of summarizing patterns in experimental data. Rather it codifies our assumption that the linearity behavior is a reliable signal.

Conjecture 1. 1. $h_k(x) \equiv Q_{\mathcal{K},k,0}(x)$ identically; consequently, $h_k(m) = C(J_m^k)$ identically.

2. $\bar{h}_k(x) \equiv Q_{\bar{\mathcal{K}},k,0}(x)$ identically; consequently, $\bar{h}_k(m) = C(j_m^k)$ identically.

3 The reciprocals of cusp forms for $SL(2, \mathbb{Z})$

Let E_{2r} denote the weight $2r$ Eisenstein series with q -series

$$1 + \gamma_r \sum_{r=1}^{\infty} \sigma_{r-1}(n) q^n$$

for certain rational numbers γ_r ; this is Rankin's notation. In our experiments, including the case $r = 1$, which is not in Rankin's list, we rely on *SageMath* to

¹⁰See the *SageMath* notebooks in the repository [7], in the folder "conjectures".

pick out the unique normalized cusp form of weight $12 + 2r$, so there is no need to specify γ_r by hand. Recall several facts:¹¹ Setting $E_0(z) = 1$, $\tau_0(n) = \tau(n)$, and $r = 0, 2, 3, 4, 5$ or 7 :

1. $\Delta(z)E_{2r}(z)$ generates the space of weight $12 + 2r$ cusp forms for $SL(2, \mathbb{Z})$.
2. Writing $\Delta_r = \Delta(z)E_{2r}(z)$ and $\Delta_r = \sum_{n=1}^{\infty} \tau_r(n)q^n$: the functions $n \mapsto \tau_r(n)$ are multiplicative.

Conjecture 2. *Suppressing the dependence upon k and r , let $d_p = d_p(k)$, $C = C(1/\Delta_r^k)$ and $o_p = \text{ord}_p(C)$.*

1. Let $r = 0$.

- (a) $o_2 = 3d_2$ and $o_3 = d_3$.
- (b) $C/3^{o_3} \equiv 1 \pmod{3}$ if and only if k is even.
- (c) $C/3^{o_3} \equiv 2 \pmod{3}$ if and only if k is odd.
- (d) i. $C \equiv 0, 1$, or $4 \pmod{5}$.
ii. $o_5 = 0$ if and only if the set of digits in the base 5 expansion of k is a subset of $\{0, 1, 2\}$.¹²
- (e) $o_7 = 0$ if and only if the set of digits in the base 7 expansion of k is a subset of $\{0, 1, 2, 3, 4\}$.

2. Let $r = 2$.¹³

- (a) $o_2 = 3d_2$.
- (b) i. $o_3 = d_3$ if and only if $k \equiv 0 \pmod{3}$.
ii. If D is a positive integer such that $D \equiv 2 \pmod{3}$, $k > D$, and, for some positive n , $k = D + 3^n$, then o_3 is constant for large n . Let $L_D = \lim_{n \rightarrow \infty} o_3$ and N_D be the smallest value of n such that $n \geq N_D \Rightarrow o_3 = L_D$. Below is a table for small D . More extensive tables are posted on GitHub [7].

D	2	5	8	11	14	17	20	23
L_D	4	5	8	5	6	8	7	8
N_D	1	2	3	3	3	3	4	3

- (c) If k is even and $k \equiv 0 \pmod{3}$, then $C/3^{o_3} \equiv 1 \pmod{3}$.
- (d) If k is odd and $k \equiv 0 \pmod{3}$, then $C/3^{o_3} \equiv 2 \pmod{3}$.
- (e) $o_5 = 0$ if and only if the set of digits in the base 5 expansion of k is a subset of $\{0, 1, 2\}$.

3. Let $r = 3$.¹⁴

¹¹See page ran-4 (page six in the proceedings volume) of Rankin's article [32].

¹²See O.E.I.S. page [19].

¹³The converses of clauses (c) and (d) are false.

¹⁴Again, the converses of clauses (c) and (d) are false.

- (a) $o_2 = 3d_2$ if and only if k is even.
- (b)
 - i. $o_3 = d_3$.
 - ii. $C/3^{o_3} \equiv 1 \pmod{3}$ if and only if k is even.
 - iii. $C/3^{o_3} \equiv 2 \pmod{3}$ if and only if k is odd.
- (c) If $o_5 = 0$, then the set of digits in the base 5 expansion of k is a subset of $\{0, 1, 2, 3\}$.
- (d) If $o_7 = 0$, then the set of digits in the base 7 expansion of k is a subset of $\{0, 1, 2, 3, 4\}$.

4. Let $r = 4$.

- (a) For all positive k , $o_2 = 3d_2$.
- (b)
 - i. For all positive k , $C \equiv 0 \pmod{3}$.
 - ii. If $k \equiv 0 \pmod{3}$, then $o_3 = d_3$.
 - iii. If $k \equiv 1 \pmod{3}$, then $o_3 = d_3$ if and only if k belongs to O.E.I.S sequence A191107 [20]¹⁵ $\{1, 4, 10, \dots\}$,
 - iv. If $k \equiv 2 \pmod{3}$ and d_3 divides o_3 , then $o_3/d_3 = 2$.
- (c)
 - i. $c \equiv 0, 1$, or $4 \pmod{5}$.
 - ii. $o_5 = 0$ if and only if the digits in the base 5 expansion of k is a subset of $\{0, 1, 2\}$.
 - iii. If $o_5 = 0$, then $C/5^{o_5} \equiv 1$ or $4 \pmod{5}$.¹⁶

5. Let $r = 5$.

- (a) If k is even, then $o_2 = 3d_2$.
- (b) If D is a positive odd integer, $k > D$ and $k = D + 2^n$, then o_2 is constant for large n . Let $L_D = \lim_{n \rightarrow \infty} o_2$ and N_D be the smallest value of n such that $n \geq N_D \Rightarrow o_2 = L$. Below is a table for small D . More extensive tables are posted on GitHub [7].

D	1	3	5	7	9	11	13	15	17	19
L_D	10	13	17	19	15	17	23	27	17	17
N_D	3	3	6	5	6	5	8	9	6	6

Remark 2. 1. A p -adic geometric view of conjectures 2.2.b (ii) and 2.5.b is that the function $k \mapsto C(1/\Delta^k)$ takes certain units k in sufficiently small disks around certain other units to circles around zero.

2. Conjectures 2.2.b (ii) and 2.5.b have only limited empirical support because the mentioned p -adic units k grow exponentially with n and, on account of drastic slowdowns for large k , our experiments tested only $k \leq 5000$. Thus for $p = 2$ and 3, we could only check $n \leq 12$ and 7, respectively. We will include tables of what empirical data we do have in the appendix.

¹⁵Description: "Increasing sequence generated by these rules: $a(1) = 1$, and if x is in a then $3x - 2$ and $3x + 1$ are in a ." *Mathematica* code: `h = 3; i = -2; j = 3; k = 1; f = 1; g = 7; a = Union[Flatten[NestList[{h # + i, j # + k} &, f, g]]]`.

¹⁶The converse is false.

4 Constant terms for $j^k, k = 1, 2, \dots$

Recall that $h_k(x)$ and $\bar{h}_k(x)$ are polynomials identified from numerical data by Lagrange interpolation conjectured to satisfy $C(J_m^k) = h_k(m)$ and $C(j_m^k) = \bar{h}_k(m)$. In this section, we illustrate connections between the divisibility patterns (described in the introduction) for the constant terms of the $j(\tau)^k = j_3(\tau)^k$ Fourier expansions on one side, and the $h_k(x)$ on the other. Let $\bar{h}_k(x)$ factor as $\bar{h}_k(x) = \nu_k \cdot p_{k,1}(x) \times p_{k,2}(x) \times \dots \times p_{k,\alpha}(x) = (\text{say}) \nu_k \cdot \tilde{p}_k(x)$ where each of the $p_{k,n} (n = 1, 2, \dots, \alpha)$ is monic and ν_k is rational. We represent O.E.I.S. sequence A005148 [28] $\{0, 1, 47, 2488, 138799, \dots\}$ as $\{a_0, a_1, \dots\}$.

Conjecture 3. 1. $\nu_k = 24a_k$.

2. $\tilde{p}_k(3)$ is always odd.

3. $\text{ord}_2(a_k) = 3d_2(k) - 3$.

4. $\text{ord}_3(\tilde{p}_k(3)) = d_3(k) - 1$.

5. From the introduction: $\text{ord}_2(C(j_3^k)) = 3d_2(k)$ and $\text{ord}_3(C(j_3^k)) = d_3(k)$.

6. We restate another observation from the article [9]. Let $o_k = \text{ord}_3(C(j_3^k)), \kappa = C(j_3^k)/3^{o_k}$, and $\rho_k = \text{mod}(\kappa, 3)$. Then $\rho_k = 1$ or 2 , according as k is even or odd, respectively.

7. (a) Let $p = 5$ or 7 and let $o = \text{ord}_p(C(j_3^k))$. Then $o = 0$ if and only if the set of digits in the base p expansion of k is a subset of $\{0, 1, 2\}$.

(b) Let $p = 11$. With notation as above, $o = 0$ if and only if the set of digits in the base p expansion of k is a subset of $\{0, 1, 2, 3, 4\}$.

Remark 3. Clause 5 of the conjecture follows from the earlier clauses. First claim: $\text{ord}_2(C(j_3^k)) = \text{ord}_2(\bar{h}_k(3)) = \text{ord}_2(\nu_k \cdot \tilde{p}_k(3)) = \text{ord}_2(24a_k) \cdot \tilde{p}_k(3) = \text{ord}_2(24) + \text{ord}_2(a_k) + \text{ord}_2(\tilde{p}_k(3)) = 3 + 3d_2(k) - 3 + 0 = 3d_2(k)$. Second claim: In their 1984 article [27], Newman, Shanks and Zagier demonstrated that $\text{ord}_3(a_k) = 0$ for all k . Therefore (under the previous clauses) $\text{ord}_3(C(j_3^k)) = \text{ord}_3(\bar{h}_k(3)) = \text{ord}_3(\nu_k) + \text{ord}_3(\tilde{p}_k(3)) = 1 + \text{ord}_3(a_k) + d_3(k) - 1 = d_3(k)$.

5 Sufficient conditions for equations (1), (2)

We construct some Laurent series (not necessarily modular, even after an appropriate substitution) such that their constant terms satisfy analogues of equation (1) or equation (2). Some conjectures in this section were tested with Monte Carlo methods.

Conjecture 4. ¹⁷

¹⁷See the folder “conjectures” in the repository [7].

1. Let $A_n = \text{lcm}(\{2 \cdot 8^{d_2(k)}\}_{k=1, \dots, n+1})$. If

$$f(x) = \sum_{k=1}^{\infty} a_k x^k$$

is in $\mathbb{Z}[[x]]$, $a_k \equiv \tau(k) \pmod{A_n}$ for $k = 1, 2, \dots, n+1$, then

$$\text{ord}_2(C(1/f(x)^n)) = 3d_2(n).$$

2. Let $B_n = \text{lcm}(\{3 \cdot 3^{d_3(k)}\}_{k=1, \dots, n+1})$. If

$$f(x) = \sum_{k=1}^{\infty} a_k x^k$$

is in $\mathbb{Z}[[x]]$, $a_k \equiv \tau(k) \pmod{B_n}$ for $k = 1, 2, \dots, n+1$, then

$$\text{ord}_3(C(1/f(x)^n)) = d_3(n).$$

3. Let $C_n = \text{lcm}(\{6 \cdot 8^{d_2(k)} \cdot 3^{d_3(k)}\}_{k=1, \dots, n+1})$. If

$$f(x) = \sum_{k=1}^{\infty} a_k x^k$$

is in $\mathbb{Z}[[x]]$, $a_k \equiv \tau(k) \pmod{C_n}$ for $k = 1, 2, \dots, n+1$, then

$$\text{ord}_2(C(1/f(x)^n)) = 3d_2(n)$$

and

$$\text{ord}_3(C(1/f(x)^n)) = d_3(n).$$

In the following conjectures, analogues to the series expansion of $\Delta(z)$ from the right sides of Ramanujan's congruences (3) – (11) are constructed. Graphical tests indicate that they are not modular forms, but they each appear to have some of the behaviors conjectured for Δ .

Conjecture 5. 1. Let $o_k = \text{ord}_2(k)$, $g_k = 8^{o_k} \cdot \sigma_{11}(k/2^{o_k})$, and

$$f(x) = \sum_{k=1}^{\infty} g_k x^k.$$

Then

$$(a) \text{ord}_2(C(1/f(x)^n)) = 3d_2(n).$$

$$(b) C(1/f(x)^n) \equiv 1 \pmod{3}.$$

2. Let A_n be as in the previous conjecture, g_k be as above, and let

$$f(x) = \sum_{k=1}^{\infty} a_k x^k,$$

where $a_k \equiv g_k \pmod{A_n}$. Then $\text{ord}_2(C(1/f(x)^n)) = 3d_2(n)$.

Conjecture 6. Let $o_2 = \text{ord}_2(k)$, $o_3 = \text{ord}_3(k)$, $g_k = k \cdot \sigma_1(k)$, and

$$f(x) = \sum_{k=1}^{\infty} g_k x^k.$$

1. If n is divisible by 4, then $\text{ord}_2(C(1/f(x)^n)) = 3d_2(n)$.
2. If n is divisible by 3, then $\text{ord}_3(C(1/f(x)^n)) = d_3(n)$.
3. If $n - 1$ is divisible by 3 and $n - 2$ is a power of 3 or twice a power of 3, then once again $\text{ord}_3(C(1/f(x)^n)) = d_3(n)$.¹⁸

Conjecture 7. Let $g_k = k^2 \cdot \sigma_1(k)$ and

$$f(x) = \sum_{k=1}^{\infty} g_k x^k.$$

1. If n is even, then $\text{ord}_2(C(1/f(x)^n)) = 3d_2(n)$.
2. For $n = 1, 2, \dots$, $\text{ord}_3(C(1/f(x)^n)) = d_3(n)$.

6 Powers of reciprocals of generating functions of certain other arithmetic functions

The functions studied in this section are constructed from certain multiplicative or additive (in the sense that $f(ab) = f(a) + f(b)$ when $\gcd(a, b) = 1$) arithmetic functions. They are not necessarily modular or consistent with analogues of equations (1) and (2).

Conjecture 8. Let

$$f_r(x) = \sum_{k=1}^{\infty} \sigma_r(k) x^k.$$

1. (a) $C(1/f_0(x)^n)$ is odd if and only if n is divisible by three.
(b) blah
2. (a) For all positive integers n , $C(1/f_1(x)^n)$ is odd.
(b) Blah
3. blah

In the following conjecture we study divisor sums with multiplicity.

Definition 3. 1. For $n = \prod_i p_i^{n_i}$, $d = \prod_i p_i^{d_i}$ with $0 \leq d_i \leq n_i$, and $\binom{a}{b}$ the usual binomial coefficient, the multiplicity of d in n is

$$\mu(d, n) := \prod_{d_i | n_i} \binom{n_i}{d_i}.$$

¹⁸For this sequence, see the O.E.I.S. page [10] of K. Brockhaus.

2.

$$\sigma_r^\mu(n) := \sum_{d|n} \mu(d, n) d^r.$$

Conjecture 9. ¹⁹ *Let*

$$f_r^\mu(x) = \sum_{k=1}^{\infty} \sigma_r^\mu(k) x^k$$

and $C_{r,n} = C(1/f_r^\mu(x)^n)$.

1. (a) $C_{r,n}$ is odd for all positive integers r and n .
(b) If r is odd and n is even, then $C_{r,n} \equiv 1 \pmod{3}$.
2. $C_{0,n}$ is even for all positive integers n .
3. (a) $C_{1,n}$ is odd for all positive integers n .
(b) $C_{1,n} \equiv 0 \pmod{3}$ if and only if n is even.
4. *blah*

7 Constant terms for $j_m^k, k = 1, 2, \dots$

By imposing restrictions on k and m , we found several narrow conjectures about constant term p orders for various primes p .

7.1 m a prime power.

20

Conjecture 10. *If p is prime and a is an integer that is larger than 2, then*

$$\text{ord}_p(C(j_{p^a}^k)) = (a-3)k + \text{ord}_p(C(j_{p^3}^k)).$$

Conjecture 11. *Let $a \geq 2$. Then $\text{ord}_2(C(j_{2^a}^2)) = 2a + 7$.*

Conjecture 12. *Let p be a prime number larger than 2 and let a be a positive integer. Then $\text{ord}_p(C(j_{p^a}^p)) = ap - 2$.*

¹⁹Clause 1 is based on substantially less data than the clauses that specify particular values of r .

²⁰Again, see the *SageMath* notebooks in the folder “conjectures” in the repository [7]. Also see O.E.I.S. pages [41],[17], [36],[45].

7.2 Other m .

Conjecture 13. If $d_2(k) = 1$, $a = \text{ord}_2(m)$, $a \geq 2$, and $o = \text{ord}_2(C(j_m^k))$, then $o = k(a + 2) + 3$.

Conjecture 14. Let $d_2(k) = 1$, $m \equiv 2 \pmod{4}$, and $a = \text{ord}_2(m) (= 1, \text{ of course.})$ Then $\text{ord}_2(C(j_m^k)) = k(a + 6) + 1 = 7k + 1$.

Now let $K_n, n = 0, 1, 2, \dots$ be the n^{th} Catalan number. (We depart from the standard notation because we have been using the letter “c” in so many other contexts.) One of several explicit formulas for K_n is

$$K_n = \frac{(2n)!}{(n+1)!n!}.$$

For n positive let $K_{1,n}$ denote the n^{th} Catalan number K such that $K \neq K_0$ and $\text{ord}_2(K) = 1$.²¹

Conjecture 15. Let k be the n^{th} positive integer such that $d_2(k) = 2$; also, $m = 4j$, ($j = 1, 2, \dots$), and $a = \text{ord}_2(m)$. Furthermore, let $o = \text{ord}_2(C(j_m^k))$ and $t = ((a + 6)k + 2 - o)/4$. Then $t = K_{1,n}$.

Conjecture 16. Let $d_2(k) = 2$, $m = 4j + 2$, $j = 1, 2, \dots$, and $a = \text{ord}_2(m)$ (again, $a = 1$.) Then $\text{ord}_2(C(j_m^k)) = (a + 6)k + 2 = 7k + 2$.

Conjecture 17. If $m \equiv 0 \pmod{3}$, then $\text{ord}_3(C(j_m^k)) = k \cdot \text{ord}_3(m) + d_3(k) - k$.

7.3 The constant terms $c(J_m^k)$.

The Fourier coefficients of the J_m are rational numbers, but typically they are not integers.

Conjecture 18.²² Let p be a prime number greater than two and let $c(J_p^p) = a/b$ (a, b relatively prime integers, b positive.) Then $b = 2^{6p-3d_2(p)}p^{2p+2}$.

References

- [1] S. Akiyama. “A note on Hecke’s absolute invariants”. In: *J. Ramanujan Math. Soc* 7.1 (1992), pp. 65–81.
- [2] M. H. Ashworth. “Congruence and identical properties of modular forms”. Ph.D. thesis supervised by A. O. L. Atkin, cited in [39]. University of Oxford, 1968.
- [3] B. C. Berndt and M. I. Knopp. *Hecke’s theory of modular forms and Dirichlet series*. Vol. 5. World Scientific, 2008.

²¹See Bottomley’s O.E.I.S. page [6].

²²See [37] and other O.E.I.S. pages cited within it.

- [4] B. C. Berndt and K. Ono. “Ramanujan’s unpublished manuscript on the partition and tau functions with proofs and commentary”. In: *Sém. Lotharingien de Combinatoire* 42 (1999), p. 63.
- [5] R. P. Boas and R. C. Buck. *Polynomial expansions of analytic functions*. Vol. 19. Springer Science & Business Media, 2013.
- [6] H. Bottomley. *The On-Line Encyclopedia of Integer Sequences*, A099628. <http://oeis.org/A099628>.
- [7] B. Brent. *GitHub files for this article*. <https://github.com/barry314159a/NewmanShanks>. 2023.
- [8] B. Brent. *Polynomial interpolation of modular forms for Hecke groups*. <http://math.colgate.edu/~integers/v118/v118.pdf>.
- [9] B. Brent. “Quadratic minima and modular forms”. In: *Experimental Mathematics* 7.3 (1998), pp. 257–274.
- [10] K. Brockhaus. *The On-Line Encyclopedia of Integer Sequences*, A164123. <http://oeis.org/A164123>.
- [11] J. D. Buckholtz. “Series expansions of analytic functions”. In: *Journal of Mathematical Analysis and Applications* 41.3 (1973), pp. 673–684.
- [12] P. F. Byrd. “Expansion of analytic functions in polynomials associated with Fibonacci numbers”. In: *Fibonacci Q.* 1 (1963), p. 16.
- [13] C. Carathéodory. *Theory of functions of a complex variable, Second English Edition*. Vol. 1. Translated by F. Steinhardt. Chelsea Publishing Company, 1958.
- [14] C. Carathéodory. *Theory of functions of a complex variable, Second English Edition*. Vol. 2. Translated by F. Steinhardt. Chelsea Publishing Company, 1981.
- [15] I. B. Frenkel. “Representations of Kac-Moody algebras and dual resonance models”. In: *Applications of group theory in physics and mathematical physics* 21 (1985), pp. 325–354.
- [16] I. B. Frenkel, J. Lepowsky, and A. Meurman. *Vertex operator algebras and the Monster*. Academic press, 1989.
- [17] J.-S. Gerasimov. *The On-Line Encyclopedia of Integer Sequences*, A176003. <http://oeis.org/A176003>.
- [18] E. Hecke. “Über die bestimmung dirichletscher reihen durch ihre funktionalgleichung”. In: *Mathematische Annalen* 112.1 (1936), pp. 664–699.
- [19] C. Kimberling. *The On-Line Encyclopedia of Integer Sequences*, A037453. <http://oeis.org/A037453>.
- [20] C. Kimberling. *The On-Line Encyclopedia of Integer Sequences*, A191107. <http://oeis.org/A191107>.
- [21] O. Kolberg. *Congruences for Ramanujan’s Function $\tau(n)$* . Norwegian Universities Press, 1962.

- [22] D. H. Lehmer. *Note on some arithmetical properties of elliptic modular functions*. Duplicated notes, University of California at Berkeley, cited in [39].
- [23] J. Lehner. “Note on the Schwarz triangle functions.” In: *Pacific Journal of Mathematics* 4.2 (1954), pp. 243–249.
- [24] J. G. Leo. *Fourier coefficients of triangle functions*, *Ph.D. thesis*. <http://halfaya.org/ucla/research/thesis.pdf>. 2008.
- [25] L. J. Mordell. “Note on certain modular relations considered by Messrs. Ramanujan, Darling, and Rogers”. In: *Proceedings of the London Mathematical Society* 2.1 (1922), pp. 408–416.
- [26] S. Murthy. *Black holes and modular forms in string theory*. <https://arxiv.org/pdf/2305.11732.pdf>. 2023.
- [27] M. Newman and D. Shanks. “On a Sequence Arising in Series for π ”. In: *Pi: A Source Book*. Springer, 2004, pp. 462–480.
- [28] S. Plouffe and N. J. A. Sloane. *The On-Line Encyclopedia of Integer Sequences*, A005148. <http://oeis.org/A005148>.
- [29] J. Raleigh. “On the Fourier coefficients of triangle functions”. In: *Acta Arithmetica* 8 (1962), pp. 107–111.
- [30] S. Ramanujan. “On certain arithmetical functions”. In: *Trans. Cambridge Philos. Soc* 22.9 (1916), pp. 159–184.
- [31] S. Ramanujan. “On certain arithmetical functions”. In: *Collected papers of Srinivasa Ramanujan*. Cambridge University Press, 2015, pp. 136–162.
- [32] R. A. Rankin. “Ramanujan’s unpublished work on congruences”. In: *Modular Functions of one Variable V: Proceedings International Conference, University of Bonn, Sonderforschungsbereich Theoretische Mathematik July 2–14, 1976*. Springer. 2006, pp. 3–15.
- [33] J.-P. Serre. *A course in arithmetic*. Springer-Verlag, 1970.
- [34] C. L. Siegel. “Berechnung von Zetafunktionen an ganzzahligen Stellen”. In: *Akad. Wiss.* 10 (1969), pp. 87–102.
- [35] C. L. Siegel. “Evaluation of zeta functions for integral values of arguments”. In: *Advanced Analytic Number Theory, Tata Institute of Fundamental Research, Bombay* 9 (1980), pp. 249–268.
- [36] N. J. A. Sloane. *The On-Line Encyclopedia of Integer Sequences*, A049001. <http://oeis.org/A049001>.
- [37] N. J. A. Sloane and A. Wilks. *The On-Line Encyclopedia of Integer Sequences*, A005187. <http://oeis.org/A005187>.
- [38] H. P. F. Swinnerton-Dyer. “Congruence properties of $\tau(n)$ ”. In: *Ramanujan revisited: proceedings of the [Ramanujan] Centenary Conference, University of Illinois at Urbana-Champaign, June 1-5, 1987*. Harcourt Brace Jovanovich, 1988, pp. 289–311.

- [39] H. P. F. Swinnerton-Dyer. “On l -adic representations and congruences for coefficients of modular forms”. In: *Modular Functions of One Variable III: Proceedings International Summer School University of Antwerp, RUCA July 17–August 3, 1972*. Springer. 1973, pp. 1–55.
- [40] H. P. F. Swinnerton-Dyer. “On l -adic representations and congruences for coefficients of modular forms (II)”. In: *Modular Functions of one Variable V: Proceedings International Conference, University of Bonn, Sonderforschungsbereich Theoretische Mathematik July 2–14, 1976*. Springer. 2006, pp. 63–90.
- [41] A. Turpel. *The On-Line Encyclopedia of Integer Sequences*, A037168. <http://oeis.org/A037168>.
- [42] Wikipedia. *j-invariant*. <https://en.wikipedia.org/wiki/J-invariant>. 2022.
- [43] J. R. Wilton. “Congruence properties of Ramanujan’s function $\tau(n)$ ”. In: *Proceedings of the London Mathematical Society* 2.1 (1930), pp. 1–10.
- [44] D. Zagier. “Appendix to ‘On a Sequence Arising in Series for π ’ by Newman and Shanks”. In: *Pi: A Source Book*. Springer, 2004, pp. 462–480.
- [45] R. Zumkeller. *The On-Line Encyclopedia of Integer Sequences*, A084920. <http://oeis.org/A084920>. 2013.

barrybrent@iphouse.com