

Polynomial interpolation of modular forms for Hecke groups

Barry Brent

20h 15 May 2021

Abstract

For $m = 3, 4, \dots$, let $\lambda_m = 2 \cos \pi/m$ and let $J_m(m = 3, 4, \dots)$ be triangle functions for the Hecke groups $G(\lambda_m)$ with Fourier expansions $J_m(\tau) = \sum_{n=-1}^{\infty} a_n(m) q_m^n$, where $q_m(\tau) = \exp 2\pi i \tau / \lambda_m$. (When normalized appropriately, J_3 becomes Klein's j -invariant $j(\tau) = 1/e^{2\pi i \tau} + 744 + \dots$) For $n = -1, 0, 1, 2$ and 3 , Raleigh [25] gave polynomials $P_n(x)$ such that $a_{-1}(m)^n q_m^{2n+2} a_n(m) = P_n(m)$ for $m = 3, 4, \dots$, and conjectured that similar relations hold for all positive integers n . Akiyama proved this conjecture in [1]. We apply work of Hecke [4] to study experimentally similar polynomial interpolations of the J_m Fourier coefficients and the Fourier coefficients of other, positive weight, modular forms for $G(\lambda_m)$. We connect these polynomials (again, only empirically) with variants of Dedekind's eta function, with the Fourier expansions of some standard Hauptmoduln, and, in the case of analogues of Eisenstein series for $SL(2, \mathbb{Z})$, with certain divisor sums.

1 Introduction

Let $\mathbb{Z}, \mathbb{Q}, \mathbb{C}$ and \mathbb{H} denote, respectively, the set of rational integers, the set of rational numbers, the set of complex numbers, and the set of complex numbers with positive imaginary parts. (We will reserve the letter τ for elements of the upper half-plane, and z for generic complex numbers.) We write $\mathbb{H}^* = \mathbb{H} \cup \mathbb{Q} \cup \{i\infty\}$, and we equip \mathbb{H}^* with the Poincaré metric. Figures T made by three geodesics of \mathbb{H}^* are called hyperbolic or circular-arc triangles. Let $\lambda_m = 2 \cos \pi/m$. For $m = 3, 4, \dots$, we define the Hecke group $G(\lambda_m)$ as the discrete group generated by the maps $z \rightarrow -1/z$ and $z \rightarrow z + \lambda_m$. The full modular group $SL(2, \mathbb{Z})$ is identical to $G(\lambda_3)$.

To define modular forms for the Hecke groups, we preview a definition from Berndt [4], which we will quote again in a later section. (We depart occasionally from Berndt's choices of variable to avoid clashes with some of our other notation.)

We say that f belongs to the space $M(\lambda, k, \gamma)$ if

1.

$$f(\tau) = \sum_{n=0}^{\infty} a_n e^{2\pi i n \tau / \lambda},$$

where $\lambda > 0$ and $\tau \in \mathbb{H}$, and

2. $f(-1/\tau) = \gamma(\tau/i)^k f(\tau)$, where $k > 0$ and $\gamma = \pm 1$.

We say that f belongs to the space $M_0(\lambda, k, \gamma)$ if f satisfies conditions 1 and 2 and if $a_n = O(n^c)$ for some real number c , as n tends to ∞ .

Members of $M(\lambda, k, \gamma)$ are known as modular forms for $G(\lambda)$ of weight k . Condition 1 tells us that they are invariant under translations $\tau \mapsto \tau + \lambda$. Next we preview Berndt's definition of cusp forms for Hecke groups. If $f \in M(\lambda, k, \gamma)$ and $f(i\infty) = 0$, then we call f a cusp form of weight k and multiplier γ with respect to $G(\lambda)$. For cusp forms, the constant terms of condition 1 vanish. We denote by $C(\lambda, k, \gamma)$ the vector space of all cusp forms of this kind.

For our purposes, Schwarz triangles T are hyperbolic triangles in \mathbb{H}^* with certain restrictions on the angles at the vertices. From a Euclidean point of view, their sides are vertical rays, segments of vertical rays, semicircles orthogonal to the real axis and meeting it at points $(r, 0)$ with r rational, or arcs of such semicircles. We choose λ, μ and ν , all non-negative, such that $\lambda + \mu + \nu < 1$; then the angles of T are $\lambda\pi, \mu\pi$, and $\nu\pi$. By reflecting T across one of its edges, we get another Schwarz triangle. The reflection between two triangles in \mathbb{H}^* is effected by a Möbius transformation, so the orbit of T under repeated reflections is associated to a collection of Möbius transformations. The group generated by these transformations is a triangle group. By the Riemann Mapping Theorem there is a conformal, onto map $\phi : T \mapsto \mathbb{H}^*$ called a triangle function.

Hecke groups [17] are triangle groups H that act properly discontinuously on \mathbb{H} . This means that for compact $K \subset \mathbb{H}$, the set $\{\mu \in H \text{ s.t. } K \cap \mu(K) \neq \emptyset\}$ is finite. Recall that $G(\lambda_m)$ is the Hecke group generated by the maps $z \mapsto -1/z$ and $z \mapsto z + \lambda_m$. Apparently it was Hecke who, also in [17], established that $G(\lambda_m)$ has the structure of a free product of cyclic groups $C_2 * C_m$, generalizing the relation [29, 8] $SL(2, \mathbb{Z}) = C_2 * C_3$.

Let $\rho = -\exp(-\pi i/m) = -\cos(\pi/m) + i \sin(\pi/m)$, and let $T_m \subset \mathbb{H}^*$ denote the hyperbolic triangle with vertices ρ, i , and $i\infty$. The corresponding angles are $\pi/m, \pi/2$ and 0 respectively. Let ϕ_{λ_m} be a triangle function for T_m . The function ϕ_{λ_m} has a pole at $i\infty$ and period λ_m . For $P, Q \in \mathbb{H}^*$, let us write $P \equiv_H Q$ when $\mu \in H$ and $Q = \mu(P)$. Then ϕ_{λ_m} extends to a function $J_m : \mathbb{H}^* \rightarrow \mathbb{H}^*$ by declaring that $J_m(P) = J_m(Q)$ if and only if $P \equiv_H Q$. J_m is a modular function for $G(\lambda_m)$.

Schwarz [28], Lehner [20], Raleigh [25] and others studied Schwarz triangle functions, which map hyperbolic triangles T in the extended upper half z -plane

onto the extended upper half w -plane. For certain $T = T_m$, a triangle function $\phi_{\lambda_m} : T \rightarrow \mathbb{H}^*$ extends to a map $J_m : \mathbb{H}^* \rightarrow \mathbb{H}^*$ invariant under modular transformations from $G(\lambda_m)$. Suitably normalized, the J_m become analogues j_m of the normalized Klein’s modular invariant

$$j(\tau) = -1/q + 744 + 196884q + \dots$$

where $q = q(\tau) = \exp(2\pi i\tau)$ and $j_3(\tau) = j(\tau)$.

With $\lambda_m = 2 \cos \pi/m$ and $q_m(\tau) = \exp(2\pi i\tau/\lambda_m)$, the original J_m have Fourier series $J_m(\tau) = \sum_{n \geq -1} a_n(m) q_m(\tau)^n$. For $n = -1, 0, 1, 2$ and 3 , Raleigh [25] gave polynomials $P_n(x)$ such that $a_{-1}(m)^n q_m^{2n+2} a_n(m) = P_n(m)$ for $m = 3, 4, \dots$, and conjectured that similar relations hold for all positive integers n . Akiyama [1] proved Raleigh’s conjecture in the passage after his (Akiyama’s) equation (6).

Hecke ([4], [17]) built families of modular forms f_m for $G(\lambda_m)$ sharing particular properties, such as their weight; we will describe his construction later. Earlier authors, whose work we will also describe, had already built modular functions (meromorphic functions invariant under the action of $G(\lambda)$, thus, of weight zero) from triangle functions.

The plan of the article is as follows. We sketch the theory of Schwartz triangles; then, the construction of their triangle functions; from the triangle functions, the modular functions for the Hecke groups; and, from them, modular forms for these groups. To this point, the article is just a summary of background material. By methods familiar from the classical case, we then construct modular forms with rational Fourier expansions, describe our modifications of them, and describe computer experiments on the modifications.¹

The experiments indicate that the Fourier expansions of several Hecke modular forms are interpolated in the sense of Raleigh by polynomials. We describe connections among these polynomials, certain divisor sums, and Hauptmoduln discussed in the Moonshine literature. We offer only conjectures; we prove no theorems of our own. All theorems that we mention are part of the existing literature.

The earliest computer code we located for calculating Fourier expansions of triangle functions for Hecke groups is that of Leo; it is based on Lehner’s construction. Leo also calculates the Fourier coefficients of weight 4 and weight 6

¹We documented our experiments in a series of *Mathematica* notebooks and accompanying data files. That material is in a *Github* repository named “interpolations” [5]. A green “Code” button on the top page of the repository contains a drop-down menu with a download option. We tried to make the remarks, clauses of conjectures, *etc.*, findable by reading the file names. Data files (in plain text format) are named less transparently, but they are defined within the documents where they were generated. A *Mathematica* notebook (“mf23.nb”) in the repository contains a searchable library of functions that may not be defined explicitly within notebooks containing code that calls them.

Hecke-analogues of classical Eisenstein series in Chapter 4 of [21]. Our code for triangle functions, which is based on Leo's, comes from the papers of Lehner and Raleigh. J. Jermann's package [18] is also concerned with modular forms of triangle groups for Hecke groups, but we did not use his code in our experiments. For definiteness, we transcribe our code defining triangle functions and modular forms in an appendix.

2 A glossary

Several special functions in this list are so closely related that one might wish to know why we have included all of them. It is for the convenience of both the reader and ourselves, because differing forms (basically just different notations) are used by Lehner and Raleigh, and we hope in this way to discuss both of them while avoiding confusion.

1. The digamma function $\psi(z) := \Gamma'(z)/\Gamma(z)$.
2. The Schwarzian derivative ([10], p. 130, equation 370.8)

$$\{w, z\} = \frac{2w'w''' - 3w''^2}{2w'^2} \quad (1)$$

for $w = w(z)$.

3. The Pochhammer symbol

$$(a)^0 := 1 \text{ and, for } n \geq 1, (a)^n := a(a+1)\dots(a+n-1) = \Gamma(a+n)/\Gamma(a).$$

4. The function c_ν given by ([10], p. 138, equation 377.3)

$$c_\nu = c_\nu(\alpha, \beta, \gamma) := \frac{(\alpha)^\nu (\beta)^\nu}{\nu! (\gamma)^\nu}, \nu \geq 0.$$

To facilitate comparison with Raleigh's equation (9¹) [25], we remark that

$$c_\nu = \frac{\Gamma(\alpha + \nu)}{\Gamma(\alpha)} \cdot \frac{\Gamma(\beta + \nu)}{\Gamma(\beta)} \cdot \frac{\Gamma(1)}{\Gamma(1 + \nu)} \cdot \frac{\Gamma(\gamma)}{\Gamma(\gamma + \nu)}. \quad (2)$$

In the terms of this article's Theorem 1 below, Raleigh is treating the case $\lambda = 0$, for which (equation (6) below) $\gamma = 1$ and the expression on the right side of (2) becomes, as in Raleigh,

$$\frac{\Gamma(\alpha + \nu)\Gamma(\beta + \nu)}{\Gamma(\alpha)\Gamma(\beta)(\nu!)^2}.$$

5. The function e_ν given by ([25], equation 9¹)

$$e_\nu = e_\nu(\alpha, \beta) := \sum_{p=0}^{\nu-1} \left(\frac{1}{\alpha + p} + \frac{1}{\beta + p} - \frac{2}{1 + p} \right).$$

Here, we are dealing with the same ambiguity present in the definition of c_ν : this is a specialization to the case $\gamma = 1$ of the e_ν for $\nu \geq 1$ given by ([10], p. 153, equation 387.5)

$$e_\nu = e_\nu(\alpha, \beta, \gamma) := \sum_{p=0}^{n-1} \left(\frac{1}{\alpha+p} + \frac{1}{\beta+p} - \frac{2}{\gamma+p} \right).$$

Unless it is explicitly indicated to be otherwise, we intend the former (Raleigh's) definition.

6. (a) Gauss's hypergeometric series given by [10], p. 138, equation (377.4),

$$F(\alpha, \beta, \gamma; z) := \sum_{\nu=0}^{\infty} c_\nu(\alpha, \beta, \gamma) z^\nu$$

F is occasionally written in [10] as ϕ_1 (for example, on, p. 152.)

- (b) As defined in the first line of [10], p. 142:

$$F_1(\alpha, \beta, \gamma; z) := F(\alpha, \beta, \gamma + 1; z).$$

- (c) Alternatively, dropping γ , as defined in [21], equation (3.5):

$$F_1(\alpha, \beta; z) := \sum_{\nu=1}^{\infty} \frac{(\alpha)_k (\beta)_\nu}{(\nu!)^2} e_\nu(\alpha, \beta).$$

It is in the latter form, defined more cryptically in [20], p. 244, that we will use F_1 ; to establish his series for the triangle functions, which we will apply below, Lehner uses this definition of F_1 , as well as certain theorems from Fricke [14]. Referring to item 4, we see that

$$F_1(\alpha, \beta; z) = \sum_{\nu=1}^{\infty} c_\nu(\alpha, \beta, 1) e_\nu(\alpha, \beta).$$

We will derive another form of $F_1(\alpha, \beta; z)$ in item 7.

7. With $F = F(\alpha, \beta, \gamma; z)$, a special function

$$F^*(\alpha, \beta, \gamma; z) := \frac{\partial F}{\partial \alpha} + \frac{\partial F}{\partial \beta} + 2 \frac{\partial F}{\partial \gamma};$$

by [10], equation (387.4) on p. 153, F^* may be written

$$F^*(\alpha, \beta, \gamma; z) = \sum_{\nu=1}^{\infty} c_\nu(\alpha, \beta, \gamma) e_\nu(\alpha, \beta, \gamma) z^\nu.$$

It follows that $F^*(\alpha, \beta, 1; z) = F_1(\alpha, \beta; z)$.

8. A special function $\phi_2^*(z)$ is defined as a certain limit ([10], p. 152, equation 386.2) but is immediately (equation 386.3) reduced to

$$\phi_2^*(z) = F(\alpha, \beta, 1; z) \log z + F^*(\alpha, \beta, 1; z).$$

9. The set $\mathcal{Q} = \{2, 5, 6, 8, 10, 11, 14, 15, 17, 18, 20, 22, 23, \dots\}$ of positive integers not represented by the quadratic form $x^2 + xy + y^2$ [30]. B. Cloitre asserts, on the same O.E.I.S. page, that \mathcal{Q} is also the set of non-negative integers n such that $\delta(n)$ is non-zero, where η is Dedekind's eta function and $\sum_n \delta(n)x^n = \eta(x^3)/\eta(x)^3$.
10. The McKay-Thompson series of class 4A, $\{1, 24, 276, 2048, \dots\}$, which is the sequence of coefficients in the q -series of a certain hauptmodul discussed in [22]. McKay relates the hauptmodul treated in that paper to Dedekind's eta function. We identified the class 4A McKay Thompson series with the \bar{a}_n of our conjecture 1 after finding it in the *On-line Encyclopedia of Integer Sequences* [35].

3 Calculation of Schwarz's inverse triangle function

Schwarz proved

Theorem 1. ([10], §374)

1. Let the half-plane $\Im z > 0$ be mapped conformally onto an arbitrary circular-arc triangle whose angles at its vertices A, B , and C are $\pi\lambda, \pi\mu$, and $\pi\nu$, and let the vertices A, B, C be the images of the points $z = 0, 1, \infty$, respectively. Then the mapping function $w(z)$ must be a solution of the third-order differential equation

$$\{w, z\} = \frac{1 - \lambda^2}{2z^2} + \frac{1 - \mu^2}{2(1 - z^2)} + \frac{1 - \lambda^2 - \mu^2 + \nu^2}{2z(1 - z)}. \quad (3)$$

2. If $w_0(z)$ is any solution of equation (3) that satisfies $w'_0(z) \neq 0$ at all interior points of the half-plane, then the function

$$w(z) = \frac{aw_0(z) + b}{cw_0(z) + d} \quad (ad - bc \neq 0)$$

is likewise a solution of equation 3.

3. Also, every solution of equation (3) that is regular and non-constant in the half-plane $\Im z > 0$ represents a mapping of this half-plane onto a circular-arc triangle with angles $\pi\lambda, \pi\mu$, and $\pi\nu$.

In Carathéodory's lexicon ([9] p. 124), a regular function is one that is differentiable on an open connected set. Carathéodory writes the left side of (3) as " $\{w, z\} = \frac{w' w''' - 3w''^2}{w'^2} = \dots$ ", but this is not the case. We infer that the Schwarzian derivative $\{w, z\}$ is intended from the automorphy property of clause 2.

Let us write

$$\alpha = \frac{1}{2}(1 - \lambda - \mu + \nu), \quad (4)$$

$$\beta = \frac{1}{2}(1 - \lambda - \mu - \nu), \quad (5)$$

and

$$\gamma = 1 - \lambda. \quad (6)$$

The solutions w of (3) are inverse to triangle functions; they are quotients of arbitrary solutions of

$$u'' + p(z)u' + q(z)u = 0 \quad (7)$$

when ([10], p. 136, equation (376.4))

$$p = \frac{1 - \lambda}{z} - \frac{1 - \mu}{1 - z}$$

and

$$q = -\frac{\alpha\beta}{z(1 - z)}.$$

Equation (7) reduces ([10], p. 137, equations 376.5-7) to the hypergeometric differential equation

$$z(1 - z)u'' + (\gamma - (\alpha + \beta + 1)z)u' - \alpha\beta u = 0. \quad (8)$$

As long as γ is not a non-positive integer, $u = F(\alpha, \beta, \gamma; z)$ is a solution of (8); it is the only solution regular at $z = 0$, and it satisfies $F(\alpha, \beta, \gamma; 0) = 1$ (final paragraph of [10], §377, p. 138.)

In [10], §§386-388 (pp. 151-155), we find that when $\gamma = 1$ and $\lambda = 0$, another, linearly independent, solution of equation (7) is $\phi_2^*(z)$. [10], Section 394, pp. 165 - 167 is devoted to the case $\lambda = 0$. The mapping function w of Theorem 1 satisfies ([10], p. 166, equation 394.4)

$$w = \frac{1}{\pi i} \left[\frac{\phi_2^*}{\phi_1} - (2\psi(1) - \psi(1 - \alpha) - \psi(1 - \beta)) \right] + i \frac{\sin \pi \mu}{\cos \pi \mu + \cos \pi \nu}. \quad (9)$$

4 Inversion of Schwarz's inverse triangle function

Following Lehner and Raleigh, we consider the Schwarz triangle T_m with vertices at $\rho = -\exp(-\pi i/m)$, i , and $i\infty$. In terms of Theorem 1, T_m has $\lambda = 0$

(an angle 0 at the vertex $i\infty$), $\mu = 1/2$ (an angle $\pi/2$ at i), and $\nu = 1/m$ (an angle π/m at ρ .) In this situation, $\gamma = 1$.

Let J_m be automorphic for $G(\lambda_m)$ with $J_m(\rho) = 0$, $J_m(i) = 1$, and $J_m(i\infty) = \infty$. In terms of Theorem 1, w and J_m are inverse functions. We are going to write down the Fourier expansion $\sum_{n=-1}^{\infty} a_n q_m(\tau)^n$ of J_m .

By clause 2 of Theorem 1, if w satisfies equations (3) and (9), so does $\tau = \tau(z) = \lambda_m w(z)/2$, and therefore

$$2\pi i\tau/\lambda_m = \frac{\phi_2^*}{\phi_1} - (2\psi(1) - \psi(1-\alpha) - \psi(1-\beta)) - \pi \sec(\pi/m).$$

Let us write $\log A_m = -2\psi(1) + \psi(1-\alpha) + \psi(1-\beta) - \pi \sec(\pi/m)$. In general, $A_m = a_{-1}(m)$ ([25], for example two lines below equation (12).) Recalling the definitions of ϕ_1 and ϕ_2^* from our glossary items 6 and 8, we find (abbreviating $J_m(\tau)$ as J_m) that

$$2\pi i\tau/\lambda_m = -\log J_m + \frac{F^*(\alpha, \beta, 1; 1/J_m)}{F(\alpha, \beta, 1; 1/J_m)} + \log A_m. \quad (10)$$

Equation (10) is equation (6) of [25], but Raleigh suppresses the subscripts. He also writes $\exp 2\pi i\tau/\lambda_m$ as x_m , so that (in our earlier notation) $x_m = q_m(\tau)$.

In Raleigh's notation, after taking exponentials,

$$x_m/A_m = \frac{1}{J_m} \exp \frac{F^*(\alpha, \beta, 1; 1/J_m)}{F(\alpha, \beta, 1; 1/J_m)}, \quad (11)$$

the right side of which has a power series in J_m with rational coefficients. Writing $X_m = x_m/A_m$ we can regard $X_m = X_m(J_m)$ as a power series in J_m with rational coefficients. Following [20] and [25], we inverted this power series to obtain one for the modular function J_m , also with rational coefficients. Let \mathcal{J} be a formal operation taking a power series $\sigma(v)$ to its inverse; that is, if $u = \sigma(v)$ then $v = \mathcal{J}(\sigma)(u)$. Let $Y_m(J)$ be a power series such that

$$Y_m(J_m) = J_m \exp \frac{F^*(\alpha, \beta, 1; J_m)}{F(\alpha, \beta, 1; J_m)} = X_m(1/J_m)$$

and hence

$$Y_m(1/J_m) = \frac{1}{J_m} \exp_m \frac{F^*(\alpha, \beta, 1; 1/J_m)}{F(\alpha, \beta, 1; 1/J_m)} = X_m(J_m),$$

so that $\mathcal{J}(Y_m)(X_m(J)) = 1/J_m$ and, therefore, $J_m = 1/\mathcal{J}(Y_m)(X_m)$.

Remark 1. We noticed several typos in [25]. Four of Raleigh's equations—(I), (10), and the two equations on p. 109 that begin “ $a_{-1}(q) = \dots$ ” (where Raleigh's q is our m)—are pairwise contradictory. From the second paragraph

on Raleigh's p. 110, we expect that $A_3 = a_{-1}(3) = 1/1728$, $A_4 = a_{-1}(4) = 1/256$, and $A_6 = a_{-1}(6) = 1/108$. These values are consistent with Raleigh's equation (10), but not with the others. We infer that all of them except (10) are incorrect. Thus, following Raleigh by writing ψ for the digamma function, $\alpha(m)$ for $(1/2 - 1/m)/2$, and $\beta(m)$ for $(1/2 + 1/m)/2$,

$$a_{-1}(m) = \exp(-2\psi(1) + \psi(1 - \alpha(m)) + \psi(1 - \beta(m)) - \pi \sec(\pi/m)). \quad (12)$$

5 Modular forms from modular functions

When the w -image of \mathbb{H}^* is T_m , the inverse of w is ϕ_{λ_m} . J_m , the extension by modularity of ϕ_{λ_m} to \mathbb{H}^* , is periodic with period λ_m and maps ρ to 0, i to 1, and $i\infty$ to ∞ ([20], equation (2).) These mapping properties allow us, following Berndt's exposition [4] of Hecke, to construct positive weight modular forms for $G(\lambda_m)$ from J_m . This section describes results of Hecke that are perhaps most easily accessible for the classical case $m = 3$ in Schoeneberg [27] and, for the general case, in Berndt [4].

5.1 The case $m = 3$.

By keeping track of the weights, zeros and poles of the constituent factors in the numerator and denominator of the fraction defining

$$f_{a,b,c} = \frac{J'^a}{J^b(J-1)^c},$$

Schoeneberg [27] (Theorem 16, p.45) demonstrates that $f_{a,b,c}$ is an entire modular form of weight $2a$ for $SL(2, \mathbb{Z})$ if $a \geq 2$, $3c \leq a$, $3b \leq 2a$, $b + c \geq a$ and a, b, c are integers. (Schoeneberg speaks of "dimension $-2a$.".) Thus he is able to write down a weight 4 entire modular form $E_4^* = f_{2,1,1}$ for $SL(2, \mathbb{Z})$ with a zero of order $\frac{1}{3}$ at $\rho = e^{2\pi i/3}$ and a weight 6 entire modular form $E_6^* = f_{3,2,1}$ for $SL(2, \mathbb{Z})$ with a zero of order $\frac{1}{2}$ at i . (Schoeneberg writes G_4^*, G_6^* .) It is well known that the (vector space) dimension of the spaces of weight 4 and 6 entire modular forms for $SL(2, \mathbb{Z})$ is equal to one, so E_4^* and E_6^* may be identified with the usual weight 4 and weight 6 Eisenstein series, up to a normalization. Finally, Schoeneberg defines the weight 12 cusp form $\Delta^* = E_4^{*3} - E_6^{*2}$ with a zero of order 1 at $i\infty$. It is a multiple of Δ .

5.2 The case $m \geq 3$.

We quote statements from Berndt [4], which is an exposition of Hecke's [16] and other writings. We depart occasionally from Berndt's choices of variable to avoid clashes with our earlier notation.

Definition 1. ([4], Definition 2.2) *We say that f belongs to the space $M(\lambda, k, \gamma)$ if*

1.

$$f(\tau) = \sum_{n=0}^{\infty} a_n e^{2\pi i n \tau / \lambda},$$

where $\lambda > 0$ and $\tau \in \mathbb{H}$, and

2. $f(-1/\tau) = \gamma \cdot (\tau/i)^k f(\tau)$, where $k > 0$ and $\gamma = \pm 1$.

We say that f belongs to the space $M_0(\lambda, k, \gamma)$ if f satisfies conditions 1 and 2, and if $a_n = O(n^c)$ for some real number c , as n tends to ∞ .

After defining the notion of a fundamental region in the usual way and defining as $G(\lambda)$ the group of linear fractional transformations generated by $\tau \mapsto -1/\tau$ and $\tau \mapsto \tau + \lambda$, Berndt states (for $\tau = x + iy$)

Theorem 2. ([4], Theorem 3.1) Let $B(\lambda) = \{\tau \in \mathbb{H} : x < \lambda/2, |\tau| > 1\}$. Then if $\lambda \geq 2$ or if $\lambda = 2 \cos(\pi/m)$, where $m \geq 3$ is an integer, $B(\lambda)$ is a fundamental region for $G(\lambda)$.

Definition 2. ([4], Definition 3.4) Let $T_A = \{\lambda : \lambda = 2 \cos(\pi/m), m \geq 3, m \in \mathbb{Z}\}$.

Berndt states in his Theorem 5.4 that $G(\lambda)$ is discrete if and only if λ belongs to T_A . This discreteness is the premise of the theory of automorphic functions generally. He embeds within the proof of his Lemma 3.1 (which we omit), the

Definition 3. τ_λ denotes the intersection in \mathbb{H} of the line $x = -\lambda/2$ and the unit circle $|\tau| = 1$.

(Berndt remarks at the top of page 35 that τ_λ is the lower left corner of $B(\lambda)$. and that $\pi\theta = \pi - \arg(\tau_\lambda)$, so that $\cos(\pi\theta) = \lambda/2$.)

To characterize Eisenstein series, we need to keep track of some analytical properties. The next definition summarizes the second paragraph of Berndt's Chapter 5. (Throughout his Chapter 5, $\lambda < 2$.)

Definition 4. Let $f \in M(\lambda, k, \gamma)$, f not identically zero.

1. $N = N_f$ counts the zeros of f on $\overline{B(\lambda)}$ with multiplicities.
2. N_f does not count zeros at τ_λ , at $\tau_\lambda + \lambda$, at i , or at $i\infty$.
3. If $\tau_0 \in \overline{B(\lambda)}$, $f(\tau_0) = 0$ and $\Re(\tau_0) = -\lambda/2$, then $f(\tau_0 + \lambda) = 0$ and N_f counts only one of the two zeros.
4. If $\tau_0 \in \overline{B(\lambda)}$, $f(\tau_0) = 0$, and $|\tau_0| = 1$, then, $f(-1/\tau_0) = 0$, and N_f counts only one of these two zeros.
5. The numbers n_λ, n_i , and n_∞ are the orders of the zeros of f at τ_λ, i and $i\infty$, respectively. The order n_∞ is measured in terms of $\exp(2\pi i \tau / \lambda)$.

The multiplier γ is given by

Theorem 3. ([4], Corollary 5.2) Let $f \in M(\lambda, k, \gamma)$ and let n_i be the order of the zero of f at $\tau = i$. Then

$$\gamma = (-1)^{n_i}.$$

The next two results tell us that the only nontrivial case in this theory is the one that we are interested in.

Theorem 4. ([4], Lemma 5.1) If $\dim M(\lambda, k, \gamma) \neq 0$,

$$N_f + n_\infty + \frac{1}{2}n_i + \frac{n_\lambda}{m} = \frac{1}{2}k \left(\frac{1}{2} - \theta \right).$$

By Berndt's equation (5.16), if $m \geq 3$ then the right side can be written as $k(m-2)/4m$.

Theorem 5. ([4], Theorem 5.2) If $\dim M(\lambda, k, \gamma) \neq 0$, then $\theta = 1/m$ where $m \geq 3$ and $m \in \mathbb{Z}$.

We are concerned with $\lambda \in T_A$. This makes $\lambda < 2$ as in all the results of Berndt's Chapter 5.

One estimate for $\dim M(\lambda, k, \gamma)$ is

Theorem 6. ([4], Theorem 5.6) If $\lambda \notin T_A$, then $\dim M(\lambda, k, \gamma) = 0$. If $\lambda = 2 \cos(\pi/m) \in T_A$, then for nontrivial $f \in M(\lambda, k, \gamma)$, the weight k has the form

$$k = \frac{4h}{m-2} + 1 - \gamma,$$

where $h \geq 1$ is an integer. Furthermore,

$$\dim M(\lambda, k, \gamma) = 1 + \left\lfloor \frac{h + (\gamma - 1)/2}{m} \right\rfloor.$$

Eliminating h , we find that

$$\dim M(\lambda, k, \gamma) = 1 + \left\lfloor k \left(\frac{1}{4} - \frac{1}{2m} \right) + \frac{\gamma}{4} - \frac{1}{4} \right\rfloor. \quad (13)$$

Berndt ([4], Remark 5.3) proves that the dimension formula above holds also when $h = 0$.

The existence of certain modular forms is provided by

Theorem 7. ([4], Theorem 5.5) Let $\lambda \in T_A$. Then there exist functions f_λ, f_i , and $f_\infty \in M(\lambda, k, \gamma)$ such that each has a simple zero at τ_λ, i , and $i\infty$, respectively, and no other zeros. Here, γ is given by [Theorem 3 (this article)], and k is determined in each case from [Theorem 4 (this article)]. Thus, $f_\lambda \in M(\lambda, 4/(m-2), 1)$, $f_i \in M(\lambda, 2m/(m-2), -1)$, and $f_\infty \in M(\lambda, 4m/(m-2), 1)$.

Remark 2. ([4], pages 47-48) By the Riemann mapping theorem there exists a function $g(\tau)$ that maps the simply connected region $B(\lambda)$ one-to-one and conformally onto \mathbb{H} . If we require that $g(\tau_\lambda) = 0$, $g(i) = 1$, and $g(i\infty) = \infty$, then g is determined uniquely.

Now we can write down f_λ , f_i , and f_∞ explicitly. The next theorem is extracted from the proof of Theorem 7. f_λ and f_i correspond to Eisenstein series and f_∞ to a cusp form. In our code, we take g to be a normalized form of J_m .

Theorem 8. ([4], page 50)

$$f_\lambda(\tau) = \left\{ \frac{g'(\tau)^2}{g(\tau)(g(\tau) - 1)} \right\}^{1/(m-2)},$$

$$f_i(\tau) = \left\{ \frac{g'(\tau)^m}{g(\tau)^{m-1}(g(\tau) - 1)} \right\}^{1/(m-2)},$$

and

$$f_\infty(\tau) = \left\{ \frac{g'(\tau)^{2m}}{g(\tau)^{2m-2}(g(\tau) - 1)^m} \right\}^{1/(m-2)}.$$

In our applications to Lehmer's problem, we will be interested in the dimensions of the weight 12 cusp spaces for $\lambda = \lambda_m = 2 \cos \pi/m$.

Definition 5. ([4], Definition 5.2) If $f \in M(\lambda, k, \gamma)$ and $f(i\infty) = 0$, then we call f a cusp form of weight k and multiplier γ with respect to $G(\lambda)$. We denote by $C(\lambda, k, \gamma)$ the vector space of all cusp forms of this kind.

Remark 3. ([4], equation (5.25))

$$\dim C(\lambda, k, \gamma) \geq \dim M(\lambda, k, \gamma) - 1.$$

Remark 4. In view of (i) Theorem 6, (ii) equation (12), (iii) Remark 2, and (iv) the fact that $\gamma = \pm 1$, we see that $\dim C(\lambda_m, 12, \gamma) > 1$ when m is greater than or equal to 12.

6 The objects in our experiments

In this section, we write down versions of the functions from Theorem 8 such that, at $m = 3$, they reduce to corresponding functions in the classical theory. The classical objects (in Serre's notation [29]), are Klein's j -invariant, the weight four Eisenstein series E_2 , the weight six Eisenstein series E_3 , and the generating function of Ramanujan's tau function, namely the normalized weight twelve cusp form Δ . Our definitions come in pairs because we find that, in this way, we can extract a little more information about the interpolating polynomials in some situations. The second definition in each pair replaces 1 by $2^6 m^3$, but (when it matters) they can be cast into the original form up to a multiplicative constant depending upon m .

We have already normalized the Fourier expansions of the J_m by replacing x_m with X_m in the expansions. Now we normalize the J_m themselves to obtain functions j_m such that $j_3 =$ the usual j function. (Recall that X_m is a function of τ , which, in this context, is a point in the upper half-plane.)

Definition 6. *Writing the Fourier expansion of $J_m(\tau)$ as $\sum_{n=-1}^{\infty} a_n(m)X_m^n$, we set $\bar{J}_m(\tau) := J_m(\tau)/a_{-1}(m) =$ (say) $1/X_m + \sum_{n=0}^{\infty} \bar{a}_n(m)X_m^n$, and $j_m(\tau) := 1/X_m + \sum_{n=0}^{\infty} \bar{a}_n(m)(2^6 m^3 X_m)^n$.*

In general, we find that the substitution $X_m \mapsto 2^6 m^3 X_m$ creates functions that behave less mysteriously than the originals when analyzing the interpolating polynomials we want to study; we do not know why. In particular, after this substitution, when the interpolating polynomials are written as products of monic irreducible factors, the leading numerical coefficients often form recognizable sequences, as we will see below. This is not typical in the absence of this substitution (which appears in the thesis of Leo [21].)

Corresponding to f_λ ,

Definition 7. 1. $H_{\lambda,m}(\tau) :=$

$$\left\{ \frac{J'_m(\tau)^2}{J_m(\tau)(J_m(\tau) - 1)} \right\}^{1/(m-2)}.$$

2. $H_{\lambda,4,m}(\tau) := H_{\lambda,m}(\tau)^{m-2}.$

3. *Writing the Fourier expansion of $H_{\lambda,4,m}(\tau)$ as $\sum_{n=0}^{\infty} h_{4,n}(m)q_m(\tau)^n$, we set $H_{\lambda,4,m}^*(\tau) := \sum_{n=-1}^{\infty} h_{4,n}(m)(2^6 m^3 q_m(\tau))^n$.*

4. $H_{\lambda,6,m}(\tau) := H_{\lambda,m}(\tau)^{\frac{3}{2}(m-2)}.$

5. $\tilde{H}_{\lambda,m}(\tau) :=$

$$\left\{ \frac{j'_m(\tau)^2}{j_m(\tau)(j_m(\tau) - 2^6 m^3)} \right\}^{1/(m-2)}.$$

6. $\tilde{H}_{\lambda,4,m}(\tau) := \tilde{H}_{\lambda,m}(\tau)^{m-2}.$

7. $\tilde{H}_{\lambda,6,m}(\tau) := \tilde{H}_{\lambda,m}(\tau)^{\frac{3}{2}(m-2)}.$

Corresponding to f_i , we state

Definition 8. 1.

$$H_{i,m}(\tau) := \left\{ \frac{J'_m(\tau)^m}{J_m(\tau)^{m-1}(J_m(\tau) - 1)} \right\}^{1/(m-2)}.$$

2. $H_{i,4,m}(\tau) := H_{i,m}(\tau)^{2(m-2)/m}.$

3. $H_{i,6,m}(\tau) := H_{i,m}(\tau)^{3(m-2)/m}$.
4. Let $\epsilon(m) = e^{-i\pi/(m-2)}$ or one, depending on whether m is odd or even, respectively.

$$\tilde{H}_{i,m}(\tau) := \epsilon(m) \left\{ \frac{j'_m(\tau)^m}{j_m(\tau)^{m-1}(j_m(\tau) - 2^6 m^3)} \right\}^{1/(m-2)}.$$

5. $\tilde{H}_{i,4,m}(\tau) := \tilde{H}_{i,m}(\tau)^{2(m-2)/m}$.
6. $\tilde{H}_{i,6,m}(\tau) := \tilde{H}_{i,m}(\tau)^{3(m-2)/m}$.

Remark 5. In view of Berndt's theorem 7 above, the functions defined in items 2, 3 and 6 of definition 7 and items 2 and 5 of definition 8 all have weight 4. Similarly, the functions defined in items 4 and 7 of definition 7 and items 3 and 6 of definition 8 have weight 6.

In our *Mathematica* code, Hecke modular forms and functions are expressed as power series in x_m , so that *Mathematica* is performing a purely formal series operation when the exponent $1/(m-2)$ is applied in the series above. Furthermore, in order to obtain a series with rational coefficients for $\tilde{H}_{i,m}$, we found we had to use the *Mathematica* command `Refine[x>0]` at this stage. The result agrees at $m=3$ with the weight six Eisenstein series $E_3 = 1 - 504 \sum_{n=1}^{\infty} \sigma_5(n) q^n$ on page 93 of [29].

Definition 9. 1. Corresponding to f_{∞} , we have

$$\Delta_{\infty,m}(\tau) := \left\{ \frac{J'_m(\tau)^{2m}}{J_m(\tau)^{2m-2}(J_m(\tau) - 1)^m} \right\}^{1/(m-2)}.$$

2. $\Delta_m^{\dagger} := H_{\lambda,m}^3 - H_{i,m}^2$.
3. $\Delta_{12,m}^{\diamond} := H_{\lambda,4,m}^3 / J_m$.
4. Writing the Fourier expansion of $\Delta_{12,m}^{\diamond}(\tau)$ as $\sum_{n=1}^{\infty} \tau_{12,m}^{\diamond} q_m(\tau)^n$, we set $\Delta_{12,m}^{\diamond\diamond}(\tau) := \sum_{n=1}^{\infty} \tau_{12,m}^{\diamond\diamond} (2^6 m^3 q_m(\tau))^n$.
- 5.

$$\tilde{\Delta}_{\infty,m}(\tau) := \left\{ \frac{j'_m(\tau)^{2m}}{j_m(\tau)^{2m-2}(j_m(\tau) - 2^6 m^3)^m} \right\}^{1/(m-2)}.$$

6. $\tilde{\Delta}_m^{\dagger} := \tilde{H}_{\lambda,m}^3 - \tilde{H}_{i,m}^2$.
7. $\tilde{\Delta}_m^{\diamond} := \tilde{H}_{\lambda,m}^3 / j_m$.

Remark 6. By definition 1, f^2 belongs to $M(\lambda, 2k, 1)$ if f belongs to $M(\lambda, k, -1)$. Therefore, the functions defined above in items 1, 2, 5, 6 and 7 belong to $M(\lambda, 12/(m-2), 1)$. On the other hand, the functions defined in items 3 and 4 belong to $M(\lambda, 12, 1)$.

7 Interpolation by polynomials of the Fourier coefficients of modified modular forms and functions for Hecke groups

In this section, we state conjectures about polynomials interpolating coefficients of modular forms for Hecke groups. Conjectures 6 and 7 bear on Lehmer's question about the existence of zeros of Ramanujan's tau function.

As before, let

$$J_m(\tau) = \sum_{n=-1}^{\infty} a_n(m) q_m(\tau)^n.$$

For integers $m \geq 3$ Raleigh showed that, for $n = 0, 1, 2, 3$, $a_n(m) = m^{-2n-2} a_{-1}(m)^{-n} P_n(m)$ where $P_n(x)$ is a polynomial with rational coefficients and degree $2n + 2$, such that the coefficient of x^n is zero when n is odd. As we said in the introduction, he conjectured, and Akiyama proved, that similar relations exist among the a_n for all positive n .

Berndt's (Hecke's) theorems 7 and 8 above make it clear that Akiyama's result extends in some way to the modular forms defined in the previous section. Our experiments were done to explore the details; the observations are summarized in the conjectures below.

Conjecture 1. *blah*

Conjecture 2. *Let the Fourier expansion of j_m be*

$$j_m = 1/q_m + \sum_{n \geq 0} c_n(m) q_m(\tau)^n.$$

1. *For each integer n greater than -2 , there exists a polynomial $C_n(x) \in \mathbb{Q}[x]$ that satisfies the relation $c_n(m) = C_n(m)$ for $m = 3, 4, \dots$, with $C_{-1}(x) \equiv 1$, $C_0(x) = 24x(x^2 + 4/3)$, and $C_1(x) = 276x^2(x^4 - 8x^2/69 - 16/23)$.*
2. *Let n be larger than one. Then*
 - (a) $C_n(x) = \phi_n(x - 2)(x + 2)x^{n+1}\gamma_n(x)$ where ϕ_n is a rational number and $\gamma_n(x)$ an irreducible monic polynomial in $\mathbb{Q}[x]$.
 - (b) $\{\phi_n\}$ is the McKay-Thompson series of class 4A (item 10 in our glossary).
 - (c) The coefficient of x^k in $\gamma_n(x)$ is non-zero if and only if k is even and not greater than $2n$.
 - (d) All of the complex roots of $\gamma_n(x)$ lie in the disk with center zero and radius $n/\log(n)$.
3. *For n greater than one, let G_n be the Galois group of $\gamma_n(x)$ over the rationals. The size of G_n is $2^n n!$ and (if n is greater than two) G_n is isomorphic*

to a permutation group on $2n$ elements $\{e_1, \dots, e_{2n}\}$ with three generators: a transposition (e_j, e_k) , a product $(e_j, e_{j'})(e_k, e_{k'})$, and a product $\Gamma_1 \Gamma_2$ of disjoint cycles Γ_1 and Γ_2 , each of length n , such that Γ_1 sends e_j to $e_{j'}$ and Γ_2 sends e_k to $e_{k'}$.

Clause 2 implies that, for m greater than or equal to three, $c_n(m)$ is nonzero. It is already known that, for all integers $n \geq -1$, the n^{th} Fourier coefficient of $j = j_3$, namely $c(n) = c_n(3)$, is positive; see, for example, page 199 in [26].

We tested clause 2(d) in several ways. We approximated the roots of the $\gamma_n(x)$ with root-finding routines and compared their complex moduli with $n/\log(n)$. We used the argument principle to count the zeros in central disks of radius $n/\log(n)$. We superimposed plots of the roots of $\gamma_n(x)$ against plots of circles with radius $n/\log(n)$ and center at the origin. An example is depicted in Figure 1. (We produced this plot in our *Mathematica* notebook “conjecture1clause2d.nb”. The words in the caption “length of data file” refer to the number of values of m that we tested to arrive at the interpolating polynomial.) We tested the γ_n for $n \leq 24$. For clause 3, we computed the Galois groups in *Magma*.

Remark 7. Like G_n in clause 3, the index- n hyperoctahedral group has size $2^n n!$ [12, 23, 15]. It seems natural to ask whether or not they are isomorphic.

We found the sequence $\{c_{4,n}\}$ mentioned below on page A008547 of [30].

Conjecture 3. Let the Fourier expansion of $H_{\lambda,4,m}^*(\tau)$ be $\sum_{n=0}^{\infty} \beta_{4,n}(m) q_m(\tau)^n$.

1. For each n there is a polynomial $B_{4,n}(x)$ with rational coefficients such that $\beta_{4,n}(m) = B_{4,n}(m)$ for $m = 3, 4, \dots$
2. If n is positive, then the degree of $B_{4,n}(x)$ is $6n$.
3. $B_{4,0}(x) \equiv 1$ and, if n is positive, then

$$B_{4,n}(x) = e_{4,n}(x^2 - 4)x^{4n}b_{4,n}(x),$$

where $e_{4,n} = 16 \sum_{\nu|n, \nu \text{ odd}} (-1)^{n-\nu} \nu^3$ and $b_{4,n}(x)$ is a monic irreducible polynomial in $\mathbb{Q}[x]$.

Conjecture 4. Let the Fourier expansion of $\tilde{H}_{\lambda,m}$ be

$$\tilde{H}_{\lambda,m} = \sum_{n=0}^{\infty} \beta_n(m) q_m(\tau)^n.$$

1. For each n there is a polynomial $B_n(x)$ with rational coefficients such that $\beta_n(m) = B_n(m)$ for $m = 3, 4, \dots$
2. If n is positive, then the degree of $B_n(x)$ is $3n - 1$.
3. $B_0(x) \equiv 1$ and $B_1(x) = 16x(x + 2)$.

4. Let \mathcal{Q} be as in item 9 of our glossary and let $e_n = 16(-1)^{n+1} \sum_{\nu|n} 1/\nu$.
If $n \in \mathcal{Q}$, then

$$B_n(x) = e_n(x^2 - 4)(x - 6)x^n b_n(x),$$

where $b_n(x)$ is a monic irreducible polynomial. Otherwise (for n greater than one) $B_n(x) = e_n(x^2 - 4)x^n b_n(x)$ where, again, $b_n(x)$ is a monic irreducible polynomial in $\mathbb{Q}[x]$.

(We identified the e_n after reading pages A034020 and A098985 in [30].)

Thus, in the range of our observations ($3 \leq m \leq 302, 0 \leq n \leq 100$), the only integer value of m such that $\tilde{H}_{\lambda,m}$ has any vanishing coefficients is six, and $\beta_n(6)$ is zero just if n is in \mathcal{Q} .

Conjecture 5. Let the Fourier expansion of $\tilde{H}_{i,m}(\tau)$ be

$$\tilde{H}_{i,m}(\tau) = \sum_{n=0}^{\infty} \delta_n(m) q_m(\tau)^n.$$

1. For each non-negative integer n , there is a polynomial $D_n(x)$ in $\mathbb{Q}[x]$ such that
 - (a) The degree of D_n is $3n$.
 - (b) $D_n(x) = a$ rational number $d_n \times$ a product of monic irreducible polynomials.
 - (c) $d_0 = 1$ and, for n a positive integer, $d_n = 24(-1)^n \sum_{\nu|n}^* \nu$. Again, the asterisk means that the sum is taken over the odd positive divisors of n .
2. $\delta_n(m) = D_n(m)$ for $m = 3, 4, \dots$
3. $D_0(x) \equiv 1$ identically, $D_1(x) = -24(x - 2/3)x^2$, and $D_2(x) = 24(x - 2/3)(x - 2)x^3(x - 14)$.
4. For n larger than two, $D_n(x) = d_n(x - 2)(x - 2/3)x^{n+1}\epsilon_n(x)$ where $\epsilon_n(x)$ is a monic irreducible polynomial in $\mathbb{Q}[x]$.

Let Δ be usual normalized discriminant, a weight 12 cusp form for $SL(2, \mathbb{Z}) = G(\lambda_3)$ with integer coefficients. Its Fourier expansion is written

$$\Delta(\tau) = \sum_{n=1}^{\infty} \tau(n) q^n$$

where $q = e^{2\pi i \tau}$ and $\tau(n)$ is Ramanujan's function. (We trust that the reader will not confuse the complex number τ with Ramanujan's function $\tau(n)$ or any of its relatives defined below.) Whether or not the equation $\tau(n) = 0$ has any solutions is (notoriously) an open question [19]. (Recently, Balakrishnan, Craig, and Ono [3] ruled out $\pm 1, \pm 3, \pm 5, \pm 7$, and ± 691 from membership in the image of tau.)

Conjecture 6. Let the Fourier expansion of $\Delta_{12,m}^{\circ\circ}(\tau)$ be

$$\Delta_{12,m}^{\circ\circ}(\tau) = \sum_{n=1}^{\infty} \tau_{12,m}^{\circ\circ}(n) q_m(\tau)^n.$$

1. There is a set of polynomials $T_{12,n}^{\circ\circ}(x)$ with coefficients in \mathbb{Q} such that $\tau_{12,m}^{\circ\circ}(n) = T_n(m)$ for each $m = 3, 4, \dots$
2. $T_{12,n}^{\circ\circ}(x) = (-1)^{n+1} 64 \tau(n) x^{n+3} t_{12,n}^{\circ\circ}(x)$, where $t_{12,n}^{\circ\circ}(x)$ is a monic irreducible polynomial over \mathbb{Q} of degree $2n - 2$.

Conjecture 7. Let the Fourier expansion of $\tilde{\Delta}_{\infty,m}(\tau)$ be

$$\tilde{\Delta}_{\infty,m}(\tau) = \sum_{n=1}^{\infty} \tilde{\tau}_{\infty,m}(n) q_m(\tau)^n.$$

1. There is a set of polynomials $\tilde{T}_{\infty,n}(x)$ with coefficients in \mathbb{Q} such that $\tilde{\tau}_{\infty,m}(n) = \tilde{T}_{\infty,n}(m)$.
2. $\tilde{T}_{\infty,1}(x) \equiv 1$ identically, and, if n is greater than one,
 - (a) $\tilde{T}_{\infty,n}(x) = \tilde{s}_{\infty,n}(x-2)^2 x^{n-1} \tilde{t}_{\infty,n}(x)$, where $\tilde{t}_{\infty,n}(x)$ is a monic irreducible polynomial over \mathbb{Q} of degree $2n - 4$ and
 - (b) $\tilde{s}_{\infty,n}$ is (in the notation of [11], Chapter 7, Theorem 7) the coefficient of q^n in the Fourier expansion of $\Delta_8(z)$ [31].
 - (c) Also ([32]),

$$\tilde{s}_{\infty,n} = (-1)^{n+1} \sum_{\substack{\nu|n \\ \nu \text{ odd}}} \nu^3.$$

This sum is the coefficient of q^n in the Fourier expansion of $E_{\infty,4}$, the unique normalized weight-4 modular form for $\Gamma_0(2)$ with simple zeros at $i\infty$ ([6], equation (2-3)); it is also the number of representations of $n - 1$ as a sum of 8 triangular numbers ([24], theorem 5.)

- (d) Finally, $\tilde{s}_{\infty,n}$ is the coefficient of q^n in the expansion of $\eta(2z)^{16}/\eta(z)^{-8}$ where $\eta(z)$ is Dedekind's function ([6], equation (2-16).)

Conjecture 8. Let the Fourier expansion of $\tilde{\Delta}_m^{\dagger}(\tau)$ be

$$\tilde{\Delta}_m^{\dagger}(\tau) = \sum_{n=1}^{\infty} \tilde{\tau}_m^{\dagger}(n) q_m(\tau)^n.$$

1. There is a set of polynomials $\tilde{T}_n^{\dagger}(x)$ with coefficients in \mathbb{Q} such that $\tilde{\tau}_m^{\dagger}(n) = \tilde{T}_n^{\dagger}(m)$.
2. $\tilde{T}_1^{\dagger}(x) = 16x(3x^2 + x + 6)$.

3. $\tilde{T}_2^\dagger(x) = -16x^2(39x^4 - 95x^3 + 66x^2 - 260x - 120)$.
4. $\tilde{T}_3^\dagger(x) = 64x^3(189x^6 - 3021x^5 + 9574x^4 - 12520x^3 + 19136x^2 - 2960x - 2208)/9$.
5. If n is greater than 3, then
 - (a) $\tilde{T}_n^\dagger(x) = \tilde{s}_n^\dagger(x-2)^2x^n\tilde{t}_n^\dagger(x)$, where $\tilde{t}_n^\dagger(x)$ is a monic polynomial, irreducible over \mathbb{Q} , of degree $2n-1$ and
 - (b) \tilde{s}_n^\dagger is (in the notation of [11], Chapter 7, Section 2) the coefficient of q^n for the theta series of the direct sum of two copies of the D_4 lattice in powers of q^2 [33].
 - (c) Also (proposition 7.5 of [22]) $\tilde{s}_n^\dagger = (-1)^n \times$ the coefficient of q^{2n} in the Fourier expansion of $\{f_8, z\}$ where $f_8(z) = \eta(4z)^{12}/(\eta(2z)^4\eta(8z)^8)$. Here f_8 is the normalized Hauptmodul for the $SL(2, \mathbb{Z})$ subgroup $\Gamma_0(8)$ and the braces stand for the Schwarzian derivative (item 2 in our glossary.)

Conjecture 9. Let the Fourier expansion of $\tilde{\Delta}_m^\diamond(\tau)$ be

$$\tilde{\Delta}_m^\diamond(\tau) = \sum_{n=1}^{\infty} \tilde{\tau}_m^\diamond(n) q_m(\tau)^n.$$

1. There is a set of polynomials $\tilde{T}_n^\diamond(x)$ with coefficients in \mathbb{Q} such that $\tilde{\tau}_m^\diamond(n) = \tilde{T}_n^\diamond(m)$.
2. $\tilde{T}_1^\diamond(x)$, $\tilde{T}_2^\diamond(x)$, and $\tilde{T}_3^\diamond(x)$ are irreducible polynomials over \mathbb{Q} of degrees 3, 6, and 9, respectively.
3. If n is greater than 3, $\tilde{T}_n^\diamond(x) = \tilde{s}_n^\diamond(x-2)x^{n-1}\tilde{t}_n^\diamond(x)$, where $\tilde{t}_n^\diamond(x)$ is a monic polynomial, irreducible over \mathbb{Q} , of degree $2n-3$. Furthermore,
 - (a) $\sum_{n=0}^{\infty} \tilde{s}_n^\diamond q(\tau)^n = \prod_{n \text{ odd}} (1 - q(\tau)^n)^{24} \times \prod_{n \equiv 2(4)} (1 - q(\tau)^n)^{-24} = \eta^{24}(\tau) \eta^{24}(4\tau) \eta^{-48}(2\tau)$.
 - (b) $\tilde{s}_n^\diamond = (-1)^{n+1} \times$ the coefficient of $q(\tau)^n$ in $(\eta(2\tau)/\eta(\tau))^{24}$.

The product decomposition in clause 3(a) above is a guess based on 43 terms of the series using Euler's method [13]; also, [2], Theorem 14.8. We find that it also appears in [36]. The second decomposition appears in [34].

Remark 8. We relate conjectures 6, 7, 8 and 9 to Lehmer's question on the existence of zeros for Ramanujan's tau function as follows. First (from clause 2 of conjecture 6), for $m = 3, 4, \dots$, $\tau_{12,m}^\diamond(n) = 0$ if and only if $\tau(n) = 0$. Second, let us write $S(x) := \sum \tau(n)x^n$, and regard this as a formal power series. Let us also write down the formal power series

1. $\tilde{S}_\infty(x) := \sum \tilde{\tau}_{\infty,3}(n)x^n$.
2. $\tilde{S}^\dagger(x) := \sum \tilde{\tau}_3^\dagger(n)x^n$.
3. $\tilde{S}^\circ(x) := \sum \tilde{\tau}_3^\circ(n)x^n$.
4. $S_{12}^\circ(x) := \sum \tau_{12,3}^\circ(n)x^n$.

Then we have (empirically) $S(x) = \tilde{S}_\infty(x) = \tilde{S}^\dagger(x) = \tilde{S}^\circ(x) = A_3 S_{12}^\circ(x/A_3) = S_{12}^\circ(1728x)/1728$. Consequently, for $T_n(x) = \tilde{T}_{\infty,n}(x), \tilde{T}_n^\dagger(x), \tilde{T}_n^\circ(x)$, or $T_{12,n}^\circ(x)$, if $T_n(3)$ is not equal to zero, neither is $\tau(n)$.

More generally:

Conjecture 10. Letting $T_n(x)$ and τ_m stand for the various polynomials and Fourier coefficients in conjectures 6 through 9, none of the $T_n(x)$ takes an integer greater than two to zero; consequently, none of the τ_m vanishes for $m = 3, 4, \dots$

For $T_n(x)$ as above, for each positive integer n , and for each integer m greater than two, let the minimum Euclidean distance to m of any complex root of $T_n(x)$ be denoted as $d(m, n)$. Conjecture 10 can be restated by saying that $d(m, n)$ is positive. Our experiments indicate, in fact

Conjecture 11. That

1. for fixed m , $d(m, n)$ decays exponentially as n increases, and,
2. if m is greater than three, then $d(m, n) > d(3, n)$.

Remark 9. While checking our calculations, we compared the Fourier expansion of $H_{\lambda,4}(x/A_4)$ (abusing notation in the obvious way) with Leo's ([21], p.54) expansion of the weight 4 Eisenstein series at $m = 4$. (Recall that $A_4 = 1/256$.) Within the range of our observations, they do coincide. The expansions (in our own notation) both begin

$$1 + 48q_4 + 624q_4^2 + 1344q_4^3 + \dots$$


We also read the O.E.I.S page [33], which mentions our expansion ([7], equation (2-1), p, 260):

$$E_{\gamma,2} = 1 + 24 \sum_{n=1}^{\infty} \sum_{\substack{\nu|n \\ \nu \text{ odd}}} \nu q^n.$$

On the O.E.I.S. page, N. J. A. Sloane comments that the sequence $\{1, 48, 624, \dots\}$ is the same as that of the coefficients of $E_{\gamma,2}^2$. It happens that $E_{\gamma,2}^2$ is a weight 4, level 2 modular form, that is, a weight 4 modular form for the $SL(2, \mathbb{Z})$ subgroup $\Gamma_0(2)$. While it is too vague to state as a conjecture, we venture a guess that, behind this observation and the double role of certain rational numbers as coefficients of classical Hauptmoduln and as coefficients in our interpolating polynomials, there is a theory connecting level N classical modular forms with Hecke modular forms.

8 Figures

8.1 Figure 1.



ROOTS OF POLYNOMIAL INTERPOLATING THE COEFFICIENT OF q_m^{23} IN THE FOURIER EXPANSION OF j_m .

References

- [1] Shigeki Akiyama. “A note on Hecke’s absolute invariants”. In: *J. Ramanujan Math. Soc* 7.1 (1992), pp. 65–81.
- [2] Tom M Apostol. *Introduction to analytic number theory*. Springer-Verlag, 1976.
- [3] Jennifer S Balakrishnan, William Craig, and Ken Ono. “Variations of Lehmer’s Conjecture for Ramanujan’s tau-function”. In: *Journal of Number Theory* (2020).
- [4] Bruce C Berndt and Marvin Isadore Knopp. *Hecke’s theory of modular forms and Dirichlet series*. Vol. 5. World Scientific, 2008.
- [5] B. Brent. *Interpolations repository*. <https://github.com/barry314159a/interpolations>.

- [6] Barry Brent. “Quadratic minima and modular forms”. In: *Experimental Mathematics* 7.3 (1998), pp. 257–274.
- [7] Barry Brent. “Quadratic minima and modular forms II”. In: *Acta Arithmetica* 96.4 (2001), pp. 381–387.
- [8] I. N. Cangül. “The group structure of Hecke groups $H(\lambda_q)$ ”. In: *Turkish Journal of Mathematics* 20.2 (1996), pp. 203–207.
- [9] Constantin Carathéodory. *Theory of functions of a complex variable, Second English Edition*. Vol. 1. Translated by F. Steinhardt. Chelsea Publishing Company., 1958.
- [10] Constantin Carathéodory. *Theory of functions of a complex variable, Second English Edition*. Vol. 2. Translated by F. Steinhardt. Chelsea Publishing Company., 1981.
- [11] John Horton Conway and Neil James Alexander Sloane. *Sphere packings, lattices and groups*. Vol. 290. Springer Science & Business Media, 2013.
- [12] Wikipedia The Free Encyclopedia. *Hyperoctahedral group*. https://en.wikipedia.org/wiki/Hyperoctahedral_group. 23 October 2020.
- [13] L Euler. “Discovery of a most extraordinary law of numbers concerning the sum of their divisors, Opera Omnia, Ser. 1”. In: *Eneström index* 2 (E175), pp. 241–253.
- [14] Robert Fricke. *Die elliptischen Funktionen und ihre Anwendungen*. Vol. 1. BG Teubner, 1922.
- [15] Piotr Graczyk, Gérard Letac, and Hélène Massam. “The hyperoctahedral group, symmetric group representations and the moments of the real Wishart distribution”. In: *Journal of Theoretical Probability* 18.1 (2005), pp. 1–42.
- [16] Erich Hecke. *Lectures on Dirichlet series, modular functions, and quadratic forms*. Vandenhoeck & Ruprecht, 1983.
- [17] Erich Hecke. “Über die bestimmung dirichletscher reihen durch ihre funktionalgleichung”. In: *Mathematische Annalen* 112.1 (1936), pp. 664–699.
- [18] Jonas Jermann. *Triangle functions on SageMath site*. https://doc.sagemath.org/html/en/reference/modfrm_hecketriangle/index.html. 2013.
- [19] D. H. Lehmer. “The vanishing of Ramanujan’s function $\tau(n)$ ”. In: *Duke Mathematical Journal* 14.2 (1947), pp. 429–433.
- [20] Joseph Lehner. “Note on the Schwarz triangle functions.” In: *Pacific Journal of Mathematics* 4.2 (1954), pp. 243–249.
- [21] John Garrett Leo. *Fourier coefficients of triangle functions, Ph.D. thesis*. <http://halfaya.org/ucla/research/thesis.pdf>. 2008.
- [22] John Mckay and Hubertus Strauss. “The q-series of monstrous moonshine & the decomposition of the head characters”. In: *Communications in Algebra* 18.1 (1990), pp. 253–278. DOI: 10.1080/00927879008823911. URL: <https://doi.org/10.1080/00927879008823911>.

- [23] GA Miller. “Groups formed by special matrices”. In: *Bulletin of the American Mathematical Society* 24.4 (1918), pp. 203–206.
- [24] Ken Ono, Sinai Robins, and Patrick T Wahl. “On the representation of integers as sums of triangular numbers”. In: *Aggregating clones, colors, equations, iterates, numbers, and tiles*. Springer, 1995, pp. 73–94.
- [25] J Raleigh. “On the Fourier coefficients of triangle functions”. In: *Acta Arithmetica* 8 (1962), pp. 107–111.
- [26] R.A. Rankin. *Modular Forms and Functions*. Cambridge University Press, Cambridge, 1977.
- [27] Bruno Schoeneberg. *Elliptic modular functions*. Springer, 1974, p. 45.
- [28] Hermann Amandus Schwarz. “Ueber diejenigen Fälle, in welchen die Gaussische hypergeometrische Reihe eine algebraische Function ihres vierten Elementes darstellt.” In: *Journal für die reine und angewandte Mathematik* 75 (1873), pp. 292–335.
- [29] Jean-Pierre Serre. *A course in arithmetic*. Springer-Verlag, 1970.
- [30] Neil J. A. Sloane. *The On-Line Encyclopedia of Integer Sequences*. <https://oeis.org/>.
- [31] Neil J. A. Sloane. *The On-Line Encyclopedia of Integer Sequences*, A002408. <http://oeis.org/A002408>.
- [32] Neil J. A. Sloane. *The On-Line Encyclopedia of Integer Sequences*, A007331. <http://oeis.org/A007331>.
- [33] Neil J. A. Sloane. *The On-Line Encyclopedia of Integer Sequences*, A008658. <http://oeis.org/A008658>.
- [34] Neil J. A. Sloane. *The On-Line Encyclopedia of Integer Sequences*, A014103. <http://oeis.org/A014103>.
- [35] Neil J. A. Sloane. *The On-Line Encyclopedia of Integer Sequences*, A097340. <https://oeis.org/A097340>.
- [36] Neil J. A. Sloane. *The On-Line Encyclopedia of Integer Sequences*, A100130. <http://oeis.org/A100130>.