

Assignment 1 for #70240413

”Statistical Machine Learning”

Kui XU, 2016311209

2017/03/23

1 Mathematics Basics

Choose one problem from the 1.1 and 1.2. A bonus would be given if you finished the both.

1.1 Calculus

The gamma function is defined by (assuming $x > 0$)

$$\Gamma(x) = \int_0^{\infty} u^{x-1} e^{-u} du. \quad (1)$$

(1) Prove that $\Gamma(x+1) = x\Gamma(x)$.

(2) Also show that

$$\int_0^1 u^{a-1} (1-u)^{b-1} du = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}. \quad (2)$$

Solution: For Question (1), we can prove it by Using integration by parts, the steps are as follows:

$$\begin{aligned} \Gamma(x+1) &= \int_0^{\infty} u^x e^{-u} du \\ &= [-u^x e^{-u}]_0^{\infty} + \int_0^{\infty} x u^{x-1} e^{-u} du \\ &= \lim_{u \rightarrow \infty} (-u^x e^{-u}) - (0e^{-0}) + x \int_0^{\infty} u^{x-1} e^{-u} du \\ &= x \int_0^{\infty} u^{x-1} e^{-u} du \\ &= x\Gamma(x) \end{aligned} \quad (3)$$

As we know, when $u \rightarrow \infty$, $-u^x e^{-u} \rightarrow 0$, so the equation is proved.

Solution: For Question (2), we know that the left of the equation is a Beta function. From the definitions, we can express the equation which we want to prove as :

$$\Gamma(a+b)B(a,b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)} \quad (4)$$

It's a double integral, the expansion formula is as follows:

$$\begin{aligned} \Gamma(a+b)B(a,b) &= \int_0^{\infty} u^{a+b-1} e^{-u} du \int_0^1 v^{a-1} (1-v)^{b-1} dv \\ &= \int_0^{\infty} \int_0^1 (uv)^{a-1} [u(1-v)]^{b-1} u e^{-u} du dv \end{aligned} \quad (5)$$

Then we do a transformation $w = uv$, $z = u(1-v)$. The inverse transformation is $u = w+z$, $v = w/(w+z)$, the corresponding ranges of them are $w \in (0, \infty)$ and $u \in (0, \infty)$. The absolute value of the Jacobian is

$$\left| \nabla \frac{\partial(u, v)}{\partial(w, z)} \right| = \frac{1}{(w+z)} \quad (6)$$

Next, we use the changed of variables to do a double integral, the equation above becomes:

$$\begin{aligned} & \int_0^\infty \int_0^\infty w^{a-1} z^{b-1} (w+z) e^{-(w+z)} \frac{1}{w+z} dw dz \\ &= \int_0^\infty \int_0^\infty w^{a-1} z^{b-1} e^{-(w+z)} dw dz \\ &= \int_0^\infty w^{a-1} e^{-w} dw \int_0^\infty z^{b-1} e^{-z} dz \\ &= \Gamma(a)\Gamma(b) \end{aligned} \quad (7)$$

Finally the equation is proved.

1.2 Optimization

Choose one problem from the following 1.3 and 1.4. A bonus would be given if you finished the both.

1.3 Stochastic Process

We toss a fair coin for a number of times and use H(head) and T(tail) to denote the two sides of the coin. Please compute the expected number of tosses we need to observe a first time occurrence of the following consecutive pattern

$$H, \underbrace{T, T, \dots, T}_k. \quad (8)$$

Solution: we assume that E is the expectation of the consecutive pattern $H, \underbrace{T, T, \dots, T}_k$, and E_T^k is the expectation of $\underbrace{T, T, \dots, T}_k$, then we have

$$\begin{cases} E = 1 + \frac{1}{2}E + \frac{1}{2}E_T^k, \\ E_T^k = E_T^{k-1} + 1 + \frac{1}{2}E_T^k + \frac{1}{2} \times 0, \end{cases} \quad (9)$$

which

1.4 Probability

Suppose $p \sim \text{Beta}(p|\alpha, \beta)$ and $x|p \sim \text{Bernoulli}(x|p)$. Show that $p|x \sim \text{Beta}(p|\alpha + x, \beta + 1 - x)$, which implies that the Beta distribution can serve as a conjugate prior to the Bernoulli distribution.

Solution: Consider calculating the posterior $p|x$, and we know the likelihood function $x|p$ and the prior p , here we use Bayes' theorem:

$$\begin{aligned} P(p|x) &= \frac{P(x|p)P(p)}{P(x)} \\ &= \frac{P(x|p)P(p)}{\int P(x|p')P(p')dp'} \end{aligned} \quad (10)$$

From the definition, $P(p) \sim \text{Beta}(p|\alpha, \beta)$ and $P(x|p) \sim \text{Bernoulli}(x|p)$, and the Beta function is

$$\text{Beta}(p|\alpha, \beta) = \frac{1}{B(\alpha, \beta)} p^{\alpha-1} (1-p)^{\beta-1} \quad (11)$$

so $P(p|x)$ should be

$$P(p|x) = \frac{P(x|p)P(p)}{\int P(x|p')P(p')dp'} \quad (12)$$

Differentiation allows for the calculation of the slope of the tangent of a curve at any given point as shown in Figure 1.




Figure 1: The plot of $f(x) = 1 - x^2$ with a tangent at $x = .5$.

Differentiation is now a technique taught to mathematics students throughout the world. In this document I will discuss some aspects of differentiation.

2 Exploring the derivative using Sage

The definition of the limit of $f(x)$ at $x = a$ denoted as $f'(a)$ is:

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} \quad (13)$$

The following code can be used in sage to give the above limit:

```
def illustrate(f, a):
    """
    Function to take a function and illustrate the limiting definition of a derivative at a given point
    """
    lst = []
    for h in srange(.01, 3, .01):
        lst.append([h, (f(a+h)-f(a))/h])
    return list_plot(lst, axes_labels=['x$', '$\frac{f(.02f+h)-f(.02f)}{h}$' % (a,a)])
```

Figure 2: The derivative of $f(x) = 1 - x^2$ at $x = .5$ converging to -1 as $h \rightarrow 0$.

If we want to plot the tangent at a point α to a function we can use the following:

$$\begin{array}{ll}
 y = ax + b & \text{(definition of a straight line)} \\
 f'(a)x + b & \text{(definition of the derivative)} \\
 f'(a)x + f(a) - f'(a)a & \text{(we know that the line intersects } f \text{ at } (a, f(a))
 \end{array}$$

We can combine this with the approach of the previous piece of code to see how the tangential line converges as the limiting definition of the derivative converges:

```
def convergetangentialline(f, a, x1, x2, nbrofplots=50, epsilon=.1):
    """
    Function to make a tangential line converge
    """
    clr = rainbow(nbrofplots)
    k = 0
    h = epsilon
    p = plot(f, x, x1, x2)
    while k < nbrofplots:
        tangent(x) = fdash(f, a, h) * x + f(a) - fdash(f, a, h) * a
        p += plot(tangent(x), x, x1, x2, color=clr[k])
        h += epsilon
        k += 1
    return p
```

The plot shown in Figure 3 shows how the lines shown converge to the actual tangent to $1 - x^2$ as $x = 2$ (the red line is the ‘closest’ curve).

Figure 3: Lines converging to the tangent curve as $h \rightarrow 0$.

Note here that the last plot is given using the **real** definition of the derivative and not the approximation.

3 Conclusions

In this report I have explored the limiting definition of the limit showing how as $h \rightarrow 0$ we can visualise the derivative of a function. The code involved <https://sage.maths.cf.ac.uk/home/pub/18/> uses the differentiation capabilities of Sage but also the plotting abilities.

There are various other aspects that could be explored such as symbolic differentiation rules. For example:

$$\frac{dx^n}{dx} = (n+1)x^n \text{ if } x \neq -1$$

Furthermore it is interesting to note that there exists some functions that **are not** differentiable at a point such as the function $f(x) = \sin(1/x)$ which is not differentiable at $x = 0$. A plot of this function is shown in Figure 4.

Figure 4: None differentiable function at $x = 0$.