

## CALCULATION OF LLL BERRY CURVATURE IN SPACE OF METRICS

D.B.

We calculate the adiabatic curvature resulting from varying the LLL wavefunctions with respect to the torus geometry, as parametrized by flat metrics

$$g(V, \tau) = \frac{V}{\tau_2} \begin{pmatrix} 1 & \tau_1 \\ \tau_1 & |\tau|^2 \end{pmatrix}$$

Starting from the Landau level wavefunctions on the torus (ASZ 14), we can calculate the Berry curvature using (ASZ 6),

$$F_{ij} = \text{Im} \sum_{\ell=1}^N \langle \partial_i \phi_\ell | \partial_j \phi_\ell \rangle,$$

where  $\ell$  indexes the single particle wavefunctions, and  $N$  is the number of particles. We will focus on obtaining this curvature for a particular index  $\ell$ . We will find that  $F$  does not depend on  $\ell$ , so the result for multiple particles filling  $N$  levels follows trivially.

The normalized lowest Landau level wavefunction for a given metric  $g(V, \tau)$  is

$$\phi_\ell(\mathbf{x}) = \frac{(2\tau_2 B)^{1/4}}{\sqrt{V}} \sum_{n=-\infty}^{\infty} e^{i\pi\tau B(\tilde{y}+n)^2} e^{-2i\pi(\varphi_2+B/2)(\tilde{y}+n)} e^{2i\pi(nB+\ell)x}$$

with  $\tilde{y} = y + y_\ell = y + \ell/B + \varphi_1/B + 1/2$ .

To compute components of  $F$  involving  $\tau_1, \tau_2$ , we note

$$\phi_\ell(\mathbf{x}) = \frac{(2\tau_2 B)^{1/4}}{V^{1/2}} \Phi_\ell(\mathbf{x})$$

with  $\Phi$  holomorphic in  $\tau$ . Therefore

$$\partial_{\tau_1} \phi_\ell(\mathbf{x}) = \frac{(2\tau_2 B)^{1/4}}{V^{1/2}} \partial_\tau \Phi_\ell(\mathbf{x})$$

and

$$\partial_{\tau_2} \phi_\ell(x, y) = \frac{(2\tau_2 B)^{1/4}}{V^{1/2}} \left[ \frac{1}{4\tau_2} \Phi_\ell(\mathbf{x}) + i \partial_\tau \Phi_\ell(\mathbf{x}) \right].$$

Explicitly,

$$\partial_\tau \Phi_\ell(\mathbf{x}) = \sum_{n=-\infty}^{\infty} \left( i\pi B(\tilde{y} + n)^2 \right) e^{i\pi\tau B(\tilde{y}+n)^2} e^{-2i\pi(\varphi_2+B/2)(\tilde{y}+n)} e^{2i\pi(nB+\ell)x}.$$

For  $F_{V, \tau_2}^\ell$ ,

$$\begin{aligned} F_{V, \tau_2}^\ell &= -F_{\tau_2, V}^\ell = \text{Im} [\langle \partial_V \phi_\ell | \partial_{\tau_2} \phi_\ell \rangle] \\ &= \text{Im} \left[ \int_{\mathbb{Q}} \sqrt{\det g} \, dx \, dy \, \partial_V \phi_\ell^*(\mathbf{x}) \partial_{\tau_2} \phi_\ell(\mathbf{x}) \right] \\ &= \text{Im} \left[ -\frac{1}{2V} \int_{\mathbb{Q}} dx \, dy \, \phi_\ell^*(\mathbf{x}) \left( \frac{1}{4\tau_2} \phi_\ell(\mathbf{x}) + i\sqrt{2\tau_2 B} \partial_\tau \Phi_\ell(\mathbf{x}) \right) \right], \end{aligned}$$

where the integral is over  $Q = [0, 1] \times [0, 1]$ . The first term in the integral is clearly real, so

$$\begin{aligned} F_{V, \tau_2}^\ell &= \frac{\sqrt{2\tau_2 B}}{2V} \text{Im} \left[ i \int_Q dx dy \Phi_\ell^*(\mathbf{x}) \partial_\tau \Phi_\ell(\mathbf{x}) \right] \\ &\equiv \frac{\sqrt{2\tau_2 B}}{2V} \text{Im} [iI_1]. \end{aligned}$$

The integral  $I_1$  is

$$\begin{aligned} I_1 &= \int_Q dx dy \Phi_\ell^*(\mathbf{x}) \partial_\tau \Phi_\ell(\mathbf{x}) = \int_Q dx dy \left[ \sum_{m=-\infty}^{\infty} e^{-i\pi\tau^* B(\tilde{y}+m)^2} e^{2i\pi(\varphi_2+B/2)(\tilde{y}+m)} e^{-2i\pi(mB+\ell)x} \right] \\ &\quad \times \left[ \sum_{n=-\infty}^{\infty} (i\pi B(\tilde{y}+n)^2) e^{i\pi\tau B(\tilde{y}+n)^2} e^{-2i\pi(\varphi_2+B/2)(\tilde{y}+n)} e^{2i\pi(nB+\ell)x} \right] \\ &= \int_Q dx dy \sum_{m,n} i\pi B(\tilde{y}+n)^2 e^{i\pi B(\tau(\tilde{y}+n)^2 - \tau^*(\tilde{y}+m)^2)} e^{2i\pi(\varphi_2+B/2)(m-n)} e^{2i\pi Bx(n-m)} \end{aligned}$$

Performing the  $x$  integration, we have, since  $B$  is an integer

$$\int_0^1 dx e^{2i\pi Bx(n-m)} = \delta_{n,m},$$

and so

$$\begin{aligned} I_1 &= \int_0^1 dy \sum_n i\pi B(\tilde{y}+n)^2 e^{i\pi B(\tau - \tau^*)(\tilde{y}+n)^2} \\ &= i\pi B \sum_{n=-\infty}^{\infty} \int_0^1 dy (\tilde{y}+n)^2 e^{-2\pi B\tau_2(\tilde{y}+n)^2} \\ &= i\pi B \sum_{n=-\infty}^{\infty} \int_{y_\ell+n}^{y_\ell+n+1} du u^2 e^{-2\pi B\tau_2 u^2} \\ &= i\pi B \int_{-\infty}^{\infty} du u^2 e^{-2\pi B\tau_2 u^2} \\ &= i\pi B \left( \frac{1}{4\sqrt{2}\pi(B\tau_2)^{3/2}} \right) \\ &= \frac{i}{4\tau_2\sqrt{2\tau_2 B}} \end{aligned}$$

Since  $I_1$  is pure imaginary,

$$F_{V, \tau_2}^\ell = \frac{\sqrt{2\tau_2 B}}{2V} \text{Im} [iI_1] = 0.$$

For  $F_{V,\tau_1}^\ell$ ,

$$\begin{aligned} F_{V,\tau_1}^\ell &= \text{Im} \left[ -\frac{\sqrt{2\tau_2 B}}{2V} \int_Q dx dy \Phi_\ell^*(\mathbf{x}) \partial_\tau \Phi_\ell(\mathbf{x}) \right] \\ &= -\frac{\sqrt{2\tau_2 B}}{2V} \text{Im} [I_1] \\ &= -\frac{1}{8\tau_2 V} \neq 0 [???] \end{aligned}$$

For  $F_{\tau_1,\tau_2}^\ell$ ,

$$\begin{aligned} F_{\tau_1,\tau_2}^\ell &= -F_{\tau_2,\tau_1}^\ell = \text{Im} [\langle \partial_{\tau_1} \phi_\ell | \partial_{\tau_2} \phi_\ell \rangle] \\ &= \text{Im} \left[ \int_Q \sqrt{\det g} dx dy \partial_{\tau_1} \phi_\ell^*(\mathbf{x}) \partial_{\tau_2} \phi_\ell(\mathbf{x}) \right] \\ &= \text{Im} \left[ \sqrt{2\tau_2 B} \int_Q dx dy \partial_\tau \Phi_\ell^*(\mathbf{x}) \left( \frac{1}{4\tau_2} \Phi_\ell(\mathbf{x}) + i \partial_\tau \Phi_\ell(\mathbf{x}) \right) \right] \\ &= \text{Im} \left[ \sqrt{2\tau_2 B} \left( \frac{1}{4\tau_2} \int_Q dx dy \partial_\tau \Phi_\ell^*(\mathbf{x}) \Phi_\ell(\mathbf{x}) + i \int_Q dx dy \partial_\tau \Phi_\ell^*(\mathbf{x}) \partial_\tau \Phi_\ell(\mathbf{x}) \right) \right] \\ &= \text{Im} \left[ \sqrt{2\tau_2 B} \left( \frac{1}{4\tau_2} I_1^* + i \int_Q dx dy \partial_\tau \Phi_\ell^*(\mathbf{x}) \partial_\tau \Phi_\ell(\mathbf{x}) \right) \right] \\ &\equiv \sqrt{2\tau_2 B} \text{Im} \left[ \frac{1}{4\tau_2} I_1^* + i I_2 \right] \end{aligned}$$

The integral  $I_2$  is

$$\begin{aligned} I_2 &= \int_Q dx dy \partial_\tau \Phi_\ell^*(\mathbf{x}) \partial_\tau \Phi_\ell(\mathbf{x}) \\ &= \int_Q dx dy \sum_{m,n} \left( -i\pi B (\tilde{y} + m)^2 \right) \left( i\pi B (\tilde{y} + n)^2 \right) e^{i\pi B (\tau(\tilde{y}+n)^2 - \tau^*(\tilde{y}+m)^2)} e^{2i\pi(\phi_2+B/2)(m-n)} e^{2i\pi B x(n-m)} \\ &= \pi^2 B^2 \sum_{n=-\infty}^{\infty} \int_0^1 dy (\tilde{y} + n)^4 e^{-2\pi B \tau_2 (\tilde{y}+n)^2} \\ &= \pi^2 B^2 \int_{-\infty}^{\infty} du u^4 e^{-2\pi B \tau_2 (u)^2} \\ &= \pi^2 B^2 \frac{3}{16\sqrt{2}\pi^2 (B\tau_2)^{5/2}} \\ &= \frac{1}{\sqrt{2\tau_2 B}} \frac{3}{16\tau_2^2}. \end{aligned}$$

So that

$$\begin{aligned}
 F_{\tau_1, \tau_2} &= \sqrt{2\tau_2 B} \operatorname{Im} \left[ \frac{1}{4\tau_2} \left( \frac{-i}{4\tau_2 \sqrt{2\tau_2 B}} \right) + i \frac{1}{\sqrt{2\tau_2 B}} \frac{3}{16\tau_2^2} \right] \\
 &= \operatorname{Im} \left[ i \left( -\frac{1}{16\tau_2} + \frac{3}{16\tau_2^2} \right) \right] \\
 &= \frac{1}{8\tau_2^2}
 \end{aligned}$$