

Quartic dispersion

D.B.

Consider the semi-classical energy $E(p_x, p_y)$. If we enforce inversion symmetry, then the most general E we can write up to an additive constant is

$$E(\mathbf{p}) = \sum_{k \text{ even}} E_k(\mathbf{p}),$$

where E_k is a homogeneous polynomial over \mathbb{R} of degree k . Define

$$\varepsilon_k(p_x) = E_k(p_x, 1),$$

then

$$E_k(p_x, p_y) = p_y^k \varepsilon_k(p_x/p_y).$$

ε_k is a polynomial of degree k in one variable over \mathbb{R} , and every polynomial in one variable over \mathbb{R} is reducible to a product of polynomials of degree at most 2. [1] (This corresponds to the fact that we can decompose a polynomial into products of factors for each of its roots, but a real polynomial might have paired complex roots.) Thus there are M possible forms of ε_k ,

$$\varepsilon_k^M(p) = \prod_{i=1}^M (A_2^{M,i} x^2 + A_1^{M,i} x + A_0^{M,i}) \prod_{j=1}^{k-2M} (B_1^{M,j} x + B_0^{M,j})$$

with $0 \leq M \leq \lceil k/2 \rceil$. For $k = 4$, there are 3 possibilities for ε_4 : 2 quadratic factors, 1 quadratic factor and 2 linear factors, or 4 linear factors.

REFERENCES

[1] https://en.wikipedia.org/wiki/Irreducible_polynomial