

## Greek Characters

Name	Symbol	Typical use(s)
alpha	$\alpha$	angle, constant
beta	$\beta$	angle, constant
gamma	$\gamma$	angle, constant
delta	$\delta$	limit definition
epsilon	$\epsilon$ or $\varepsilon$	limit definition
theta	$\theta$ or $\vartheta$	angle
pi	$\pi$ or $\pi$	circular constant
phi	$\phi$ or $\varphi$	angle, constant

## Named Sets

empty set	$\emptyset$
real numbers	$\mathbf{R}$
ordered pairs	$\mathbf{R}^2$
integers	$\mathbf{Z}$
positive integers	$\mathbf{Z}_{>0}$
positive real numbers	$\mathbf{R}_{>0}$

## Set Symbols

Meaning	Symbol
is a member	$\in$
subset	$\subset$
intersection	$\cap$
union	$\cup$
set minus	$\setminus$

## Intervals

For numbers  $a$  and  $b$ , we define the intervals:

$$(a, b) = \{x \in \mathbf{R} \mid a < x < b\}$$

$$[a, b) = \{x \in \mathbf{R} \mid a \leq x < b\}$$

$$(a, b] = \{x \in \mathbf{R} \mid a < x \leq b\}$$

$$[a, b] = \{x \in \mathbf{R} \mid a \leq x \leq b\}$$

## Logic Symbols

Meaning	Symbol
negation	$\neg$
and	$\wedge$
or	$\vee$
implies	$\implies$
equivalent	$\equiv$
for all	$\forall$
there exists	$\exists$

## Exponents

For  $a, b > 0$  and  $m, n$  real:

$$a^0 = 1, \quad 0^a = 0$$

$$1^a = 1, \quad a^n a^m = a^{n+m}$$

$$a^n / a^m = a^{n-m}, \quad (a^n)^m = a^{n \cdot m}$$

$$a^{-m} = 1/a^m, \quad (a/b)^m = a^m / b^m$$

## Trigonometric Identities

$$\sin(x)^2 + \cos(x)^2 = 1$$

$$2 \cos(x)^2 = 1 + \cos(2x)$$

$$2 \sin(x)^2 = 1 - \cos(2x)$$

$$\sin(x+y) = \sin(x) \cos(y) + \cos(x) \sin(y)$$

$$\cos(x+y) = \cos(x) \cos(y) - \sin(x) \sin(y)$$

$$\operatorname{arccot}(x) = \arctan(1/x)$$

$$\operatorname{arccsc}(x) = \arcsin(1/x)$$

$$\operatorname{arcsec}(x) = \arccos(1/x)$$

## Limits

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1 \quad \lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x} = 0$$

$$\lim_{x \rightarrow -\infty} e^x = 0 \quad \lim_{x \rightarrow -\infty} e^x = 0$$

$$\lim_{x \rightarrow -\infty} \ln(x) = -\infty \quad \lim_{x \rightarrow 0^+} \ln(x) = -\infty$$

## Derivatives

### Specific cases

$F(x)$	$F'(x)$
$\cos(x)$	$-\sin(x)$
$\sin(x)$	$\cos(x)$
$\tan(x)$	$\sec(x)^2$
$\sec(x)$	$\sec(x) \tan(x)$
$\csc(x)$	$-\cot(x) \csc(x)$
$\cot(x)$	$-\csc(x)^2$
$\arccos(x)$	$-1/\sqrt{1-x^2}$
$\arcsin(x)$	$1/\sqrt{1-x^2}$
$\arctan(x)$	$1/(x^2+1)$
$\exp(x)$	$\exp(x)$
$\ln(x)$	$1/x$

### General Cases

$F(x)$	$F'(x)$
$af(x) + bg(x)$	$af'(x) + bg'(x)$
$f(x)g(x)$	$f'(x)g(x) + f(x)g'(x)$
$1/g(x)$	$-g'(x)/g(x)^2$
$f(x)/g(x)$	$(g(x)f'(x) - f(x)g'(x))/g(x)^2$
$f(g(x))$	$g'(x)f'(g(x))$
$f^{-1}(x)$	$1/f'(f^{-1}(x))$

## Antiderivatives

$$\int a \, dx = ax$$

$$\int x^a \, dx = \frac{1}{1+a} x^{a+1}, \quad \text{if } a \neq -1$$

$$\int \frac{1}{x} \, dx = \ln|x|$$

$$\int \cos(x) \, dx = \sin(x)$$

$$\int \sin(x) \, dx = -\cos(x)$$

$$\int \tan(x) \, dx = \ln|\sec(x)|$$

$$\int \sec(x) \, dx = \ln|\tan(x) + \sec(x)|$$

$$\int \csc(x) \, dx = -\ln|\csc(x) + \cot(x)|$$

$$\int \cot(x) \, dx = \ln|\sin(x)|$$

$$\int 2|x| \, dx = x|x|$$

$$\int 2[x] \, dx = (2x-1)[x] - [x]^2$$

$$\int 2\lceil x \rceil \, dx = (2x+1)\lceil x \rceil - \lceil x \rceil^2$$

## Sums

For  $k, m, n \in \mathbf{Z}_{>0}$

$$\sum_{k=0}^{n-1} 1 = n$$

$$\sum_{k=0}^{n-1} k = \frac{(n-1)n}{2}$$

$$\sum_{k=0}^{n-1} k^2 = \frac{(n-1)n(2n-1)}{6}$$


$$\sum_{k=0}^{n-1} x^k = \frac{1-x^n}{1-x}, \quad x \neq 1$$

## Logarithms


$$\log_a(x) = \frac{1}{\ln(a)} \ln(x)$$

## Famous Triangles


### The 30-60-90 triangle



## The 45-45-90 triangle



## Laws of Cosine & Sine




**Law of cosine:**  $C^2 = A^2 + B^2 - 2AB \cos(\gamma)$

**Law of sines:**  $\frac{\sin \alpha}{A} = \frac{\sin \beta}{B} = \frac{\sin \gamma}{C}$

**Area:**  $\text{Area} = \frac{1}{2}hB = \frac{1}{2}AB \sin(\gamma)$

## Volumes

### Right Circular Cylinder




**Volume:**  $V = \pi r^2 h$

**Area:** (not including circular ends)

$$A = 2\pi r h$$

### Cone



**Volume:**  $V = \frac{1}{3}\pi r^2 h$

**Area** (not including circular base)


$$A = \pi r \left( r + \sqrt{r^2 + h^2} \right)$$

### Sphere

**Area:**  $A = 4\pi r^2$

**Volume:**  $V = \frac{4\pi}{3}r^3$

## Unit Circle



## Applications

Arclength of curve  $y = f(x)$  with  $a \leq x \leq b$

$$= \int_a^b \sqrt{1 + f'(x)^2} dx$$

For the region  $Q$  of the  $xy$  plane given by

$$Q = \{(x, y) \mid f(x) \leq y \leq g(x) \wedge a \leq x \leq b\},$$

we have

$$\text{Area}(Q) = \int_a^b g(x) - f(x) dx$$

Assuming  $0 \leq f(x)$  and rotating about the  $x$ -axis

$$\text{Vol}(Q) = \pi \int_a^b g(x)^2 - f(x)^2 dx$$

Assuming  $0 \leq a < b$  and rotating about the  $y$ -axis

$$\text{Vol}(Q) = 2\pi \int_a^b x(g(x) - f(x)) dx$$

Centroid


$$\text{Area}(Q) \times \bar{x} = \int_a^b x(g(x) - f(x)) dx$$

$$\text{Area}(Q) \times \bar{y} = \frac{1}{2} \int_a^b (g(x)^2 - f(x)^2) dx$$

For the region  $Q$  of the  $xy$  plane given by

$$Q = \{(x, y) \mid f(y) \leq x \leq g(y) \wedge a \leq y \leq b\},$$

interchange  $x$  and  $y$  in *all* the previous formulas.

Revised August 5, 2022. Barton Willis is the author of this work. This work is licensed under Attribution 4.0 International (CC BY 4.0)  For the current version of this document, visit 