

Greek Characters

| Name | Symbol | Typical use(s) |
|---------|-----------------------------|-------------------|
| alpha | α | angle, constant |
| beta | β | angle, constant |
| gamma | γ | angle, constant |
| delta | δ | limit definition |
| epsilon | ϵ or ε | limit definition |
| theta | θ or ϑ | angle |
| pi | π or π | circular constant |
| phi | ϕ or φ | angle, constant |

Named Sets

| | |
|-----------------------|-------------------|
| empty set | \emptyset |
| real numbers | \mathbf{R} |
| ordered pairs | \mathbf{R}^2 |
| integers | \mathbf{Z} |
| positive integers | $\mathbf{Z}_{>0}$ |
| positive real numbers | $\mathbf{R}_{>0}$ |

Set Symbols

| Meaning | Symbol |
|--------------|-------------|
| is a member | \in |
| subset | \subset |
| intersection | \cap |
| union | \cup |
| set minus | \setminus |

Intervals

For numbers a and b , we define the intervals:

$$(a, b) = \{x \in \mathbf{R} \mid a < x < b\}$$

$$[a, b) = \{x \in \mathbf{R} \mid a \leq x < b\}$$

$$(a, b] = \{x \in \mathbf{R} \mid a < x \leq b\}$$

$$[a, b] = \{x \in \mathbf{R} \mid a \leq x \leq b\}$$

Logic Symbols

| Meaning | Symbol |
|--------------|-------------------|
| negation | \neg |
| and | \wedge |
| or | \vee |
| implies | \Rightarrow |
| equivalent | \equiv |
| iff | \Leftrightarrow |
| for all | \forall |
| there exists | \exists |

Exponents

For $a, b > 0$, $x \in \mathbf{R}$, and m, n real:

$$a^0 = 1, \quad 0^a = 0$$

$$1^a = 1, \quad a^n a^m = a^{n+m}$$

$$a^n / a^m = a^{n-m}, \quad (a^n)^m = a^{n \cdot m}$$

$$a^{-m} = 1/a^m, \quad (a/b)^m = a^m / b^m$$

$$\sqrt{x^2} = |x|$$

Trigonometric Identities

$$\cos(x)^2 + \sin(x)^2 = 1$$

$$2 \cos(x)^2 = 1 + \cos(2x)$$

$$2 \sin(x)^2 = 1 - \cos(2x)$$

$$\sin(x+y) = \sin(x) \cos(y) + \cos(x) \sin(y)$$

$$\cos(x+y) = \cos(x) \cos(y) - \sin(x) \sin(y)$$

$$\operatorname{arccot}(x) = \pi/2 - \arctan(x)$$

$$\operatorname{arccsc}(x) = \pi/2 + \arcsin(1/x)$$

$$\operatorname{arcsec}(x) = \arccos(1/x)$$

Limits

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1 \quad \lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x} = 0$$

$$\lim_{x \rightarrow \infty} e^x = \infty \quad \lim_{x \rightarrow -\infty} e^x = 0$$

$$\lim_{x \rightarrow \infty} \ln(x) = \infty \quad \lim_{x \rightarrow 0^+} \ln(x) = -\infty$$

Derivatives

Specific cases

| $F(x)$ | $F'(x)$ |
|-----------------------------|--------------------|
| $\cos(x)$ | $-\sin(x)$ |
| $\sin(x)$ | $\cos(x)$ |
| $\tan(x)$ | $\sec(x)^2$ |
| $\sec(x)$ | $\sec(x) \tan(x)$ |
| $\csc(x)$ | $-\cot(x) \csc(x)$ |
| $\cot(x)$ | $-\csc(x)^2$ |
| $\arccos(x)$ | $-1/\sqrt{1-x^2}$ |
| $\arcsin(x)$ | $1/\sqrt{1-x^2}$ |
| $\arctan(x)$ | $1/(x^2+1)$ |
| $\cosh(x)$ | $\sinh(x)$ |
| $\sinh(x)$ | $\cosh(x)$ |
| $\tanh(x)$ | $1/\cosh(x)^2$ |
| $\operatorname{arccosh}(x)$ | $1/\sqrt{x^2-1}$ |
| $\operatorname{arsinh}(x)$ | $1/\sqrt{1+x^2}$ |
| $\operatorname{arctanh}(x)$ | $1/(1-x^2)$ |
| $\exp(x)$ | $\exp(x)$ |
| $\ln(x)$ | $1/x$ |

General Cases

| $F(x)$ | $F'(x)$ |
|-----------------|----------------------------------|
| $af(x) + bg(x)$ | $af'(x) + bg'(x)$ |
| $f(x)g(x)$ | $f'(x)g(x) + f(x)g'(x)$ |
| $1/g(x)$ | $-g'(x)/g(x)^2$ |
| $f(x)/g(x)$ | $(g(x)f'(x) - f(x)g'(x))/g(x)^2$ |
| $f(g(x))$ | $g'(x)f'(g(x))$ |
| $f^{-1}(x)$ | $1/f'(f^{-1}(x))$ |

Antiderivatives

$$\int a \, dx = ax$$

$$\int x^a \, dx = \frac{1}{1+a} x^{a+1}, \quad \text{if } a \neq -1$$

$$\int \frac{1}{x} \, dx = \ln|x|$$

$$\int \cos(x) \, dx = \sin(x)$$

$$\int \sin(x) \, dx = -\cos(x)$$

$$\int \tan(x) \, dx = \ln|\sec(x)|$$

$$\int \sec(x) \, dx = \ln|\tan(x) + \sec(x)|$$

$$\int \csc(x) \, dx = -\ln|\csc(x) + \cot(x)|$$

$$\int \cot(x) \, dx = \ln|\sin(x)|$$

$$\int 2|x| \, dx = x|x|$$

Sums

For $k, m, n \in \mathbf{Z}_{>0}$

$$\sum_{k=0}^{n-1} 1 = n$$

$$\sum_{k=0}^{n-1} k = \frac{(n-1)n}{2}$$

$$\sum_{k=0}^{n-1} k^2 = \frac{(n-1)n(2n-1)}{6}$$


$$\sum_{k=0}^{n-1} x^k = \frac{1-x^n}{1-x}, \quad x \neq 1$$

Logarithms


$$\log_a(x) = \frac{\ln(x)}{\ln(a)}$$

Famous Triangles


The 30-60-90 triangle



The 45-45-90 triangle



Laws of Cosine & Sine



Law of cosine: $C^2 = A^2 + B^2 - 2AB \cos(\gamma)$

Law of sines: $\frac{\sin(\alpha)}{A} = \frac{\sin(\beta)}{B} = \frac{\sin(\gamma)}{C}$

Area: $\text{Area} = \frac{1}{2}hB = \frac{1}{2}AB \sin(\gamma)$

Hyperbolic Functions

$$2 \cosh(x) = \exp(x) + \exp(-x)$$


$$2 \sinh(x) = \exp(x) - \exp(-x)$$

$$\tanh(x) = \cosh(x)/\sinh(x)$$

$$\cosh(x)^2 - \sinh(x)^2 = 1$$

Volumes

Right Circular Cylinder




Volume: $V = \pi r^2 h$

Area: (not including circular ends)

$$A = 2\pi r h$$

Cone



Volume: $V = \frac{1}{3}\pi r^2 h$

Area (not including circular base)


$$A = \pi r \sqrt{r^2 + h^2}$$

Sphere

Area: $A = 4\pi r^2$

Volume: $V = \frac{4\pi}{3}r^3$

Unit Circle



Applications

Arclength of curve $y = f(x)$ with $a \leq x \leq b$

$$= \int_a^b \sqrt{1 + f'(x)^2} dx$$

For the region Q of the xy plane given by

$$Q = \{(x, y) \mid f(x) \leq y \leq g(x) \wedge a \leq x \leq b\},$$

we have

$$\text{Area}(Q) = \int_a^b g(x) - f(x) dx$$

Assuming $0 \leq f(x)$ and rotating about the x -axis

$$\text{Vol}(Q) = \pi \int_a^b g(x)^2 - f(x)^2 dx$$

Assuming $0 \leq a < b$ and rotating about the y -axis

$$\text{Vol}(Q) = 2\pi \int_a^b x(g(x) - f(x)) dx$$

Centroid

$$\text{Area}(Q) \times \bar{x} = \int_a^b x(g(x) - f(x)) dx$$

$$\text{Area}(Q) \times \bar{y} = \frac{1}{2} \int_a^b (g(x)^2 - f(x)^2) dx$$

For the region Q of the xy plane given by

$$Q = \{(x, y) \mid f(y) \leq x \leq g(y) \wedge a \leq y \leq b\},$$

interchange x and y in *all* the previous formulas.

Revised February 18, 2023. Barton Willis is the author of this work. This work is licensed under Attribution 4.0 International (CC BY 4.0) For the current version of this document, visit