

MATH 365
Review for Exam 3

Name: _____
Row and Seat: _____

1. Find the radius of convergence for each series

10 (a) $\sum_{k=0}^{\infty} \frac{1}{1+k^2} x^k$

10 (b) $\sum_{k=0}^{\infty} \frac{(k!)^2}{(3k)!} x^k$

10 2. Use the bound

$$\left| \int_{\gamma} F(z) \, dz \right| \leq \max_{z \in \gamma} |F(z)| \times \text{length}(\gamma)$$

to find an *upper bound* for

$$\int_L \frac{1}{x-i} \, dx,$$

where L is the line segment from $-1 - 5i$ to $1 + 5i$.

3. Write a critique of Billy's work

$$\begin{aligned}\int_0^{2\pi} \frac{1}{4 + \cos(x)} dx &= \frac{2 \arctan\left(\frac{3 \sin(x)}{\sqrt{15}(\cos(x)+1)}\right)}{\sqrt{15}} \bigg|_{x=0}^{x=2\pi} \\ &= 0 - 0 \\ &= 0.\end{aligned}$$

Fact: Everywhere $\frac{2 \arctan\left(\frac{3 \sin(x)}{\sqrt{15}(\cos(x)+1)}\right)}{\sqrt{15}}$ is differentiable, its derivative is the integrand. Here is a graph of the antiderivative.




Figure 1: Graph of antiderivative on the interval $[0, 2\pi]$.

4. Find the *numerical value* of each sum.

10

(a) $\sum_{k=0}^{\infty} \left(\frac{2}{3}\right)^{k+1}$

$$\boxed{10} \quad (b) \sum_{k=2}^{\infty} \left(\frac{2}{3}\right)^{k+1}$$

5. Find the *numerical value* for each contour integral, where C is the unit circle.

$$\boxed{10} \quad (a) \int_C \frac{1}{z-5i} dz.$$

$$\boxed{10} \quad (\text{b}) \int_C \frac{1}{z - i/5} dz.$$

$\boxed{10}$ 6. For $w > 0$, find the value of

$$\int_{-\infty}^{\infty} \frac{1}{4 + x^2} e^{iwx} dx. \tag{1}$$

- 10 7. My friend Pauline claims that since the function $x \in \mathbf{C} \mapsto e^{e^{ix}}$ is entire, we have

$$\int_0^{2\pi} e^{e^{ix}} dx = 0.$$

Pauline is correct that the function $x \mapsto e^{e^{ix}}$ is entire. But is Pauline correct about the value of the integral? If not, find the correct value for this definite integral.