

Can I do X?

MATH 202

April 15, 2024

"The law is reason unaffected by desire."

ARISTOTLE

Our problem

We would like to find an explicit formula for the sequence c that is defined recursively by

$$c_n = \begin{cases} 0 & n = 0 \text{ or } n = 1 \\ c_{n-1} + c_{n-2} + 8 & n \in \mathbf{Z}_{\geq 2} \end{cases} \quad (1)$$

👉 The first eleven terms of the sequence c are

$$0, 0, 8, 16, 32, 56, 96, 160, 264, 432, 704 \quad (2)$$

👉 To complete this task, we need three new tools

Tool 1: Binomial Coefficients

For positive integers n and k with $n \geq k$, we define

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} \quad (3)$$

- 👉 Spoken $\binom{n}{k}$ is “ n choose k .”
- 👉 For a finite set A with n members, $\binom{n}{k}$ is the number of subsets of A that have k members.

Tool 2: Product rule for n -th derivatives

For smooth functions f and g and a nonnegative integer n , the n -th derivative of the product of f with g is

$$D^n(fg) = \sum_{k=0}^n \binom{n}{k} f^k g^{n-k}. \quad (4)$$

Tool 3: n -th derivative of a power series

Let c be a sequence and let n be a nonnegative integer. Then

$$D_x^n \sum_{k=0}^{\infty} c_k x^k \Big|_{x=0} = n! c_n. \quad (5)$$

👉 For this to be true, it has to be the case that the radius of convergence of the series is nonzero.

Tool 4: n -th derivative of a rational function

Let $a \in \mathbf{R}$ and n a nonnegative integer. Then

$$D_x^n \left(\frac{1}{x-a} \right) \Big|_{x=0} = -\frac{n!}{a^{n+1}}. \quad (6)$$

Step 0

Multiply the recursion $c_n = c_{n-1} + c_{n-2} + 8$ by z^n and sum from $n = 2$ to ∞ . This gives

$$\sum_{n=2}^{\infty} c_n z^n = \sum_{n=2}^{\infty} c_{n-1} z^n + \sum_{n=2}^{\infty} c_{n-2} z^n + \sum_{n=2}^{\infty} 8z^n \quad (7)$$

👉 The lowest sum index is two because the recursion $c_n = c_{n-1} + c_{n-2} + 8$ is only valid for $n \geq 2$.

Step 1

If needed shift each sum index to make the each summand involve c_k .

$$\sum_{n=2}^{\infty} c_n z^n = \sum_{n=1}^{\infty} c_n z^{n+1} + \sum_{n=0}^{\infty} c_n z^{n+2} + \sum_{n=2}^{\infty} 8z^n \quad (8)$$

👉 The lowest sum index is two because the recursion $c_n = c_{n-1} + c_{n-2} + 8$ is only valid for $n \geq 2$.

Step 2

Use the know values of c_0 and c_1 to extend the lower sum index of each sum to zero.

$$\sum_{n=0}^{\infty} c_n z^n = z \sum_{n=0}^{\infty} c_n z^n + z^2 \sum_{n=0}^{\infty} c_n z^n + \sum_{n=2}^{\infty} 8z^n \quad (9)$$

👉 If c_0 and c_1 were nonzero, we'd have a few more terms!

Step 3

Define $G(z) = \sum_{n=0}^{\infty} c_n z^n$ and $F(x) = \sum_{n=2}^{\infty} 8z^n$. We have

$$G(z) = zG(z) + z^2G(z) + F(z) \quad (10)$$

So

$$G(z) = \frac{1}{1 - z - z^2} F(x). \quad (11)$$

Step 3

From G , determine c_k . We have

$$\begin{aligned} c_n &= \frac{1}{n!} D_z^n \frac{1}{1-z-z^2} F(x)|_{x=0}, \\ &= \frac{1}{n!} \sum_{k=0}^n \binom{n}{k} D_z^k \left(\frac{1}{1-z-z^2} \right) D_z^{n-k} F(z)|_{z=0} = \end{aligned}$$