

## Greek Characters

| Name    | Symbol                      | Typical use(s)    |
|---------|-----------------------------|-------------------|
| alpha   | $\alpha$                    | angle, constant   |
| beta    | $\beta$                     | angle, constant   |
| gamma   | $\gamma$                    | angle, constant   |
| delta   | $\delta$                    | limit definition  |
| epsilon | $\epsilon$ or $\varepsilon$ | limit definition  |
| theta   | $\theta$ or $\vartheta$     | angle             |
| pi      | $\pi$ or $\pi$              | circular constant |
| phi     | $\phi$ or $\varphi$         | angle, constant   |

## Named Sets

|                       |                   |
|-----------------------|-------------------|
| empty set             | $\emptyset$       |
| real numbers          | $\mathbf{R}$      |
| ordered pairs         | $\mathbf{R}^2$    |
| integers              | $\mathbf{Z}$      |
| positive integers     | $\mathbf{Z}_{>0}$ |
| positive real numbers | $\mathbf{R}_{>0}$ |

## Set Symbols

| Meaning      | Symbol      |
|--------------|-------------|
| is a member  | $\in$       |
| subset       | $\subset$   |
| intersection | $\cap$      |
| union        | $\cup$      |
| set minus    | $\setminus$ |

## Intervals

For numbers  $a$  and  $b$ , we define the intervals:

$$(a, b) = \{x \in \mathbf{R} \mid a < x < b\}$$

$$[a, b) = \{x \in \mathbf{R} \mid a \leq x < b\}$$

$$(a, b] = \{x \in \mathbf{R} \mid a < x \leq b\}$$

$$[a, b] = \{x \in \mathbf{R} \mid a \leq x \leq b\}$$

## Logic Symbols

| Meaning      | Symbol     |
|--------------|------------|
| negation     | $\neg$     |
| and          | $\wedge$   |
| or           | $\vee$     |
| implies      | $\implies$ |
| equivalent   | $\equiv$   |
| for all      | $\forall$  |
| there exists | $\exists$  |

## Exponents

For  $a, b > 0$  and  $m, n$  real:

$$a^0 = 1, \quad 0^a = 0$$

$$1^a = 1, \quad a^n a^m = a^{n+m}$$

$$a^n / a^m = a^{n-m}, \quad (a^n)^m = a^{n \cdot m}$$

$$a^{-m} = 1/a^m, \quad (a/b)^m = a^m / b^m$$

## Trigonometric Identities

$$\sin(x)^2 + \cos(x)^2 = 1$$

$$2 \cos(x)^2 = 1 + \cos(2x)$$

$$2 \sin(x)^2 = 1 - \cos(2x)$$

$$\sin(x+y) = \sin(x) \cos(y) + \cos(x) \sin(y)$$

$$\cos(x+y) = \cos(x) \cos(y) - \sin(x) \sin(y)$$

$$\operatorname{arccot}(x) = \arctan(1/x)$$

$$\operatorname{arccsc}(x) = \arcsin(1/x)$$

$$\operatorname{arcsec}(x) = \arccos(1/x)$$

## Limits

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1 \quad \lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x} = 0$$

$$\lim_{x \rightarrow -\infty} e^x = 0 \quad \lim_{x \rightarrow -\infty} e^{-x} = 0$$

$$\lim_{x \rightarrow -\infty} \ln(x) = -\infty \quad \lim_{x \rightarrow 0^+} \ln(x) = -\infty$$

## Derivatives

### Specific cases

| $F(x)$       | $F'(x)$            |
|--------------|--------------------|
| $\cos(x)$    | $-\sin(x)$         |
| $\sin(x)$    | $\cos(x)$          |
| $\tan(x)$    | $\sec(x)^2$        |
| $\sec(x)$    | $\sec(x) \tan(x)$  |
| $\csc(x)$    | $-\cot(x) \csc(x)$ |
| $\cot(x)$    | $-\csc(x)^2$       |
| $\arccos(x)$ | $-1/\sqrt{1-x^2}$  |
| $\arcsin(x)$ | $1/\sqrt{1-x^2}$   |
| $\arctan(x)$ | $1/(x^2+1)$        |
| $\exp(x)$    | $\exp(x)$          |
| $\ln(x)$     | $1/x$              |

### General Cases

| $F(x)$          | $F'(x)$                          |
|-----------------|----------------------------------|
| $af(x) + bg(x)$ | $af'(x) + bg'(x)$                |
| $f(x)g(x)$      | $f'(x)g(x) + f(x)g'(x)$          |
| $1/g(x)$        | $-g'(x)/g(x)^2$                  |
| $f(x)/g(x)$     | $(g(x)f'(x) - f(x)g'(x))/g(x)^2$ |
| $f(g(x))$       | $g'(x)f'(g(x))$                  |
| $f^{-1}(x)$     | $1/f'(f^{-1}(x))$                |

## Antiderivatives

$$\int a \, dx = ax$$

$$\int x^a \, dx = \frac{1}{1+a} x^{a+1}, \quad \text{if } a \neq -1$$

$$\int \frac{1}{x} \, dx = \ln|x|$$

$$\int \cos(x) \, dx = \sin(x)$$

$$\int \sin(x) \, dx = -\cos(x)$$

$$\int \tan(x) \, dx = \ln|\sec(x)|$$

$$\int \sec(x) \, dx = \ln|\tan(x) + \sec(x)|$$

$$\int \csc(x) \, dx = -\ln|\csc(x) + \cot(x)|$$

$$\int \cot(x) \, dx = \ln|\sin(x)|$$

$$\int 2|x| \, dx = x|x|$$

$$\int 2[x] \, dx = (2x-1)[x] - [x]^2$$

$$\int 2\lceil x \rceil \, dx = (2x+1)\lceil x \rceil - \lceil x \rceil^2$$

## Sums

For  $k, m, n \in \mathbf{Z}_{>0}$

$$\sum_{k=0}^{n-1} 1 = n$$

$$\sum_{k=0}^{n-1} k = \frac{(n-1)n}{2}$$

$$\sum_{k=0}^{n-1} k^2 = \frac{(n-1)n(2n-1)}{6}$$


$$\sum_{k=0}^{n-1} x^k = \frac{1-x^n}{1-x}, \quad x \neq 1$$

## Logarithms


$$\log_a(x) = \frac{1}{\ln(a)} \ln(x)$$

## Famous Triangles


### The 30-60-90 triangle



## The 45-45-90 triangle



## Laws of Cosine & Sine




**Law of cosine:**  $C^2 = A^2 + B^2 - 2AB \cos(\gamma)$

**Law of sines:**  $\frac{\sin \alpha}{A} = \frac{\sin \beta}{B} = \frac{\sin \gamma}{C}$

**Area:**  $\text{Area} = \frac{1}{2}hB = \frac{1}{2}AB \sin(\gamma)$

## Volumes

### Right Circular Cylinder




**Volume:**  $V = \pi r^2 h$

**Area:** (not including circular ends)

$$A = 2\pi r h$$

### Cone



**Volume:**  $V = \frac{1}{3}\pi r^2 h$

**Area** (not including circular base)


$$A = \pi r \left( r + \sqrt{r^2 + h^2} \right)$$

### Sphere

**Area:**  $A = 4\pi r^2$

**Volume:**  $V = \frac{4\pi}{3}r^3$

## Unit Circle



## Applications

Arclength of curve  $y = f(x)$  with  $a \leq x \leq b$

$$= \int_a^b \sqrt{1 + f'(x)^2} dx$$

For the region  $Q$  of the  $xy$  plane given by

$$Q = \{(x, y) \mid f(x) \leq y \leq g(x) \wedge a \leq x \leq b\},$$

we have

$$\text{Area}(Q) = \int_a^b g(x) - f(x) dx$$

Assuming  $0 \leq f(x)$  and rotating about the  $x$ -axis

$$\text{Vol}(Q) = \pi \int_a^b g(x)^2 - f(x)^2 dx$$

Assuming  $0 \leq a < b$  and rotating about the  $y$ -axis

$$\text{Vol}(Q) = 2\pi \int_a^b x(g(x) - f(x)) dx$$

Centroid


$$\text{Area}(Q) \times \bar{x} = \int_a^b x(g(x) - f(x)) dx$$

$$\text{Area}(Q) \times \bar{y} = \frac{1}{2} \int_a^b (g(x)^2 - f(x)^2) dx$$

For the region  $Q$  of the  $xy$  plane given by

$$Q = \{(x, y) \mid f(y) \leq x \leq g(y) \wedge a \leq y \leq b\},$$

interchange  $x$  and  $y$  in *all* the previous formulas.

Revised May 5, 2022. Barton Willis is the author of this work. This work is licensed under Attribution 4.0 International (CC BY 4.0)  For the current version of this document, visit 