

Derivative divides integration

AKA the U substitution

Chain rule swapperoo

Recall the chain rule; for differentiable functions F and g in function notation, we have

$$(F \circ g)' = g' F' \circ g.$$

Equivalently in formula notation, the chain rule is

$$\frac{d}{dx}(F(g(x))) = g'(x)F'(g(x)).$$

Fact

Up to an additive constant, the antiderivative “undoes” the derivative; thus

$$\int g'(x)F'(g(x)) \, dx = F(g(x)) + c,$$

where $c \in \mathbf{R}$.

Match this

If we can match an integrand to $g'(x)F'(g(x))$ and we know an antiderivative of F , **we win**. An example

$$\int 2x \sin(x^2) \, dx.$$

The match isn't particularly hidden:

- ✓ $g(x) = x^2 \implies g'(x) = 2x,$
- ✓ $F' = \sin \implies F = -\cos.$

Thus

$$\int 2x \sin(x^2) \, dx = F(g(x)) + c = -\cos(x^2) + c.$$

Out with the old, in with the new

Let's re-do the problem $\int 2x \sin(x^2) dx$, but organize our work differently.

- ✓ The argument of \sin is x^2 . Let's define a new variable $u = x^2$. Then

$$du = \frac{du}{dx} dx = 2x dx.$$

- ✓ We now need to write $2x \sin(x^2) dx$ *entirely* in terms of the new variable u .
- ✓ When I say entirely, I mean *entirely*. This includes expressing dx in terms of du .
- ✓ Grouping the factor of $2x$ together with dx , we have

$$2x \sin(x^2) dx = \sin(x^2)(2x dx) = \sin(u) du.$$

- ✓ So

$$\int 2x \sin(x^2) dx = \int \sin(u) du = -\cos(u) = -\cos(x^2).$$

- ✓ For the step $-\cos(u) = -\cos(x^2)$ we reverted to the “original” integration variable.

Example redux

Let's try $\int x \exp(x^2) dx$

- ✓ The argument of \exp is x^2 . Let's define a new variable $u = x^2$. Then

$$du = \frac{du}{dx} dx = 2x dx.$$

- ✓ Unlike the previous problem, the integrand is missing a factor of 2 for a complete matching. No big deal; we have

$$[du = 2x dx] = \left[\frac{1}{2} du = x dx \right]$$

- ✓ So

$$x \exp(x^2) dx = \exp(x^2)(x dx) = \frac{1}{2} \exp(u) du.$$

- ✓ So

$$\int x \exp(x^2) dx = \int \frac{1}{2} \exp(u) du = \frac{1}{2} \exp(u) = \frac{1}{2} \exp(x^2).$$

- ✓ In step $\exp(u) = \exp(x^2)$ we reverted to the “original” integration variable.

Derivative divides

- ✓ When we match $u = x^2$ to find $\int x \exp(x^2) dx$, the derivative of u , that is $\frac{du}{dx}$, doesn't exactly match the remaining factor of x , but it does match up to a multiplicative factor of 2.
- ✓ Since the quotient of $\frac{du}{dx}$ divided by the the remaining factor of x is a constant, the method is called *derivative divides*.
- ✓ But calculus books call the method *U substitution*
- ✓ If you don't like my explanations, no problem; watch <https://www.youtube.com/watch?v=8B31SAk1nD8>.

Finding You

Heuristic

- ✓ The integrand should be a product.
- ✓ Choose u to be an expression that is “inside” a function with a known antiderivative.
- ✓ The derivative of u times a constant should match the remaining factors of the integrand.
- ✓ A heuristic is a guiding principle that often works, but occasionally fails.

Matching examples

$\int x^2 \cos(x^3) dx$. The cosine has a known antiderivative. The expression inside cosine is x^3 . So choose $u = x^3$. Then $\frac{du}{dx} = 3x^2$ matches the remaining factor x^2 in the integrand up to a multiplicative factor. **We win.**

$$[u = x^3] = [du = 3x^2 dx] = \left[x^2 dx = \frac{1}{3} du \right].$$

So

$$\int x^2 \cos(x^3) dx = \int \frac{1}{3} \cos(u) du = \frac{1}{3} \sin(u) = \frac{1}{3} \sin(x^3).$$

Actually, we are losers

Actually, we are **losers**. Why? Because we didn't check our work:

$$\frac{d}{dx} \left(\frac{1}{3} \sin(x^3) \right) = \frac{1}{3} 3x^2 \cos(x^3) = x^2 \cos(x^3)$$

The integrand is $x^2 \cos(x^3)$. So our answer is OK.

- ✔ Anytime we fail to check our work, we **are losers**.
- ✔ *When you check your work, you only cry once. But when we don't, we cry many times.*

Example

$\int x \exp(-x^2) dx$. The function \exp has a known antiderivative. The expression inside \exp is $-x^2$. So choose $u = -x^2$. Then $\frac{du}{dx} = -2x$ matches the remaining factor x in the integrand up to a multiplicative factor. **We win.**

$$[u = -x^2] = [du = -2x dx] = \left[x dx = -\frac{1}{2} du \right].$$

So

$$\int x \exp(-x^2) dx = \int -\frac{1}{2} \exp(u) du = -\frac{1}{2} \exp(u) = -\frac{1}{2} \exp(-x^2).$$

Example

$\int \sqrt{5x+7} \, dx$. Oops! The integrand isn't a product. Are we losers? No way. The integrand is $\int 1 \times \sqrt{5x+7} \, dx$.

The square root has a known antiderivative, so choose $u = 5x + 7$. Then $\frac{du}{dx} = 5$ matches the remaining factor 1 in the integrand up to a multiplicative factor. **We win.**

$$[u = 5x + 7] = [du = 5dx] = \left[dx = \frac{1}{5} du \right].$$

So

$$\int \sqrt{5x+7} \, dx = \int \frac{1}{5} \sqrt{u} \, du = \frac{1}{5} \times \frac{2}{3} u^{3/2} = \frac{2}{15} (5x+7)^{3/2}.$$

Fake matches

Let's try the problem

$$\int x \exp(x^4) dx.$$

Since \exp has a known antiderivative, for u choose what is inside \exp . Thus

$$[u = x^4] = [du = 4x^3 dx] = \left[dx = \frac{1}{4x^3} du \right]$$

So far, Okie-dokie.

$$\int x \exp(x^4) dx = \int x \exp(u) \frac{1}{4x^3} du = \int \exp(u) \frac{1}{4x^2} du$$

- ❗ Since $\int \exp(u) \frac{1}{4x^2} du$ depends on both the old variable x and the new variable u , we **have not completed our task of out with the old and in with the new.**
- ❗ We could push on this a bit using $x = u^{1/4}$ to eliminate the remaining terms involving x . This gives

$$\int \frac{1}{4\sqrt{u}} \exp(u) du.$$

Occult (that is hidden) matches

$$\int \frac{x}{\sqrt{1-x^4}} dx$$

- ✓ We know the antiderivative of reciprocal square root. That suggests

$$u = 1 - x^4$$

But then

$$du = -4x^3 dx.$$

- ✓ The substitution yields

$$\int \frac{x}{\sqrt{1-x^4}} dx = \int -\frac{x}{\sqrt{u}} \frac{du}{4x^3} = \int -\frac{1}{\sqrt{u}} \frac{du}{4x^2}$$

- ✓ As yet, we have a yucky mixture of old and new. This doesn't look promising. Should we bail out, or try some thing new?

The rule of holes ¹

- ✓ Let's bailout.
- ✓ We also know the antiderivative of $\int \frac{1}{\sqrt{1-x^2}} dx$.
- ✓ This suggests choosing $1 - x^4 = 1 - u^2$; equivalently $x^2 = u$. Then

$$du = 2x dx.$$

- ✓ So

$$\int \frac{x}{\sqrt{1-x^4}} dx = \int \frac{1}{2} \frac{1}{\sqrt{1-u^2}} du = \frac{1}{2} \sin^{-1}(u) = \frac{1}{2} \sin^{-1}(x^2)$$

¹When you are in a hole, digging faster isn't the answer.