

NAMED SETS

empty set	\emptyset
real numbers	\mathbf{R}
ordered pairs	\mathbf{R}^2
integers	\mathbf{Z}
positive integers	$\mathbf{Z}_{>0}$
positive real numbers	$\mathbf{R}_{>0}$

EXPONENTS

For $a, b \in \mathbf{R}_{>0}$, $x \in \mathbf{R}$, and $m, n \in \mathbf{R}$,

$$\begin{aligned} a^0 &= 1, & 0^a &= 0 \\ 1^a &= 1, & a^n a^m &= a^{n+m} \\ a^n / a^m &= a^{n-m}, & (a^n)^m &= a^{n \cdot m} \\ a^{-m} &= 1/a^m, & (a/b)^m &= a^m / b^m \\ \sqrt{x^2} &= |x| \end{aligned}$$

TRIGONOMETRIC IDENTITIES

$$\begin{aligned} (\cos(x))^2 + (\sin(x))^2 &= 1 \\ 2(\cos(x))^2 &= 1 + \cos(2x) \\ 2(\sin(x))^2 &= 1 - \cos(2x) \\ (\cos(x))^2 - (\sin(x))^2 &= \cos(2x) \\ \sin(x+y) &= \sin(x)\cos(y) + \cos(x)\sin(y) \\ \cos(x+y) &= \cos(x)\cos(y) - \sin(x)\sin(y) \\ \operatorname{arccot}(x) &= \pi/2 - \arctan(x) \quad \operatorname{dom}(\operatorname{arccot}) = (0, \pi) \\ \operatorname{arccsc}(x) &= \arcsin(1/x) \\ \operatorname{arcsec}(x) &= \arccos(1/x) \\ \arcsin(x) + \arccos(x) &= \pi/2 \\ \operatorname{arcsec}(x) + \operatorname{arccsc}(x) &= \pi/2 \end{aligned}$$

HYPERBOLIC FUNCTIONS

$$\begin{aligned} 2 \cosh(x) &= \exp(x) + \exp(-x) \\ 2 \sinh(x) &= \exp(x) - \exp(-x) \\ \tanh(x) &= \cosh(x)/\sinh(x) \\ \cosh(x)^2 - \sinh(x)^2 &= 1 \end{aligned}$$

LOGARITHMS

$$\log_a(x) = \frac{\ln(x)}{\ln(a)}$$

DERIVATIVES

Specific cases

$F(x)$	$F'(x)$
$\cos(x)$	$-\sin(x)$
$\sin(x)$	$\cos(x)$
$\tan(x)$	$\sec(x)^2$
$\sec(x)$	$\sec(x)\tan(x)$
$\csc(x)$	$-\cot(x)\csc(x)$
$\cot(x)$	$-\csc(x)^2$
$\arccos(x)$	$-1/\sqrt{1-x^2}$
$\arcsin(x)$	$1/\sqrt{1-x^2}$
$\arctan(x)$	$1/(x^2+1)$
$\cosh(x)$	$\sinh(x)$
$\sinh(x)$	$\cosh(x)$
$\tanh(x)$	$1/\cosh(x)^2$
$\operatorname{arccosh}(x)$	$1/\sqrt{x^2-1}$
$\operatorname{arsinh}(x)$	$1/\sqrt{1+x^2}$
$\operatorname{arctanh}(x)$	$1/(1-x^2)$
$\exp(x)$	$\exp(x)$
$\ln(x)$	$1/x$

General Cases

$F(x)$	$F'(x)$
$af(x) + bg(x)$	$af'(x) + bg'(x)$
$f(x)g(x)$	$f'(x)g(x) + f(x)g'(x)$
$1/g(x)$	$-g'(x)/g(x)^2$
$f(x)/g(x)$	$(g(x)f'(x) - f(x)g'(x))/g(x)^2$
$f(g(x))$	$g'(x)f'(g(x))$
$f^{-1}(x)$	$1/f'(f^{-1}(x))$

ANTIDERIVATIVES

$$\begin{aligned} \int a \, dx &= ax \\ \int x^a \, dx &= \frac{1}{1+a} x^{a+1}, \quad \text{if } a \neq -1 \\ \int \frac{1}{x} \, dx &= \ln|x| \\ \int \cos(x) \, dx &= \sin(x) \\ \int \sin(x) \, dx &= -\cos(x) \\ \int \tan(x) \, dx &= \ln|\sec(x)| \\ \int \sec(x) \, dx &= \ln|\tan(x) + \sec(x)| \\ \int \csc(x) \, dx &= -\ln|\csc(x) + \cot(x)| \\ \int \cot(x) \, dx &= \ln|\sin(x)| \\ \int |x| \, dx &= x|x|/2 \end{aligned}$$

SUMS

For $n \in \mathbf{Z}_{>0}$

$$\begin{aligned} \sum_{k=0}^{n-1} 1 &= n \\ \sum_{k=0}^{n-1} k &= \frac{(n-1)n}{2} \\ \sum_{k=0}^{n-1} k^2 &= \frac{(n-1)n(2n-1)}{6} \\ \sum_{k=0}^{n-1} x^k &= \frac{1-x^n}{1-x}, \quad x \neq 1 \\ \sum_{k=0}^{\infty} z^k &= \begin{cases} \frac{1}{1-z} & z \in (-1, 1) \\ \infty & z \in [1, \infty] \end{cases} \end{aligned}$$

When $z \in (-\infty, -1]$, the series $\sum_{k=0}^{\infty} z^k$ diverges.

APPLICATIONS

Arclength of curve $y = f(x)$ with $a \leq x \leq b$

$$= \int_a^b \sqrt{1 + f'(x)^2} \, dx$$

For the region Q of the xy plane given by

$$Q = \{(x, y) \mid f(x) \leq y \leq g(x) \wedge a \leq x \leq b\},$$

we have

$$\operatorname{Area}(Q) = \int_a^b g(x) - f(x) \, dx$$

Assuming $0 \leq f(x)$ and rotating about the x -axis

$$\operatorname{Vol}(Q) = \pi \int_a^b g(x)^2 - f(x)^2 \, dx$$

Assuming $0 \leq a < b$ and rotating about the y -axis

$$\operatorname{Vol}(Q) = 2\pi \int_a^b x(g(x) - f(x)) \, dx$$

Centroid

$$\operatorname{Area}(Q) \times \bar{x} = \int_a^b x(g(x) - f(x)) \, dx$$

$$\operatorname{Area}(Q) \times \bar{y} = \frac{1}{2} \int_a^b (g(x)^2 - f(x)^2) \, dx$$


For the region Q of the xy plane given by

$$Q = \{(x, y) \mid f(y) \leq x \leq g(y) \wedge a \leq y \leq b\},$$


interchange x and y in *all* the previous formulas.

FAMOUS TRIANGLES


The 30-60-90 triangle



The 45-45-90 triangle



LAWS OF COSINE & SINE




Law of cosine: $C^2 = A^2 + B^2 - 2AB \cos(\gamma)$

Law of sines: $\frac{\sin(\alpha)}{A} = \frac{\sin(\beta)}{B} = \frac{\sin(\gamma)}{C}$

Area: $\operatorname{Area} = 1/2 h B = 1/2 AB \sin(\gamma)$

VOLUMES

Right Circular Cylinder



Volume: $V = \pi r^2 h$


Area: (not including circular ends) $A = 2\pi r h$

Sphere

Area:
 $A = 4\pi r^2$

Volume:
 $V = \frac{4\pi}{3} r^3$

Cone



$$\text{Volume: } V = \frac{1}{3}\pi r^2 h$$

Area (not including circular base)

$$A = \pi r \sqrt{r^2 + h^2}$$

P-SERIES, DIVERGENCE TEST, RATIO TEST, COMPARISON, & AST

The series $\sum_{k=1}^{\infty} \frac{1}{k^p}$ converges when $p \in (1, \infty)$; otherwise it diverges.

If $\lim_{k \rightarrow \infty} a_k \neq 0$, the series $\sum a_k$ diverges.

Let a be a sequence with $0 \notin \text{range}(a)$. Define $L = \lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right|$.

- $L \in [0, 1) \Rightarrow \sum |a_k|$ converges.
- $L \in (1, \infty) \Rightarrow \sum a_k$ diverges.

Let a and b be positive sequences. Define $L = \lim_{k \rightarrow \infty} \frac{a_k}{b_k}$.

- If $L \in \mathbf{R}_{>0}$ and $\sum a_k$ converges then $\sum b_k$ converges.
- If $L \in \mathbf{R}_{>0}$ and $\sum a_k$ diverges then $\sum b_k$ diverges.
- If $L = 0$ and $\sum b_k$ converges, then $\sum a_k$ converges.
- If $L = \infty$ and $\sum b_k$ diverges, then $\sum a_k$ diverges.

Let a be a positive and eventually decreasing sequence. Then $\sum (-1)^k a_k$ converges iff $\lim_{k \rightarrow \infty} a_k = 0$.

TAYLOR AND MACLAURIN SERIES

If a function F is infinitely differentiable at a , its Taylor series centered at a is

$$\sum_{k=0}^{\infty} \frac{F^{(k)}(a)}{k!} (x-a)^k.$$

When a is zero, the Taylor series is also known as the MacLaurin Series.

POLAR TO CARTESIAN

$$y = r \sin(\theta)$$

For $r > 0$ and $0 \leq \theta < 2\pi$

$$r = \sqrt{x^2 + y^2}$$

$$\theta = \begin{cases} 2\pi - \arccos(x/r) & \text{if } y < 0 \\ \arccos(x/r) & \text{if } y \geq 0 \end{cases}$$

INTEGRATE POWERS OF TRIG

Let $m, n \in \mathbf{Z}_{\geq 0}$. Then


- $\int \cos(x)^{2m} \sin(x)^{2n} dx = \int \left(\frac{1 + \cos(2x)}{2} \right)^m \left(\frac{1 - \cos(2x)}{2} \right)^n dx$
- $\int \cos(x)^{2m+1} \sin(x)^n dx = \int (1 - z^2)^m z^n dz$, where $z = \sin(x)$
- $\int \cos(x)^m \sin(x)^{2n+1} dx = \int z^m (1 - z^2)^n dz$, where $z = \cos(x)$

- $\int \sec(x)^n dx = \frac{1}{n-1} \sec(x)^{n-2} \tan(x) + \frac{n-2}{n-1} \int \sec(x)^{n-2} dx$, provided $n \neq 1$.
- $\int \tan(x)^{2m+1} \sec(x)^n dx = \int (z^2 - 1)^m z^{n-1} dz$, where $z = \sec(x)$
- $\int \tan(x)^{2m} \sec(x)^n dx = \int (\sec(x)^2 - 1)^m \sec(x)^n dx$.

TRIG SUBSTITUTIONS

- $\int F(x, (1-x^2)^{n/2}) dx$, use $x = \sin(\theta)$, where $\theta \in [-\pi/2, \pi/2]$, then integrate $\int F(\sin(\theta), \cos(\theta)^n) \cos(\theta) d\theta$
- $\int F(x, (1+x^2)^{n/2}) dx$, use $x = \sinh(\theta)$, where $\theta \in \mathbf{R}$, then integrate $\int F(\sinh(\theta), \cosh(\theta)^n) \cosh(\theta) d\theta$
- $\int F(x, (x^2-1)^{n/2}) dx$, use $x = \sec(\theta)$, then integrate $\int F(\sec(\theta), \tan(\theta)^n) \sec(\theta), \tan(\theta) d\theta$

UNIT CIRCLE



Revised September 28, 2023. Barton Willis is the author of this work. This work is licensed under Attribution 4.0 International (CC BY 4.0). For the current version of this document, visit 