

In class work **8** has questions **1** through **4** with a total of **15** points. Turn in your work at the end of class *on paper*. This assignment is due *Thursday 21 September 13:20*.

**Notice:**  $\cos(x)^2$  means  $(\cos(x))^2$ . It *doesn't* mean  $\cos(x^2)$ . Our textbook writes this expression as  $\cos^2(x)$ . Either way is OK.


- 2 1. Find the area of the region  $\{(x, y) | 0 \leq y \leq \sin(x)^2 \text{ and } 0 \leq x \leq \pi\}$ .

**Solution:**

$$\begin{aligned}
 \text{Area} &= \int_0^{\pi} \sin(x)^2 dx, && \text{(area formula)} \\
 &= \int_0^{\pi} \frac{1}{2} - \frac{1}{2} \cos(2x) dx, && \text{(double angle)} \\
 &= \frac{1}{2}x - \frac{1}{4} \sin(2x) \Big|_0^{\pi}, && \text{(known antiderivatives)} \\
 &= \frac{\pi}{2}. && \text{(algebra)}
 \end{aligned}$$

- 2 2. Use Desmos to graph  $y = \cos(x)^3$  on the interval  $[0, \pi]$ . Based on the graph, make a pretty good guess for the numerical value of  $\int_0^{\pi} \cos(x)^3 dx$ . Duplicate the graph here and justify your guess.

**Solution:**



The area below the x-axis appears to equal the area above the x-axis. My guess is that  $\int_0^\pi \cos(x)^3 dx = 0$ .

- 2 3. Find the numerical value of  $\int_0^\pi \cos(x)^3 dx$ .

**Solution:** Let's first find an antiderivative

$$\begin{aligned}\int \cos(x)^3 dx &= \int \cos(x)(1 - \sin(x)^2) dx, \\ &= \int 1 - z^2 dz, & (z = \sin(x) \text{ and } dz = \cos(x)dx) \\ &= z - \frac{1}{3}z^3, \\ &= \sin(x) - \frac{1}{3}\sin(x)^3\end{aligned}$$

So  $\int_0^\pi \cos(x)^3 dx = 0$ .

3 4. Use the identities

$$\begin{aligned}\sin(x) \cos(y) &= \frac{\sin(y+x) - \sin(y-x)}{2}, \\ \sin(x) \sin(y) &= -\frac{\cos(y+x) - \cos(y-x)}{2}, \\ \cos(x) \cos(y) &= \frac{\cos(y+x) + \cos(y-x)}{2}.\end{aligned}$$

to find the values of each of the following definite integrals

2 (a)  $\int_0^{2\pi} \sin(5x) \cos(x) \, dx$ .

**Solution:**

$$\int_0^{2\pi} \sin(5x) \cos(x) \, dx = \int_0^{2\pi} \frac{\sin(6x) + \sin(4x)}{2} \, dx = -\frac{\cos(6x)}{12} - \frac{\cos(4x)}{8} \Big|_0^{2\pi} = 0.$$

Actually for any integer  $n$ , we have  $\int_0^{2\pi} \sin(nx) \, dx = 0$ .

2 (b)  $\int_0^{2\pi} \cos(5x) \cos(x) \, dx$ .

**Solution:** Let's use the nice fact that for any nonzero integer  $n$ , we have  $\int_0^{2\pi} \cos(nx) \, dx = 0$

$$\int_0^{2\pi} \cos(5x) \cos(x) \, dx = \int_0^{2\pi} \frac{\cos(6x) + \cos(4x)}{2} \, dx = 0.$$

2 (c)  $\int_0^{2\pi} \cos(5x)^2 \, dx$ .

**Solution:** Again, use the nice fact that for any nonzero integer  $n$ , we have  $\int_0^{2\pi} \cos(nx) \, dx = 0$ .

$$\int_0^{2\pi} \cos(5x)^2 \, dx = \int_0^{2\pi} \frac{1}{2} + \frac{1}{2} \cos(10x) \, dx = \pi.$$