

CS 302 Recitation 3

November 1, 2020

Problem 1

- Simplify the given regular expression heuristically.

Problem 1

- Simplify the given regular expression heuristically.

-

$$(aa^*(aa^*b + b) + b + aa^*b + \varepsilon)(aa^*(aa^*b + b))^*$$

Problem 1

- Simplify the given regular expression heuristically.

-

$$(aa^*(aa^*b + b) + b + aa^*b + \varepsilon)(aa^*(aa^*b + b))^*$$

-

$$P = aa^*$$

Problem 1

- Simplify the given regular expression heuristically.

$$(aa^*(aa^*b + b) + b + aa^*b + \varepsilon)(aa^*(aa^*b + b))^*$$

$$P = aa^*$$

$$(P(Pb + b) + \varepsilon + Pb + b)(P(Pb + b))^*$$

Problem 1

- Simplify the given regular expression heuristically.

$$(aa^*(aa^*b + b) + b + aa^*b + \varepsilon)(aa^*(aa^*b + b))^*$$

$$P = aa^*$$

$$(P(Pb + b) + \varepsilon + Pb + b)(P(Pb + b))^*$$

$$(P(Pb + b) + \varepsilon + Pb + b)(P(Pb + b))^*$$

Problem 1

- Simplify the given regular expression heuristically.

$$(aa^*(aa^*b + b) + b + aa^*b + \varepsilon)(aa^*(aa^*b + b))^*$$

$$P = aa^*$$

$$(P(Pb + b) + \varepsilon + Pb + b)(P(Pb + b))^*$$

$$(P(Pb + b) + \varepsilon + Pb + b)(P(Pb + b))^*$$

- Distributive Law

$$(P(Pb + b) + \varepsilon)(P(Pb + b))^* + (Pb + b)(P(Pb + b))^*$$

Problem 1 cont.

- $(P(Pb + b) + \varepsilon)(P(Pb + b))^* + (Pb + b)(P(Pb + b))^*$

Problem 1 cont.



$$(P(Pb + b) + \epsilon)(P(Pb + b))^* + (Pb + b)(P(Pb + b))^*$$

- By $(\epsilon + L)L^* = L^*$,

$$(P(Pb + b))^* + (Pb + b)(P(Pb + b))^*$$

Problem 1 cont.



$$(P(Pb + b) + \varepsilon)(P(Pb + b))^* + (Pb + b)(P(Pb + b))^*$$

- By $(\varepsilon + L)L^* = L^*$,

$$(P(Pb + b))^* + (Pb + b)(P(Pb + b))^*$$

- Since $Pb + b = aa^*b + b = (aa^* + \varepsilon)b = a^*b$,

$$(aa^*a^*b)^* + a^*b(aa^*a^*b)^*$$

Problem 1 cont.

•

$$(P(Pb + b) + \varepsilon)(P(Pb + b))^* + (Pb + b)(P(Pb + b))^*$$

• By $(\varepsilon + L)L^* = L^*$,

$$(P(Pb + b))^* + (Pb + b)(P(Pb + b))^*$$

• Since $Pb + b = aa^*b + b = (aa^* + \varepsilon)b = a^*b$,

$$(aa^*a^*b)^* + a^*b(aa^*a^*b)^*$$

• $a^*a^* = a^*$

$$(aa^*b)^* + a^*b(aa^*b)^*$$

Problem 1 cont. (2)



$$(a\textcolor{red}{a}^*b)^* + a^*b(a\textcolor{red}{a}^*b)^*$$

Problem 1 cont. (2)



$$(a\textcolor{red}{a}^*b)^* + a^*b(a\textcolor{red}{a}^*b)^*$$

- Distributive Law

$$(a^*b + \varepsilon)(aa^*b)^*$$

Problem 1 cont. (2)



$$(a\textcolor{red}{a}^*b)^* + a^*b(a\textcolor{red}{a}^*b)^*$$

- Distributive Law

$$(a^*b + \varepsilon)(aa^*b)^*$$

- Distributive Law

$$((\textcolor{red}{aa}^* + \varepsilon)b + \varepsilon)(aa^*b)^*$$

Problem 1 cont. (2)



$$(a\textcolor{red}{a}^*b)^* + a^*b(a\textcolor{red}{a}^*b)^*$$

- Distributive Law

$$(a^*b + \varepsilon)(aa^*b)^*$$

- Distributive Law

$$((\textcolor{red}{a}a^* + \varepsilon)b + \varepsilon)(aa^*b)^*$$

- Distributive Law

$$(\textcolor{red}{a}a^*\textcolor{red}{b} + \textcolor{red}{b} + \varepsilon)(aa^*b)^*$$

Problem 1 cont. (2)



$$(a\textcolor{red}{a}^*b)^* + a^*b(a\textcolor{red}{a}^*b)^*$$

- Distributive Law

$$(a^*b + \varepsilon)(a\textcolor{red}{a}^*b)^*$$

- Distributive Law

$$((\textcolor{red}{a}\textcolor{red}{a}^* + \varepsilon)b + \varepsilon)(a\textcolor{red}{a}^*b)^*$$

- Distributive Law

$$(\textcolor{red}{a}\textcolor{red}{a}^*b + \textcolor{red}{b} + \varepsilon)(a\textcolor{red}{a}^*b)^*$$

- By $(\varepsilon + L)L^* = L^*$,

$$(\textcolor{red}{b} + \varepsilon)(a\textcolor{red}{a}^*b)^*$$

Problem 2

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

1) $x.y^i.z \in A, i = 0, 1, 2, \dots$

2) $|y| > 0$,

3) $|x.y| \leq p$

- Prove that language $A_1 = \{0^n 1^m 0^n \mid m, n \geq 0\}$ is regular or not.

Problem 2

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Prove that language $A_1 = \{0^n 1^m 0^n \mid m, n \geq 0\}$ is regular or not.

•

$$s = x.y.z = 0^p 10^p$$

Problem 2

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Prove that language $A_1 = \{0^n 1^m 0^n \mid m, n \geq 0\}$ is regular or not.

•

$$s = x.y.z = 0^p 10^p$$

- By condition 3, x consists of all 0's, and y consists of all 0's.

$$x = 0^a, y = 0^b, z = 0^c 10^p$$

Problem 2

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Prove that language $A_1 = \{0^n 1^m 0^n \mid m, n \geq 0\}$ is regular or not.

•

$$s = x.y.z = 0^p 10^p$$

- By condition 3, x consists of all 0's, and y consists of all 0's.

$$x = 0^a, y = 0^b, z = 0^c 10^p$$

- For $x.y.z \in A_1$, $x.y.y.z = 0^{a+2b+c} 10^p$

Problem 2

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Prove that language $A_1 = \{0^n 1^m 0^n \mid m, n \geq 0\}$ is regular or not.

•

$$s = x.y.z = 0^p 10^p$$

- By condition 3, x consists of all 0's, and y consists of all 0's.

$$x = 0^a, y = 0^b, z = 0^c 10^p$$

- For $x.y.z \in A_1$, $x.y.y.z = 0^{a+2b+c} 10^p$
- Contradicts with Condition 1!

Problem 3

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Prove that language $A_2 = \{a^{2^n} \mid n \geq 0\}$ is regular or not.

Problem 3

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Prove that language $A_2 = \{a^{2^n} \mid n \geq 0\}$ is regular or not.

•

$$s = x.y.z = a^{2^p}$$

Problem 3

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Prove that language $A_2 = \{a^{2^n} \mid n \geq 0\}$ is regular or not.

•

$$s = x.y.z = a^{2^p}$$

- By definition of the language, the string from language A_2 with minimum length greater than s , is $a^{2^{p+1}}$

$$|a^{2^{p+1}}| = |a^{2 \cdot 2^p}| = 2 \cdot |s|$$

Problem 3

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Prove that language $A_2 = \{a^{2^n} \mid n \geq 0\}$ is regular or not.

•

$$s = x.y.z = a^{2^p}$$

- By definition of the language, the string from language A_2 with minimum length greater than s , is $a^{2^{p+1}}$

$$|a^{2^{p+1}}| = |a^{2 \cdot 2^p}| = 2 \cdot |s|$$

- Since $x.y.z = a^{2^{p+1}} \in A_2$, the statement $x.y.y.z \in A_2$ must also be true, by condition 1. Contradiction! Since $|x.y| \leq p, |y| > 0$,

Problem 3

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Prove that language $A_2 = \{a^{2^n} \mid n \geq 0\}$ is regular or not.

•

$$s = x.y.z = a^{2^p}$$

- By definition of the language, the string from language A_2 with minimum length greater than s , is $a^{2^{p+1}}$

$$|a^{2^{p+1}}| = |a^{2 \cdot 2^p}| = 2 \cdot |s|$$

- Since $x.y.z = a^{2^p} \in A_2$, the statement $x.y.y.z \in A_2$ must also be true, by condition 1. Contradiction! Since $|x.y| \leq p, |y| > 0$,
- $x.y.y.z$ can be neither a^{2^p} nor $a^{2^{p+1}}$

Problem 4

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

① $1) x.y^i.z \in A, i = 0, 1, 2, \dots$

② $2) |y| > 0,$

③ $3) |x.y| \leq p$

- Prove that language $A_3 = \{0^m 1^n \mid m \neq n \text{ and } m, n \geq 0\}$ is regular or not.

Problem 4

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

① 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$

② 2) $|y| > 0$,

③ 3) $|x.y| \leq p$

- Prove that language $A_3 = \{0^m 1^n \mid m \neq n \text{ and } m, n \geq 0\}$ is regular or not.
- The basic idea:

$$s = x.y.z = 0^p 1^d, d > p$$

Problem 4

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Prove that language $A_3 = \{0^m 1^n \mid m \neq n \text{ and } m, n \geq 0\}$ is regular or not.
- The basic idea:

$$s = x.y.z = 0^p 1^d, d > p$$

- By condition 3, x consists of all 0's, and y consists of all 0's.

$$x = 0^a, y = 0^b, z = 0^c 1^d, a + b + c = p$$

Problem 4

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Prove that language $A_3 = \{0^m 1^n \mid m \neq n \text{ and } m, n \geq 0\}$ is regular or not.
- The basic idea:

$$s = x.y.z = 0^p 1^d, d > p$$

- By condition 3, x consists of all 0's, and y consists of all 0's.

$$x = 0^a, y = 0^b, z = 0^c 1^d, a + b + c = p$$

- Find a $x.y^i.z = 0^{a+bi+c} 1^d$ s.t. $a + bi + c = d$ and prove by contradiction.

Problem 4 cont.

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Find a $x.y^i.z = 0^{a+bi+c}1^d$ s.t. $a + bi + c = d$

Problem 4 cont.

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Find a $x.y^i.z = 0^{a+bi+c}1^d$ s.t. $a + bi + c = d$
- Problem:

$$bi = d - a - c$$

Problem 4 cont.

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Find a $x.y^i.z = 0^{a+bi+c}1^d$ s.t. $a + bi + c = d$
- Problem:

$$bi = d - a - c$$

- We must choose such a d that it satisfies this equation for each a, b, c .

Problem 4 cont.

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Find a $x.y^i.z = 0^{a+bi+c}1^d$ s.t. $a + bi + c = d$
- Problem:

$$bi = d - a - c$$

- We must choose such a d that it satisfies this equation for each a, b, c .

•

$$d = p + p!$$

$$x = 0^a, y = 0^b, z = 0^c 1^{p+p!}, a + b + c = p$$

Problem 4 cont.

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Find a $x.y^i.z = 0^{a+bi+c}1^d$ s.t. $a + bi + c = d$
- Problem:

$$bi = d - a - c$$

- We must choose such a d that it satisfies this equation for each a, b, c .
-

$$d = p + p!$$

$$x = 0^a, y = 0^b, z = 0^c 1^{p+p!}, a + b + c = p$$

- For $i = \frac{p!}{b} + 1$:

$$x.y^{\frac{p!}{b}+1}.z = 0^a 0^{p!+b} 0^c 1^{p!} = 0^{a+b+c+p!} 1^{p+p!} = 0^{p+p!} 1^{p+p!}$$

Problem 4 altn.

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

① $1) x.y^i.z \in A, i = 0, 1, 2, \dots$

② $2) |y| > 0,$

③ $3) |x.y| \leq p$

- Let C be the complement of language A_3 . By closure property, if A_3 is regular, C is also regular.

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- ❶ 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- ❷ 2) $|y| > 0$,
- ❸ 3) $|x.y| \leq p$

- Let C be the complement of language A_3 . By closure property, if A_3 is regular, C is also regular.

-

$$B = \{0^n 1^n \mid n \geq 0\} = C \cap 0^* 1^*$$

By closure property, if C and $0^* 1^*$ are regular, B is also regular.

Problem 4 altn.

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- 2) $|y| > 0$,
- 3) $|x.y| \leq p$

- Let C be the complement of language A_3 . By closure property, if A_3 is regular, C is also regular.

-

$$B = \{0^n 1^n \mid n \geq 0\} = C \cap 0^* 1^*$$

By closure property, if C and $0^* 1^*$ are regular, B is also regular.

- B is not regular $\wedge 0^* 1^*$ is regular $\Rightarrow C$ is not regular.

Problem 4 altn.

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

- ❶ 1) $x.y^i.z \in A, i = 0, 1, 2, \dots$
- ❷ 2) $|y| > 0$,
- ❸ 3) $|x.y| \leq p$

- Let C be the complement of language A_3 . By closure property, if A_3 is regular, C is also regular.

•

$$B = \{0^n 1^n \mid n \geq 0\} = C \cap 0^* 1^*$$

By closure property, if C and $0^* 1^*$ are regular, B is also regular.

- B is not regular $\wedge 0^* 1^*$ is regular $\Rightarrow C$ is not regular.
- We proved A_3 is not regular.

Problem 5

Pumping Lemma

For pumping length p and $s = x.y.z \in A$,

① $1) x.y^i.z \in A, i = 0, 1, 2, \dots$

② $2) |y| > 0,$

③ $3) |x.y| \leq p$

- Prove that language $A_4 = \{w \mid w \text{ has an equal number of occurrences of } 01\text{'s and } 10\text{'s as substrings}\}$ is regular or not.

Problem 5 cont.

- It is regular!

Problem 5 cont.

- It is regular!

